

DRAFT

KEY HABITATS

Oregon's State Wildlife Action Plan

3	Contents	
4	KEY HABITATS	9
5	Strategy Habitat Methodology	9
6	ASPEN WOODLANDS	10
7	ECOREGIONS	10
8	CHARACTERISTICS.....	10
9	CONSERVATION OVERVIEW	10
10	[Spotlight] Beaver Habitat and Beaver Modified Habitat	11
11	LIMITING FACTORS AND RECOMMENDED APPROACHES.....	12
12	Limiting Factor: Altered Fire Regimes	12
13	Limiting Factor: Overgrazing.....	12
14	Limiting Factor: Invasive Species.....	13
15	Limiting Factor: Drought.....	13
16	RESOURCES FOR MORE INFORMATION	14
17	COASTAL DUNES	15
18	ECOREGIONS	15
19	CHARACTERISTICS.....	15
20	CONSERVATION OVERVIEW	15
21	LIMITING FACTORS AND RECOMMENDED APPROACHES.....	16
22	Limiting Factor: European Beachgrass and other Invasive Plants	16
23	Limiting Factor: Development	16
24	Limiting Factor: Recreational Impacts.....	16
25	HABITAT CHANGE TRENDS ANALYSIS	16
26	RESOURCES FOR MORE INFORMATION.....	17
27	REFERENCES.....	17
28	ESTUARIES	18
29	ECOREGIONS	18
30	CHARACTERISTICS.....	18
31	Physical Environment	19

32	Biological Community	23
33	CONSERVATION OVERVIEW.....	24
34	LIMITING FACTORS AND RECOMMENDED APPROACHES	25
35	Limiting Factor: Increased Shoreline Development, Land Use Conversion, and Altered or	
36	Blocked Tidal Flow	25
37	Limiting Factor: Alteration of Freshwater Inputs into Estuaries.....	25
38	Limiting Factor: Degraded Water Quality	26
39	Limiting Factor: Non-Native and Invasive Species.....	26
40	Limiting Factor: Management and Planning Needs	28
41	Limiting Factor: Loss of Habitat Complexity	28
42	Limiting Factor: Climate Change	29
43	Limiting Factor: Oil Spills and Hazardous wastes.....	30
44	RESOURCES FOR MORE INFORMATION	30
45	REFERENCES.....	31
46	FLOWING WATER AND RIPARIAN HABITATS	33
47	ECOREGIONS	33
48	CHARACTERISTICS.....	33
49	Flowing Water Habitats.....	33
50	Riparian Habitats	34
51	CONSERVATION OVERVIEW	35
52	[Spotlight] Beaver Habitat and Beaver-Modified Habitat	36
53	Oregon Planning and Regulatory Background for Flowing Waters.....	36
54	Oregon’s Planning and Regulatory Background for Riparian Habitats	37
55	LIMITING FACTORS AND RECOMMENDED APPROACHES	37
56	Limiting Factor: Water Quantity.....	37
57	Limiting Factor: Invasive Aquatic Species	38
58	Limiting Factor: Passage Barriers and Channel Complexity.....	39
59	Limiting Factor: Pollution.....	39
60	Limiting Factor: Water Temperature.....	40
61	Limiting Factor: Sedimentation	41

62	Limiting Factor: Loss of Riparian Habitat, Floodplain Function, and Habitat Complexity..	42
63	Limiting Factor: Riparian Habitat Degradation	43
64	Limiting Factor: Invasive Plants in Riparian Habitat	43
65	RESOURCES FOR MORE INFORMATION	44
66	GRASSLANDS	45
67	ECOREGIONS	45
68	CHARACTERISTICS.....	45
69	ECOREGIONAL CHARACTERISTICS	45
70	Blue Mountains.....	45
71	Coast Range.....	45
72	Columbia Plateau	46
73	Klamath Mountains	46
74	West Cascades.....	46
75	Willamette Valley	47
76	CONSERVATION OVERVIEW	47
77	LIMITING FACTORS AND RECOMMENDED APPROACHES.....	48
78	Limiting Factor: Altered Fire Regimes	48
79	Limiting Factor: Invasive Species.....	48
80	Limiting Factor: Land Use Conversion	49
81	Limiting Factor: Land Management Conflicts	50
82	Limiting Factor: Loss of Oak Savannas	50
83	HABITAT CHANGE TRENDS ANALYSIS	51
84	RESOURCES FOR MORE INFORMATION	51
85	REFERENCES.....	52
86	LATE SUCCESSIONAL MIXED CONIFER FORESTS	53
87	ECOREGIONS	53
88	CHARACTERISTICS.....	53
89	ECOREGIONAL CHARACTERISTICS	53
90	Blue Mountains.....	53

91	Coast Range	53
92	East Cascades.....	54
93	Klamath Mountains	54
94	West Cascades.....	54
95	CONSERVATION OVERVIEW	55
96	[Spotlight] Pileated Woodpecker	56
97	LIMITING FACTORS AND RECOMMENDED APPROACHES	57
98	Limiting Factor: Loss of Structural Habitat Elements	57
99	Limiting Factor: Loss of Late Successional Stand Size and Connectivity.....	57
100	HABITAT CHANGE TRENDS ANALYSIS	58
101	RESOURCES FOR MORE INFORMATION	58
102	REFERENCES.....	58
103	NATURAL LAKES	60
104	ECOREGIONS	60
105	CHARACTERISTICS.....	60
106	CONSERVATION OVERVIEW.....	61
107	LIMITING FACTORS AND RECOMMENDED APPROACHES	61
108	Limiting Factor: Drought	61
109	Limiting Factor: Water Quality	62
110	Limiting Factor: Habitat Loss.....	63
111	Limiting Factor: Invasive species	63
112	Limiting Factor: Wildfires	64
113	RESOURCES FOR MORE INFORMATION.....	65
114	OAK HABITATS	67
115	ECOREGIONS	67
116	CHARACTERISTICS.....	67
117	ECOREGIONAL CHARACTERISTICS	67
118	Coast Range	67
119	East Cascades.....	68

120	Klamath Mountains	68
121	West Cascades	68
122	Willamette Valley	68
123	CONSERVATION OVERVIEW	69
124	[SPOTLIGHT] Acorn Woodpeckers	70
125	LIMITING FACTORS AND RECOMMENDED APPROACHES	70
126	Limiting Factor: Fire Suppression and Fir Encroachment	70
127	Limiting Factor: Land Use Conversion and Continued Habitat Loss	71
128	Limiting Factor: Loss of Habitat Structure	72
129	Limiting Factor: Invasive Species and Diseases	72
130	Limiting Factor: Climate Change	73
131	HABITAT TRENDS ANALYSIS	73
132	RESOURCES FOR MORE INFORMATION	73
133	REFERENCES	74
134	PONDEROSA PINE WOODLANDS	75
135	ECOREGIONS	75
136	CHARACTERISTICS	75
137	ECOREGIONAL CHARACTERISTICS	75
138	Blue Mountains	75
139	East Cascades	76
140	Klamath Mountains	76
141	CONSERVATION OVERVIEW	76
142	LIMITING FACTORS AND RECOMMENDED APPROACHES	77
143	Limiting Factor: Altered Fire Regimes and Addressing Risk of Uncharacteristically Severe	
144	Wildfire	77
145	Limiting Factor: Loss of Size and Connectivity of Large-structure Ponderosa Pine Habitats	
146	78
147	Limiting Factor: Invasive Species	79
148	RESOURCES FOR MORE INFORMATION	79
149	SAGEBRUSH HABITATS	80

150	ECOREGIONS	80
151	CHARACTERISTICS	80
152	ECOREGIONAL CHARACTERISTICS	80
153	Blue Mountains.....	80
154	Columbia Plateau	81
155	East Cascades	81
156	Northern Basin and Range	81
157	CONSERVATION OVERVIEW	82
158	LIMITING FACTORS AND RECOMMENDED APPROACHES	83
159	Limiting Factor: Altered Fire Regimes.....	83
160	Limiting Factor: Invasive Species.....	84
161	Limiting Factor: Damage to Microbiotic Soil Crusts.....	85
162	Limiting Factor: Conversion to Other Land Uses	85
163	Limiting Factor: Loss of Habitat Connectivity	86
164	HABITAT CHANGE TRENDS ANALYSIS	86
165	RESOURCES FOR MORE INFORMATION.....	87
166	REFERENCES.....	88
167	WETLANDS.....	89
168	ECOREGIONS	89
169	CHARACTERISTICS	89
170	[Spotlight] Beaver Habitat and Beaver Modified Habitat	90
171	ECOREGIONAL CHARACTERISTICS	91
172	Blue Mountains.....	91
173	Coast Range	92
174	Columbia Plateau	92
175	East Cascades	92
176	Klamath Mountains	93
177	Northern Basin and Range	93
178	West Cascades.....	93

179	Willamette Valley	94
180	CONSERVATION OVERVIEW.....	94
181	[SPOTLIGHT] White-faced Ibis.....	95
182	LIMITING FACTORS AND RECOMMENDED APPROACHES	95
183	Limiting Factor: Habitat Loss.....	95
184	Limiting Factor: Drought	97
185	Limiting Factor: Water Quantity	97
186	Limiting Factor: Degraded Water Quality	98
187	Limiting Factor: Invasive Species.....	98
188	RESOURCES FOR MORE INFORMATION.....	99
189	NEARSHORE HABITATS	100
190	ECOREGIONS	100
191	CHARACTERISTICS.....	100
192	Neritic Habitat (Open Water).....	100
193	Soft Bottom Subtidal.....	103
194	Rocky Subtidal.....	104
195	Rocky Shore.....	110
196	Sandy Beaches	114
197	CONSERVATION OVERVIEW.....	116
198	LIMITING FACTORS AND RECOMMENDED APPROACHES.....	116
199	Limiting Factor: Public Awareness	116
200	Limiting Factor: Climate Change and Disruption and Disturbance Regimes.....	117
201	Limiting Factor: Land Use Changes (Marine Spatial Planning).....	117
202	Limiting Factor: Pollution and Water Quality.....	118
203	Limiting Factor: Non-native and Invasive Species	119
204	RESOURCES FOR MORE INFORMATION.....	119
205	SPECIALIZED AND LOCAL HABITATS	120
206	ALPINE HABITATS: MEADOWS, DWARF SHRUBLANDS, ALPINE TUNDRA, AND	
207	WHITEBARK PINE	120
208	AQUATIC VEGETATION BEDS	120

209	ASH FLOWS, ASH BEDS AND LAVA FIELDS.....	121
210	BALDS AND BLUFFS	121
211	BAYS	122
212	BITTERBRUSH COMMUNITIES.....	122
213	CANYON SHRUBLANDS.....	123
214	CAVES AND OLD MINES.....	123
215	CHAPARRAL AND CEANOTHUS SHRUBLAND	123
216	EELGRASS BEDS	124
217	FEN PEATLANDS	124
218	FOREST OPENINGS	125
219	UNIQUE GRASSLAND HABITATS.....	126
220	GREASEWOOD FLATS AND WASHES	126
221	INLAND DUNES.....	126
222	INTERDUNAL LAKES AND WETLANDS.....	127
223	INTERTIDAL MUDFLATS	127
224	KELP BEDS.....	128
225	MOUNTAIN MAHOGANY WOODLAND AND SHRUBLAND.....	128
226	OFF-CHANNEL HABITAT	129
227	PORT ORFORD CEDAR FORESTS.....	129
228	ROCK HABITATS: CLIFFS, RIMROCK, ROCK OUTCROPS, AND TALUS.....	129
229	ROCKY SHORES, TIDEPOOLS, AND OFFSHORE ROCKS (E.G., SEA STACKS).....	130
230	SALT DESERT SCRUB.....	130
231	SAND SPITS, SAND BARS, AND SPARSELY VEGETATED ISLANDS.....	131
232	SPRINGS, SEEPS, AND HEADWATERS.....	131
233	SPRING-FED STREAMS	132
234	WESTERN JUNIPER SAVANNA WITH MATURE TREES; LATE SUCCESSIONAL WESTERN	
235	JUNIPER WOODLANDS.....	132
236	WESTERN LARCH FOREST AND WOODLAND	133
237		

238 KEY HABITATS

239 Key Habitats are habitats of conservation concern within Oregon that provide important
240 benefits to **Species of Greatest Conservation Need**. There are 12 Key Habitats within
241 Oregon’s State Wildlife Action Plan, **designated by ecoregion**, including habitats found in
242 the Nearshore ecoregion. The SWAP also describes **Specialized and Local Habitats** that
243 represent important landscape features not adequately addressed through the 12 Key
244 Habitats. Each Key Habitat includes a general description, conservation overview, and a list
245 of limiting factors and recommended approaches. This information is intended to provide a
246 broad summary of the habitat and its most significant conservation needs. Conditions may
247 vary by site, watershed, or ecoregional level based on differences in soil, climate, and
248 management history. Local conditions will need to be considered when determining site-
249 appropriate conservation actions.

250 **Strategy Habitat Methodology**

251 In Oregon’s original State Wildlife Action Plan, the Oregon Conservation Strategy (released
252 in 2006), Key Habitats were determined in a two-step process. First, best available and
253 most recent (in 2006) vegetation maps were compared to historical vegetation maps from
254 1850 to indicate vegetation types experiencing high degrees of loss since European
255 settlement in Oregon. Second, similar vegetation types were classified into “habitats”,
256 which were then evaluated for historical importance at the ecoregional scale, emphasizing
257 the amount of remaining habitat being managed for conservation values, known limiting
258 factors and potential issues impacting habitats, ecological similarity of habitats, and the
259 importance of each habitat to Species of Greatest Conservation Need. The habitats
260 determined to be of the most importance throughout the state were defined as Key
261 Habitats, and were designated by ecoregion. Nearshore Habitats describe the Coastal and
262 Marine Ecological Classification Standard (CMECS) habitat classification approach. See
263 **Appendix - Marine Habitat Classification** for more information.

264

265 ASPEN WOODLANDS

266 Aspen (*Populus tremuloides*) woodlands are woodland and/or forest communities
267 dominated by aspen trees with a forb, grass, and/or shrub understory. Aspen woodlands
268 also occur within conifer forests.

269 ECOREGIONS

270 Aspen woodlands are a Key Habitat in the Northern Basin and Range, Blue Mountains, and
271 East Cascades ecoregions. Small pockets of aspen can also be found in the Klamath
272 Mountains and Willamette Valley ecoregions.

273 CHARACTERISTICS

274 In open sagebrush habitat, aspens typically form woodland and/or forest communities,
275 dominated by aspen trees with a forb, grass, and/or shrub understory. In forested mountain
276 habitats, aspen typically occur within conifer forests. In drier landscapes, aspen primarily
277 occur in riparian areas or in moist microsites. Aspen habitats evolved in areas that
278 historically experienced fire. Given sufficient moisture and light, aspen will sprout annually,
279 with a tendency to sprout more vigorously after disturbance, like wildfire. Within a stand,
280 aspen trees reproduce vegetatively, producing clonal root sprouts arising from a parental
281 root system. While the aspen clone or genet may last for thousands of years, individual
282 trees may only live for 100-150 years. Without disturbance, aspen stands tend to decrease
283 in size (total acres covered) and may be lost to competition from encroaching conifer trees.

284 CONSERVATION OVERVIEW

285 Aspen woodlands are on the edge of their range in Oregon and are more common further
286 east in the Rocky Mountains and north into Canada. One of the few deciduous trees found
287 in eastern Oregon, Aspen Woodlands are especially important in the Northern Basin and
288 Range and Blue Mountains ecoregions. In a landscape dominated by shrubs and grasses,
289 aspen provide significant vertical structure that is useful as nesting and roosting sites for
290 birds and bats and cover for wildlife. Aspen stands also generally have high invertebrate
291 prey diversity and densities. Further, Aspen woodlands provide fawning and calving habitat,
292 security cover, and forage for mule deer and Rocky Mountain elk. Other wildlife that use
293 aspen include black bear, porcupine, beaver, rabbit, and grouse. Tree Swallows,
294 woodpeckers, and other birds nest in aspen cavities.

295 Throughout the west, there is concern over the loss of aspen habitats and the lack of aspen
296 regeneration and recruitment in remnant stands. The Northern Basin and Range ecoregion
297 has lost a large percentage of its aspen woodlands since the 1800s. Aspen stands often
298 depend on natural fire and disturbance to reduce competition from conifers and stimulate
299 the growth of sprouts from roots. Chronic overgrazing can prevent overstory recruitment,
300 allow invasive plant species to establish, and degrade understory plant communities.
301 Overgrazing can also cause erosion, ultimately lowering the water table, which negatively
302 impacts aspen habitats. Many existing aspen trees are reaching the end of their natural life
303 cycle, and without the appropriate recruitment of young aspen, many stands will be lost
304 completely. Aspen typically do not occur in the hottest, driest portions of the Northern
305 Basin and Range ecoregion. As the climate changes, warming temperatures and alterations
306 to hydrologic regimes may impact aspen life cycles and the distribution of this Key Habitat.

307 **[Spotlight] Beaver Habitat and Beaver Modified Habitat**

308 Beavers are widely distributed across Key Habitats statewide, including Flowing Water &
309 Riparian, Wetland, and Aspen Woodlands. **Beaver habitat**, or habitat for beaver, is the
310 specific combination of water, food, cover, and space that beaver need to support their
311 survival on the landscape through time. Beaver are semi-aquatic species that require still
312 or slow-moving, perennial water at stable depths for cover, protection from predators,
313 access to food resources, and food storage in the winter. Beavers are slow on land and
314 prefer to forage within 100 feet of their water source. They need sufficient early seral stage
315 stream buffers of deciduous and herbaceous riparian vegetation for food and foraging
316 activities. Beavers are highly territorial and require adequate lateral and longitudinal
317 habitat quality and stability to support their occupancy on the landscape. In rivers and
318 stream networks, one beaver family unit (on average two adults, two sub-adults, and two
319 kits) needs approximately 0.5 to 1.5 linear stream miles for ample space to survive,
320 reproduce, and thrive. **Beaver habitat**, habitat for beaver, supports the building blocks that
321 beaver need to create **beaver-modified habitats**, or habitat by beaver.

322 **Beaver-modified habitat**, or habitat by beaver, are the specific conditions beaver create
323 when they alter their terrestrial and aquatic habitat to improve their fitness and survival.
324 Habitat modifications include denning, damming and ponding water, creating canals or
325 side-channels, importing woody and vegetative materials into flowing water and wetlands,
326 and changing the structure of riparian vegetative communities. This suite of habitat
327 modifications and their cumulative effects can provide benefits such as increased
328 complexity and connectivity of Key Habitats and habitat, structure, and refugia for SGCN.
329 Nevertheless, beaver activity can also result in flooding, loss of vegetation, economic loss
330 on working lands, and conflict with private landowners. Actions focused on beaver habitat

331 and beaver-modified habitats should also include efforts to mitigate negative impacts and
332 reduce potential conflicts.

333 Habitat limitations for beaver such as declining surface water availability, altered
334 floodplain disturbance regimes, conversion and loss of wet meadow and wetland habitats,
335 and altered riparian vegetation communities are also primary limiting factors for many
336 SGCN.

337 LIMITING FACTORS AND RECOMMENDED APPROACHES

338 **Limiting Factor: Altered Fire Regimes**

339 Aspen stands often depend on natural fire and disturbance to reduce competition from
340 conifers and stimulate the growth of sprouts from roots. Fire suppression has resulted in
341 conifer encroachment and lack of reproduction in aspen communities.

342 **Recommended Approach**

343 Carefully reintroduce natural fire regimes using site-appropriate prescriptions, accounting
344 for the area size and vegetation characteristics that affect resiliency and resistance to
345 disturbance. Prescribed fire has been successful with regenerating aspen groves by
346 increasing sprouting. Use mechanical treatment methods (e.g., masticating, cutting for
347 firewood) to control encroaching conifers. Apply treatments appropriately with respect to
348 season, size, and location. Pursue landscape level treatments, working to restore
349 connectivity of aspen communities. The inclusion of mechanical ground disturbance to
350 stimulate the growth of sprouts from root structures may be one approach to offsetting the
351 lack of fire, but the results of this type of treatment are less predictable.

352 **Limiting Factor: Overgrazing**

353 Overgrazing has limited aspen recruitment through direct consumption or trampling of
354 sprouts and indirect effects such as limiting water availability. When conditions are over-
355 grazed, aspen may sprout but not fully grow into trees. Heavy cattle and ungulate pressure
356 can also impact the soil, herbaceous layer, and recruitment. The direct consumption of
357 aspen and terminal buds tends to be the greatest when sites are used by multiple species
358 such as cattle, sheep, deer, and elk.

359 **Recommended Approach**

360 Limit over-grazing. Use fencing and exclosures to encourage reproduction at high priority
361 sites until trees exceed browse height. Grove protection may be necessary for up to 10

362 years if elk are also present. Implement grazing plans to maintain aspen health, such as
363 limiting grazing during spring and summer.

364 **Limiting Factor: Invasive Species**

365 Invasive plants, introduction of non-native pasture grasses, and historical overgrazing have
366 altered the understory of many aspen stands. Invasive plants may also limit aspen
367 suckering by crowding out and overtopping young sprouts. Juniper have reduced soil
368 moisture in many rangeland aspen groves, increasing the presence of more drought-
369 tolerant upland plants. Prolonged intensive grazing by livestock can lead to increased
370 noxious weeds if grasses and sedges are overutilized. Stewardship actions intended to help
371 aspen (e.g., conifer removal, fire) often stimulate noxious weeds, which thrive in disturbed
372 and open areas.

373 **Recommended Approach**

374 Emphasize prevention, risk assessment, early detection, and quick control to prevent new
375 invasive species from becoming fully established. Control invasive plants using site-
376 appropriate herbicides and methods. Reintroduce native bunchgrasses and flowering
377 plants at priority restoration sites. Minimize soil disturbance in high priority areas to prevent
378 the establishment of invasive plants.

379 **Limiting Factor: Drought**

380 Persistent drought is already occurring in regions and climate change is increasing the
381 frequency and severity of extreme weather events, including heatwaves and droughts.
382 Climate change models predict that more frequent, longer, and more severe regional
383 drought conditions will increase as summer precipitation continues to decrease,
384 exacerbating wildfire risk and reducing water availability. Drought has been shown to
385 increase aspen mortality and reduce recruitment, which could lead to long-term declines
386 in aspen habitat. The interactive effects of increased drought, chronic grazing, and fire
387 suppression are an increasing concern in Oregon.

388 **Recommended Approach**

389 Implement monitoring to detect changes in regeneration, growth, and mortality in drought
390 prone areas. Manage grazing to reduce effects on the water table. Reduce or eliminate
391 encroaching vegetation that compete with aspen for water. Actions that reduce the
392 impacts of other stressors will also improve aspen resilience to drought.

393 RESOURCES FOR MORE INFORMATION

394 [Land Manager’s Guide to Aspen Management in Oregon](#)

395 [US Forest service Guide on Managing Aspen](#)

396 [Guide to Quaking Aspen Ecology and Management \(2017\)](#)

397

DRAFT

398 COASTAL DUNES

399 Occurring along the Oregon coastline, coastal dunes provide habitat for species that prefer
400 open, sandy habitats with a high degree of disturbance from winds and tides.

401 ECOREGIONS

402 Coastal dunes are a Key Habitat in the Coast Range ecoregion.

403 CHARACTERISTICS

404 The Coastal Dunes Key Habitat includes beaches, foredunes, sand spits, deflation plains,
405 and active dunes to stabilizing back dunes. The vegetation varies from sparse to forested,
406 as influenced by sand scour, deposition, movement, and erosion. Species composition is
407 also influenced by salt spray, storm tidal surges, wind abrasion, and substrate stability.
408 Beaches and sandspits are directly impacted by tidal action and are unvegetated.
409 Foredunes generally have unstable sand and sparse to moderate vegetative cover,
410 including native beachgrass, seashore bluegrass, gray beach peavine, largehead sedge,
411 beach morning glory, yellow sand verbena, and silver beachweed. In dunes with greater
412 sand stability, red fescue, seashore lupine, beach pea, coastal strawberry, dune tansy,
413 beach knotweed, and pearly everlasting are dominant. Over time, with plant succession,
414 dunes convert to coastal prairies and grasslands, then to shrublands dominated by salal,
415 and evergreen huckleberry, and eventually to forests dominated by shore pine, Sitka
416 spruce, western hemlock, and Douglas-fir.

417 CONSERVATION OVERVIEW

418 Coastal dune communities have been altered dramatically through the introduction and
419 spread of non-native European beachgrasses, which outcompetes native vegetation and
420 stabilizes foredunes. The stabilized foredunes block movement of sand inland and
421 artificially accelerate plant succession toward shrubland and forest. Dunes artificially
422 stabilized by non-native beachgrasses have contributed to commercial and residential
423 development of sandy habitats that were once naturally active, shifting shoreline
424 ecosystems. In Oregon, almost all of coastal dunes have been altered from their natural
425 state since 1850.

426 Species that live in coastal dune habitats generally prefer open, sandy environments with a
427 high degree of disturbance from winds and tides. Species of Greatest Conservation Need
428 associated with coastal dunes include the Western Snowy Plover, pacific marten, pink
429 sand verbena, Wolf's evening primrose, silvery phacelia, seaside gilia, and Siuslaw hairy-
430 necked tiger beetle.

431 LIMITING FACTORS AND RECOMMENDED APPROACHES

432 **Limiting Factor: European Beachgrass and other Invasive Plants**

433 European beachgrass stabilizes dunes, resulting in changes in vegetative communities and
434 loss of open sandy habitats that are vital to native species. Stabilized dunes are vulnerable
435 to other invasive non-native species, such as hybrid beachgrasses, Scotch broom, and
436 gorse, which displace native plants and animals and accelerate succession.

437 **Recommended Approach**

438 Use mechanical and chemical treatment to control European beachgrass in priority areas,
439 such as western snowy plover nesting areas and near pink sand verbena populations. Build
440 on existing restoration efforts to control beachgrass. Control key invasive non-native plants
441 using site-appropriate tools, such as mechanical (e.g., mowing, girdling, hand-pulling),
442 chemical, and biological control (for gorse) treatments.

443 **Limiting Factor: Development**

444 Stabilized dunes are targeted for development for residential housing, which leads to
445 habitat loss and increased direct/indirect impacts to wildlife through disturbance.

446 **Recommended Approach**

447 Use voluntary cooperative approaches, such as financial incentives, Candidate
448 Conservation Agreements with Assurances, and conservations easements with private
449 landowners to maintain dune habitats. Work with agency partners to support and
450 implement **Statewide Land Use Goal 18**, “Beaches and Dunes”.

451 **Limiting Factor: Recreational Impacts**

452 In some areas, recreational use can disturb wildlife habitat (e.g., western snowy plover
453 nesting areas). Off-highway vehicles can also impact vegetation and disturb wildlife.

454 **Recommended Approach**

455 Work with land managers to direct recreational use away from sensitive areas. Close areas
456 to access during sensitive or vulnerable periods. Provide recreational users with
457 information on coastal dune conservation issues and low impact uses.

458 HABITAT CHANGE TRENDS ANALYSIS

459 **Loss of Coastal Dunes**

460 To investigate loss of coastal dune habitat, the Institute of Natural Resources (INR)
461 compared the total area and spatial overlap of vegetation classes in two baseline maps
462 (1855-1910 and 2016). The analysis showed loss of coastal dune habitat over time. By
463 2016, the total area of coastal dunes had declined by 24% when compared to historical
464 data. There was also evidence of significant shifts in where open dune habitat is located,
465 with some previously open dunes becoming vegetated and stabilized, and new open sand
466 dunes established where dunes was previously stabilized.

467 RESOURCES FOR MORE INFORMATION

468 [Oregon Coastal Management Program](#)

469 [Oregon Dunes Cooperative Weed Management Area: Management Plan](#)

470 [An analysis of coastal sand dune management in Oregon \(United States\) from the 19th](#)
471 [to the 21st century.](#)

472 [Oregon Dunes Restoration Collaborative](#)

473 REFERENCES

474 Brunner, R. and E. Gaines. 2025. Oregon Vegetation Change 1851-2023. Trends analysis
475 conducted for Oregon Department of Fish and Wildlife. Institute for Natural Resources,
476 Portland State University, Portland, OR, USA.

477 Weidemann, A.M., L.J. Dennis, and F.H. Smith. 1999. *Plants of Oregon Coastal Dunes*.
478 Oregon State University Press, 120 p.

479 OPRD's Ocean Shores Management Plan:

480 https://www.oregon.gov/oprd/PRP/Documents/PRP_PLA_OS_FinalOceanShoresMP052305
481 [.pdf](#)

482 Snowy Plover Habitat Conservation Plan:

483 https://www.oregon.gov/oprd/PCB/Documents/WSP-HCP_08182010-web.pdf

484

485 ESTUARIES

486 Estuaries are broadly defined as partially enclosed coastal bodies of tidally influenced
487 water with one or more inputs of freshwater, and with a free or intermittent connection to
488 the open sea. Estuaries typically occur at locations where freshwater from rivers, streams,
489 or creeks meets saltwater from the nearshore ocean, creating a tidal basin that
490 experiences frequent flooding and draining and periodic changes in salinity and other water
491 parameters. Freshwater tidal estuaries can also occur in large floodplain rivers, such as the
492 Columbia River, that are strongly influenced by riverine and estuarine hydrology.


493 ECOREGIONS

494 Estuaries are identified as a Key Habitat within the Coast Range and Nearshore ecoregions.

495 CHARACTERISTICS

496 Estuaries are characterized by the mixing of fresh and salt water within a semi-enclosed
497 tidal basin, by the flux and dynamics of sediments and nutrients, and by the composition
498 and functions of distinct biological communities. The spatial extent of each Oregon estuary
499 begins on the seaward side where it meets the ocean and extends upstream and inland to
500 where the average difference between tidal water levels is 0.2 ft (0.06 m). In many cases,
501 the estuarine tidal basin encompasses a marine-dominated zone, a mixing zone, and a
502 brackish-to-fresh zone that can extend many miles inland away from the ocean.

503 Oregon's statewide framework for management planning within estuaries (Goal 16) seeks
504 to recognize and protect the unique environmental, economic, and social values of each
505 estuary, and (where appropriate) develop and restore the long-term environmental,
506 economic and social values, diversity, and benefits. The statewide planning framework
507 classifies estuaries as development (deep or shallow draft), conservation, or natural, which
508 define the prominent use or activities in the estuary and specify allowed locations for
509 various uses (Figure 1). All of Oregon's estuaries are crucial to the coastal and nearshore
510 ecology and support a diversity of habitats and species.



511

512 **Figure 1.** Oregon’s major estuaries are classified into four levels for development and
 513 planning purposes.

514 **Physical Environment**

515 Several distinct geomorphic types of estuaries occur on the Oregon coast and along the
 516 Columbia River. The geological and hydrodynamic forces that created each estuary differ
 517 from place to place, and the physical environment that maintains them varies substantially
 518 over space and time. Estuaries have been grouped using a number of different
 519 classification schemes that may account for differences in geomorphology, region, or the
 520 relative importance of marine and watershed inputs. These classifications include:

521 *River-dominated drowned river mouth estuaries:* (i.e., Columbia River, Necanicum,
522 Nehalem, Nestucca, Salmon River, Siletz, Alsea, Siuslaw, Umpqua River, Coquille River,
523 Rogue River, Chetco River) The mouths of these river-dominated estuaries were inundated
524 by rising sea levels, and they are characterized by substantial in-flows of freshwater that
525 drain coastal watersheds. The strong riverine input has a primary influence on the shape of
526 the tidal basin, level of salinity, sediment dynamics, and ecological characteristics of the
527 waters and shoreline habitats, rather than marine forces such as the ebb and flow of daily
528 tides.

529 *Tide-dominated drowned river mouth estuaries:* (i.e., Tillamook Bay, Yaquina Bay, Coos
530 Bay) These are low-lying coastal areas where a former river valley was flooded by rising sea
531 levels, and the geomorphology of the estuarine tidal basin is primarily shaped and
532 influenced by strong tidal currents and only weakly influenced by river flows. These
533 estuarine tidal basins are typically very broad and shallow, and contain numerous inlets,
534 sloughs, submerged aquatic vegetation (eelgrass and saltmarshes) and submersible lands
535 such as tideflats, mudflats, and shoals.

536 *Bar built basins and lagoons:* (i.e., Netarts Bay, Sand Lake, Lake Lytle, Smith Lake) These
537 bar-built estuaries and lagoons are formed by periodic deposition of sand and other
538 sediments to create a restriction or semi-permanent barrier to inundation by saltwater. Bar
539 built basins and lagoons typically contain calm waters and protected habitats that are
540 isolated to some extent from the driving forces of nearshore ocean waters.

541 *Blind drowned river mouth estuaries:* (i.e., New River, Sixes River, Elk River, Pistol River,
542 Winchuck River) These estuaries were formed when small coastal river valleys were
543 inundated and flooded by rising sea levels, but the openings to the ocean are partially or
544 completely blocked by a natural barrier such as sandbars or sandy berms. These “blind”
545 estuaries do not have a permanent open connection to the sea.

546 *Tidally restricted coastal creeks:* (i.e., Beaver Creek, Yachats River, Siltcoos, numerous
547 others) These small estuaries occur in areas where rivers, small coastal creeks and
548 streams empty into the ocean, typically across gravel bars or sand. At some times of the
549 year, the outflow from these coastal creeks may be partially impaired and the protected
550 waters can become influenced by the tides.


551 *Marine coves, inlets, and harbors:* (Depoe Bay, Sunset Bay, others) These small marine-
552 dominated coves or sheltered inlets have narrow entrances that protect them from the
553 direct forces or waves and wind, and they are often accompanied by minor outflows from
554 small freshwater creeks or streams.

555 Oregon’s estuarine habitats are characterized and described using the **Coastal and**
556 **Marine Ecological Classification Standard** (CMECS; 2018), a federal classification

557 system that provides a common framework for presenting, classifying, and interpreting
558 spatial data and observational information. The CMECS framework is used to both
559 enhance scientific understanding and advance ecosystem-based resource management.
560 (see **Appendix - Marine Habitat Classification**)


561 The CMECS Oregon Estuarine Aquatic System is composed of riverine subsystems (tidal
562 riverine, diked) and the more saline subsystems found lower in the estuary (coastal, diked,
563 open water). These subsystems are divided where the average salinity during the annual
564 low flow period is less than 0.5 practical salinity units. Aquatic species that inhabit the
565 tidal riverine coastal and tidal riverine open water subsystems differ greatly from those that
566 inhabit the more saline coastal and open water subsystems in all tidal zones.

567 Oregon estuaries are also classified by their CMECS Geoform Components (Figure 2 and
568 3). Geoforms are structural features of the estuarine ecosystem that are geologic in origin,
569 including sloughs, tidal inlets, tidal channels, creeks, deltas, fans, shoreline fans, flats,
570 islands, lagoons, marsh platform, natural levees, and shores. Biogenic geoforms also exist
571 in Oregon estuaries, and include shell beds, burrows in tideflats, and areas of extensive
572 bioturbation. CMECS also recognizes the classification of anthropogenic altered areas as
573 geoforms (i.e., shorelines hardened by rip-rap structures, artificial aquaculture structures,
574 man-made levees, docks and piers, dredge deposits, dredged and excavated channels, fill
575 areas, harbors, marinas, boat ramps). Similarly, Oregon's estuarine habitats include a
576 diversity of CMECS Substrate Components, including natural bedrock, gravel, sand, and
577 mud as well as anthropogenic substrates such as breakwaters, rock jetties, bridge support
578 structures, or artificial materials (pilings) used for construction of docks and piers.



579

580 **Figure 2.** Map of Yaquina Bay depicting CMECS Geoform Components of geologic and
 581 anthropogenic origin.



582

583 **Figure 3.** Map of Yaquina Bay with CMECS Biotic Components.

584 **Biological Community**

585 Oregon estuaries encompass a broad diversity of highly complex, productive habitat that is
586 critical for many species of fish and wildlife, including salmon, rockfish, perch, sculpin,
587 crab, shrimp, bay clams, infaunal invertebrates, marine mammals, and birds. By some
588 estimates, Oregon estuaries support some component of the life cycle for up to three-
589 quarters of all harvested species of fishes, largely due to the high productivity and diversity
590 of habitats, including those provided by eelgrass beds. Rates of primary production in
591 estuarine habitats are very high, and both the emergent vegetation (macroalgae, eelgrass,
592 marsh plants) and microscopic algae (diatoms, others) produce tremendous amounts of
593 organic material that supports the base of the estuarine food web.

594 Eelgrass beds are particularly important because they provide several essential ecosystem
595 functions, including foraging areas and shelter for young fish and invertebrates, production
596 of organic material and detritus, food for migratory waterfowl, and spawning surfaces for
597 species such as the Pacific herring. Eelgrass beds also reduce erosion along the shorelines
598 of Oregon estuaries where the blades and roots trap sediment, stabilize soft
599 unconsolidated substrata, and dissipate the force of wind and waves. Eelgrass improves
600 estuarine water quality by producing oxygen, filtering polluted runoff, absorbing excess
601 nutrients, and reducing localized concentrations of carbon dioxide (See Specialized and
602 Local Habitats). Tidal marshes are also an ecologically productive component of biological
603 communities in Oregon estuaries. Organic materials produced in tidal marshes are broken
604 down by microbial processes to serve as food for many organisms, which in turn are eaten
605 by larger ones as they are distributed throughout the estuary with the tides. Tidal swamps
606 provide complex habitat with layered vegetation, including low-growing herbaceous
607 plants, shrubs, and trees. In addition, tidal swamps generate large quantities of above- and
608 below-ground woody debris, and they provide deep, sheltered tidal channels and deep
609 soils rich in organic matter.

610 Many other species of fish and wildlife also use estuaries. Elk herds graze in tidal marshes
611 and shelter in tidal swamps, bears forage in tidal swamps, river otters build dens, racoons
612 forage along the shore, and rails, snipe, and songbirds nest in the dense vegetation.
613 Estuaries also provide important wintering habitat for waterfowl, including the Black Brant,
614 and migration stopover feeding areas for many shorebirds. Native eelgrass is an important
615 component of an estuary, providing important habitat for several Species of Greatest
616 Conservation Need (SGCN) and other species of conservation interest, including **Black**
617 **Brant, Dungeness crab, black rockfish, copper rockfish, and kelp greenling.** Beds of
618 **eelgrass** provide important spawning substrate for herring (an important forage fish
619 species), blue mud shrimp, native Olympia oysters and native littleneck clams.

620 CONSERVATION OVERVIEW

621 Tidal channels, inlets, sloughs, tideflats, marshes, embayments, and sandy barrier spits
622 that characterize Oregon estuaries are dynamic coastal and riverine systems that respond
623 readily to disturbance by natural and anthropogenic events. The long-term health and
624 sustained productivity of these estuaries are of conservation. Particular attention should
625 be focused on the chronic adverse effects of anthropogenic disturbances and ecological
626 stressors (i.e., invasive species, industrial contaminants, aquaculture operations, habitat
627 alterations, shoreline development, and recreational activities) on the physical structure
628 and ecological functions of estuarine habitats.

629 The spatial extent of Oregon estuaries and tidal wetlands has been significantly reduced
630 over the past 150 years due to road building, diking and filling, development of shoreline
631 municipalities and industries, and conversion of historic tidal wetlands to shoreline
632 agricultural purposes. Oregon's historic estuarine areas have been lost due to
633 anthropogenic disturbance, and even greater losses of historic estuarine habitat have
634 occurred within low-lying estuarine tidal basins that were altered to accommodate
635 shoreline dairy operations. Large expanses of historic forested tidal wetlands (>90%) have
636 been lost, along with substantial losses of salt and freshwater marshes and other tidal
637 wetlands that were diked, drained, and converted to agricultural purposes. Shrub habitat
638 and forested tidal wetlands were historically common around the perimeter of Oregon
639 estuaries, and these habitats were also heavily impacted and experienced substantial
640 habitat loss.

641 In accordance with state planning laws (**Goal 16**), local government comprehensive plans
642 and zoning ordinances have been prepared for all of Oregon's estuaries. Additionally, both
643 estuaries and eelgrass beds are habitat types that have been designated as a Habitat Area
644 of Particular Concern under National Marine Fisheries Service's (NMFS) **Essential Fish**
645 **Habitat** regulations for salmon and groundfish species, designations that require federal
646 agencies to consult with the NMFS before actions are taken.

647 Oregon's remaining estuarine habitats provide a broad diversity of valuable ecological
648 benefits and services, including protection of shorelines from erosion, cycling of nutrients,
649 trapping of sediments, improvement of water quality, production of aquatic vegetation
650 beds, generation of organic material to support food webs, provision of nursery areas and
651 forage sites for fish and shorebirds, and provision of protected waters for recreational and
652 commercial harvest of fish and shellfish. Efforts to conserve healthy estuarine areas and
653 restore degraded habitats will benefit many species, including several commercially
654 important fish and wildlife species.

655 LIMITING FACTORS AND RECOMMENDED APPROACHES

656 **Limiting Factor: Increased Shoreline Development, Land Use Conversion, and Altered or** 657 **Blocked Tidal Flow**

658 Oregon’s estuarine habitats have been altered and lost to a variety of causes, including
659 large-scale dredge and fill operations, diking, ditching, installation of tide gates, residential
660 and industrial development, and drainage of wetlands for dairy operations and other
661 agricultural purposes. Additional estuarine habitat has been lost due to inadequate
662 hydrologic flow through culverts under roads and railroads, creation of log storage areas,
663 and construction of levees, roadways, bridge structures, pilings, docks, and boat
664 launches. Some types of commercial shellfish mariculture practices impact estuarine
665 habitats by disruption of sediment dynamics and causing disturbance to eelgrass beds
666 and their associated communities. Shoreline development projects in the marine-
667 dominated regions of estuaries can impact habitats through the building and maintenance
668 of jetties, piers, breakwaters, marinas, and navigation channels, and disposal of dredge
669 materials can bury and/or alter estuarine habitats and impact nearshore SGCN.

670 **Recommended Approach**

671 Provide technical assistance and incentives to local municipalities, counties, and
672 landowners to protect, conserve, enhance, and restore estuaries. Participate in the
673 planning for state and federal permits associated with dredging of estuary navigation
674 channels and identify mitigation actions necessary to offset unavoidable damages and
675 disturbance. Where appropriate, work to restore hydrology to tidal wetlands by removing
676 dredge spoil materials, opening dikes and levees, filling ditches, and replacing undersized
677 culverts. Continue successful education programs focused on recognizing the beneficial
678 functions and services provided by estuaries. Work with agency partners to support and
679 implement existing land use regulations that preserve and restore habitats. For example,
680 refer to seasonal in-water work windows for estuaries designed to minimize impacts to
681 out-migrating salmon. Continue to develop and refine “best management practices” for
682 commercial shellfish mariculture operations within estuaries. Monitor, maintain and
683 restore eelgrass beds and forested wetlands as key habitat features. (KCI: **Land Use**
684 **Changes**)

685 **Limiting Factor: Alteration of Freshwater Inputs into Estuaries**

686 The amount and timing of freshwater inputs into estuaries are critical to maintaining the
687 hydrological regime that supports delicate estuarine ecosystems. Disruption of freshwater
688 delivery systems can contribute to decreased flushing, inundation of floodplains,
689 increased sedimentation, decreased residence time of water (which reduces the filtering
690 benefits of estuaries), altered fish community dynamics, and/or increased stress on
691 juvenile fish, nekton, or other animals. Changes in hydrological regimes can also make

692 estuaries more susceptible to the establishment and invasion by non-native species as
693 well as accumulation of marine debris and waterborne pollutants.

694 **Recommended Approach**

695 Evaluate the potential impacts of water diversions away from estuaries (e.g., for
696 agriculture, residential, or industrial purposes) on floodplain dynamics and other functions
697 of estuarine systems. Prioritize watersheds and tidal basins for the acquisition of sufficient
698 instream flows.

699 **Limiting Factor: Degraded Water Quality**

700 Water quality in estuaries is frequently degraded by both point and non-point sources of
701 pollution. The sources of degraded waters may originate from the nearshore Pacific
702 Ocean, within the estuary, and/or from sources in the adjacent watershed. Marine waters
703 that flood into estuaries may be impaired (acidified) by elevated concentrations of carbon
704 dioxide or hypoxic (low) levels of dissolved oxygen. In addition, marine waters are
705 periodically contaminated by fuel oil spills, diesel, and other hydrocarbons released by
706 vessels at sea. Contaminated runoff from residential, agricultural lands, commercial
707 forest land, failing septic systems, animal waste, and storm events can enter estuaries and
708 negatively affect water quality. Estuarine water temperatures can become elevated by
709 dredging, sedimentation, stormwater runoff, and altered patterns of tidal circulation. Other
710 discharges, including polluted runoff from commercial boatyards and marinas, discharges
711 from commercial seafood processors, and shore-based cleaning operations, all can
712 contribute to poor estuarine water quality. Estuaries are also susceptible to increased
713 loads of fecal indicator bacteria that can enter the tidal basin from multiple sources.
714 Stormwater runoff that collects water from impervious surfaces and roadways can
715 contribute fertilizers, herbicides, sediments, oil and grease, and other pollutants directly
716 into estuaries and bays.

717 **Recommended Approach**

718 Continue current efforts to consider the impacts of local land-use planning decisions on
719 estuarine water quality. Support efforts of the Oregon Department of Environmental
720 Quality (DEQ) to assess water quality and develop Total Maximum Daily Loads and water
721 quality management plans where necessary to address issues. Continue coordination to
722 ensure that plans and goals consider impacts to water quality sufficient to protect fish and
723 wildlife in addition to other goals (i.e., recreation). Work with cities to improve stormwater
724 management from impervious surfaces, and work with the Oregon Department of
725 Transportation (ODOT), County roadmasters, and industrial forest landowners to reduce
726 stormwater and sediment delivery from roads. Prioritize restoration of eelgrass beds,
727 saltmarshes, and forested and scrub-shrub estuarine wetlands to assist with buffering and
728 filtering water that enters estuaries. (KCI: **Water Quality and Quantity and Pollution**)

729 **Limiting Factor: Non-Native and Invasive Species**

730 Introduced, non-native, and invasive species present a substantial threat to the
731 biodiversity of Oregon’s estuarine habitats. Large estuaries that support maritime trade
732 and commercial mariculture activities (such as the Columbia River and Coos estuary) are
733 particularly vulnerable to colonization by new species of invertebrates, fishes, and plants.
734 Dredge spoils deposited within estuaries provide new habitat that can be rapidly colonized
735 by non-native species, and hydroelectric projects on rivers that flow into estuaries disrupt
736 freshwater inflows and the ecology of estuarine communities. It is estimated that over 100
737 non-native species have become established into the Coos estuary. Many of these species
738 are cryptic, but some displace native species and have the potential to alter habitat
739 structure and energy flow through the estuarine habitats and communities.

740 Commercial shipping vessels transport large volumes of ballast water from one port to
741 another, and they function as vectors for the introduction of living marine organisms. For
742 example, the purple varnish clam was probably transported via ballast water from Japan to
743 British Columbia before 1993. By 1997, this bivalve spread to Oregon, presumably via
744 natural transport of larvae by ocean currents.

745 Some non-native species have been introduced deliberately into Oregon as cultivated
746 seafood products (i.e., Pacific oysters and Kumamoto oysters), while others have become
747 established as inadvertent hitchhikers associated with commercial mariculture
748 operations. For example, large sections of Oregon’s estuarine tideflats have been
749 colonized over the past 35 years by Japanese eelgrass, which takes root in the upper region
750 of muddy tideflats and may compete with native eelgrass. Other undesirable species
751 associated with mariculture operations include seaweeds, predatory oyster drills (snails),
752 mud blister worms and colonial tunicates.

753 The European green crab became established in Oregon estuaries in the mid-1990s, and
754 populations persisted at low abundance for about 20 years. Following a substantial marine
755 heatwave and several successive periods of warm ocean temperatures, the population of
756 European green crab increased rapidly to the point where they are abundant in the mid and
757 upper regions of Oregon estuaries where they prey on small native clams, worms and
758 juvenile flatfish. Other examples of non-native invasive animals found in Oregon estuaries
759 include the parasitic Griffen’s isopod (which has been linked to declines of native blue
760 mud shrimp populations), the New Zealand mudsnail, and the New Zealand burrowing
761 isopod. Invasive species can also be introduced into estuaries through recreational or
762 commercial boating, or the aquarium trade where they have the potential to spread quickly
763 because they have no natural predators or competitors. An extensive list of non-native and
764 invasive species that have been found in the Nearshore ecoregion, including in estuaries,
765 can be found in **Appendix – Nearshore Species**.

766 **Recommended Approach**

767 Emphasize prevention, risk assessment, early detection, and quick control to prevent new
768 invasive species from becoming fully established. Control key invasive plants using site-

769 appropriate tools, such as hand-pulling, covering with geotextile cloth, repeated mowing,
770 flooding, and/or herbicides focusing on spot treatment. Monitor estuaries for potential
771 invasive species, and use site-appropriate methods to detect, trap, and control newly
772 established species (i.e. mud blister worms) for which management can be most effective.
773 Work with state and federal partners to implement existing ballast water regulations,
774 including development of potential methods to treat and disinfect ballast water. Work with
775 partners to limit the spread of invasive species that have become established and
776 naturalized. Explore options to allow for increased harvest of species suitable for human
777 consumption such as purple varnish clams and European green crab. (KCI: **Invasive**
778 **Species**)

779 **Limiting Factor: Management and Planning Needs**

780 Many jurisdictions and agencies have management authority and interest in Oregon
781 estuaries, which can make land-use planning, decision-making for permits, and other
782 actions more complex and difficult. In Oregon, cities, counties, port districts, and many
783 state agencies have planning and management responsibilities for estuaries. In addition,
784 the federal government and Oregon coastal Tribes have some level of management
785 authority for activities in estuaries.

786 **Recommended Approach**

787 Coordination among agencies is a high priority. Because estuarine issues are complex,
788 clear identification and communication of conservation opportunities, goals, and threats
789 should precede management actions, ensuring that all interests are considered and
790 coordinated. Develop a process to provide advanced notice and share information among
791 federal, tribal, state and county agencies to assist with conservation, protection,
792 enhancement, and restoration of estuarine habitats.

793 Develop and implement science-based management strategies for estuarine resources.
794 Expand upon management objectives previously identified and further develop plans that
795 identify restoration or conservation targets for individual estuaries. Encourage and assist in
796 estuarine research to identify data and knowledge needed for management planning.

797 **Limiting Factor: Loss of Habitat Complexity**

798 Habitat complexity provides refugia for estuarine fish and wildlife. Complex habitat
799 supports diverse ecological communities, contributing to resiliency to climate change
800 impacts. Removal or loss of large, downed trees not only reduces habitat complexity but
801 also insect production and food and cover for juvenile salmonids. Disconnection of
802 habitats from the tidal basin and floodplain interrupts the natural transition zones between
803 the aquatic, intertidal, and upland ecosystems. Dredging, ditching, channelization, and
804 filling in estuaries alters marine and freshwater inputs and reduces habitat function. In-
805 water (e.g., pilings, jetties, seawalls) or overwater (e.g., roadways, dikes, levees, mooring

806 buoys, floating docks) structures can reduce habitat complexity, as can bayside
807 development that extends into intertidal areas. Natural factors can also reduce habitat
808 complexity, such as damage or movement caused by seasonal runoff or significant storm
809 events, especially where the estuary has already been compromised, and floodplains have
810 been lost.

811 **Recommended Approach**

812 Ensure that permit application reviews consider alternative sites and practices to avoid and
813 minimize impacts and provide full and effective mitigation to offset unavoidable damages.
814 Encourage and participate in cooperative efforts and incentives to promote habitat
815 complexity in estuaries and consider the scale of development proposals in reference to
816 historical and future baselines. Prioritize conservation and restoration efforts to restore
817 floodplain connectivity, tidal marshes, and forested wetlands, and to conserve eelgrass.
818 Increase outreach and education about the importance of habitat complexity, including the
819 benefits of increased complexity associated with recovering populations of native Olympia
820 oysters.

821 **Limiting Factor: Climate Change**

822 Climate change is expected to have significant impacts to Oregon estuaries. Rising sea
823 levels are expected to more fully inundate estuarine tidal basins, resulting in changes to
824 the delivery of marine-derived nutrients and tidal hydrology, shifts in water temperatures,
825 disruption of salinity regimes, advancement of the tidal prism, changes in the deposition
826 and erosion of sediments, and losses of tidal wetlands and submerged aquatic vegetation
827 (coastal squeeze). Acidified ocean waters are impacting estuaries and contribute to
828 biogeochemical shifts in the composition of estuarine waters and difficulties in shell-
829 building for estuarine bivalves. Shifts in habitat conditions within estuaries may contribute
830 to increased colonization by non-native species, and alteration of estuarine food webs.
831 Further inland, warming and drying conditions in coastal watersheds may impact the
832 characteristics of freshwater flows into estuaries.

833 **Recommended Approach**

834 Use emerging models of future sea level rise and changing salinity regimes to inform
835 conservation actions in estuaries. Work with property owners, land use planners, and
836 restoration practitioners to focus attention on vulnerable areas. Support efforts to restore
837 natural processes of tidal exchange and sediment deposition, which will enable tidal
838 wetlands to maintain their elevation relative to rising sea levels. Support efforts to re-
839 connect floodplains to adjacent uplands by removing barriers, placement of large woody
840 debris, and planting of riparian areas. Conserve areas that will become new marshes and
841 forested wetlands with sea level rise. Inform communities about climate change impacts
842 and support community preparedness. (KCI: **Climate Change**)

843 **Limiting Factor: Oil Spills and Hazardous wastes**

844 Oregon estuaries are susceptible to periodic exposure and contamination by fuel oil,
845 petroleum products, creosote, and other hazardous materials. Hazmat spills are of
846 particular concern in deep-draft estuaries that support transport, loading, and unloading
847 of large commercial vessels, and areas with busy marinas that provide for refueling and
848 berths for commercial and recreational vessels. Estuarine tidal basins have also been
849 contaminated by legacy pollutants (heavy metals, oil and grease, etc.) and industrial
850 waste, and some sites are treated as USEPA “Superfund Cleanup” sites (i.e. Port of
851 Portland, Tongue Point). All of Oregon’s estuarine areas are at risk from oils spills that
852 occur in the ocean or along the open coast because buoyant hydrocarbons may enter
853 estuarine tidal basins on flooding tides. If a spill occurs, accumulation of oil and hazardous
854 materials can have long lasting impacts.

855 **Recommended Approach**

856 Participate in the periodic review and updates to the Oregon Geographic Response Plans
857 and oil spill contingency plans and ensure that the maps for the coast and estuaries
858 contain up-to-date information regarding living marine and estuarine resources. Maintain
859 status as emergency Hazwoper Responders and participate in interagency drills and
860 training exercises. Work with the Oregon Department of Geology and Mineral Industries,
861 Oregon Department of Environmental Quality, the US Coast Guard, and local emergency
862 officials to identify hazardous material use and storage sites in high-risk areas and seek
863 ways to minimize these risks. Coordinate with agencies to periodically communicate
864 about Hazardous material storage, transportation, and response issues to decrease
865 environmental risks and increase understanding of the impacts of Hazmat spills. (KCI:
866 **Pollution**)

867 **RESOURCES FOR MORE INFORMATION**

868 [Oregon Coastal Atlas Estuary Data Viewer, and background on CMECS classification](#)
869 [system](#)

870 [Pacific Marine and Estuarine Fish Habitat Partnership](#)

871 [National Water Quality Assessment Program](#)

872 [South Slough National Estuarine Research Reserve](#)

873 [ODFW Workshop on Estuaries, Climate Change, and Conservation Planning \(2010\)](#)

874 REFERENCES

- 875 Barbier, E.B., S.D. Hacker, C. Kennedy, E.W. Koch, A.C. Stier, and B.R. Silliman. 2011. The
876 value of estuarine and coastal ecosystem services. *Ecological Monographs* 81: 169-193.
- 877 Bottom, D.L., K.K. Jones, T.J. Cornwell, A. Gray, and C.A. Simenstad. 2005. Patterns of
878 Chinook salmon migration and residency in the Salmon River estuary (Oregon). *Estuarine,
879 Coastal and Shelf Science* 64: 79-93.
- 880 Brophy, L.S., C.M. Greene, V.C. Hare, B. Holycross, A. Lanier, W.N. Heady, K. O'Connor, H.
881 Imaki, T. Haddad, and R. Dana. 2019. Insights into estuary habitat loss in the western
882 United States using a new method for mapping maximum extent of tidal wetlands. *PLOS
883 ONE* 14(8): e0218558
- 884 Brown, C.A. and R.J. Ozretich. 2009. Coupling between the coastal ocean and Yaquina Bay,
885 Oregon: importance of oceanic inputs relative to other nitrogen sources. *Estuaries and
886 Coasts* 32:219–237.
- 887 Good, J.W. 2000. Summary and current status of Oregon's estuarine ecosystems. In,
888 Oregon State of the Environment Report. Oregon Sea Grant and Oregon State University,
889 Technical Supplement to the progress Board (September 2000)
890 https://www.oregon.gov/DSL/WW/Documents/soer_ch33.pdf
- 891 Flitcroft, R. et al., 2016. Expect the unexpected: place-based protections can lead to
892 unforeseen benefits. *Aquatic Conservation: Mar. Freshwater. Ecosystem.* 26 (Suppl. 1): 39–
893 59.
- 894 Hessian-Lewis, M.L., S.D. Hacker. 2013. Upwelling-influence, macroalgal blooms, and
895 seagrass production; temporal trends from latitudinal and local scales in northeast Pacific
896 estuaries. *Limnology and Oceanography* 58:1103-1112.
- 897 Hickey, B.M. and N.S. Banas. 2003. Oceanography of the U.S. Pacific Northwest coastal
898 ocean and estuaries with application to coastal ecology. *Estuaries* 28:1010-1031.
- 899 Howard, B.R., F.T. Francis, I.M. Cote, T.W Therriault. 2019. Habitat alteration by invasive
900 European green crab (*Carcinus maenas*) causes eelgrass loss in British Columbia, Canada.
901 *Biological Invasions* 21: 3607-3618.
- 902 Lee, Il, H., and C.A. Brown. 2009. Classification of regional patterns of environmental
903 drivers and benthic habitats in Pacific northwest estuaries. U.S. Environmental Protection
904 Agency / Office of Research and Development. National Health and Environmental Effects
905 Research Laboratory, Western Ecology Division. EPA/600/R-09/140.

- 906 Mapping efforts of Oregon's estuaries utilizing CMECS components is underway and an
907 online tool for viewing estuary maps is available at:
908 <http://www.coastalatlantlas.net/estuarymaps/>
- 909 Rumrill, S.S. 2006. Ecology of the South Slough Estuary: Site Profile of the South Slough
910 National Estuarine Research Reserve. OR Department of State Lands / NOAA. 197 pp.
- 911 Rumrill, S.S., D.C. Sowers. 2008. Concurrent assessment of eelgrass beds (*Zostera marina*)
912 and saltmarsh communities along the estuarine gradient of the South Slough, Oregon.
913 *Journal of Coastal Research* 2008(10055): 121-134.
- 914 Simenstad, C.A., Burke, J.L., O'Connor, J.E., Cannon, C., Heatwole, D.W., Ramirez, M.F.,
915 Waite, I.R., Counihan, T.D., and Jones, K.L., 2011, Columbia River Estuary Ecosystem
916 Classification—Concept and Application: U.S. Geological Survey Open-File Report 2011-
917 1228, 54 p.
- 918 Tallis, H.M., J.L. Ruesink, B. Dumbauld, S. Hacker, and L.M. Wisheart. 2009. Oysters and
919 aquaculture practices affect eelgrass density and productivity in a Pacific Northwest
920 estuary. *Journal of Shellfish Research* 28: 251–261.
- 921 Wagner, E., B.R. Dumbauld, S.D. Hacker, A.C. Trimble, L.M. Wisheart, J.L. Ruesink. 2012.
922 Density-dependent effects of an introduced oyster, *Crassostrea gigas*, on a native intertidal
923 seagrass, *Zostera marina*. *Marine Ecology Progress Series* 468:149-160.
- 924 Yamada, S.B. 2001. European green crab – a global invader. Oregon Sea Grant, Corvallis.
925 ORESU-B-01-001, 123 pp.
926

927 FLOWING WATER AND RIPARIAN HABITATS

928 Flowing Water and Riparian Habitats include all naturally occurring flowing freshwater
929 streams and rivers throughout Oregon as well as the adjacent riparian habitat.

930 ECOREGIONS

931 Flowing Water and Riparian Habitats are identified as a Key Habitat in all ecoregions.

932 CHARACTERISTICS

933 **Flowing Water Habitats**

934 Flowing creeks, streams, and rivers are a key feature of the Oregon landscape and our
935 natural resources heritage. They support diverse ecosystems and fisheries, and provide
936 significant social, economic, and recreational values. Healthy flowing freshwater systems
937 are crucial to support iconic Pacific Northwest salmon and steelhead as well as
938 amphibians, aquatic insects, and other Species of Greatest Conservation Need (SGCN).
939 People also rely on healthy waterways to irrigate crops, generate hydroelectric power,
940 operate manufacturing plants, treat wastewater, and for drinking water.

941 Natural, freshwater flowing waters are dynamic systems that typically start as small, high
942 elevation creeks that merge with mid-elevation streams and then combine to form large
943 river systems. Flowing waters are fed by a variety of sources, including melting glaciers and
944 snow, direct runoff from the surrounding landscape or watershed, and via groundwater
945 discharge such as springs. Flowing water habitat includes perennial, intermittent, and
946 ephemeral creeks, streams, and rivers. Perennial waterways are those that flow year-
947 round, whereas intermittent waterways only flow part of the year, typically during the wet
948 season, and ephemeral waterways only flow during a short period after a precipitation
949 event. Protection of habitats surrounding perennial, intermittent, and ephemeral creeks,
950 streams, and rivers helps to minimize impacts to flowing waters while providing benefits to
951 water temperature and water quality.

952 Healthy streams include structural variability essential for SGCN to meet their life cycle
953 needs. For example, pools and riffles provide a range of stream flows, and backwater
954 alcoves and side channels are essential for refugia during high flows. Ephemeral streams,
955 though flowing only during snowmelt or rain events, provide important refugia for
956 anadromous and local fish species during spring high waters. The shape and dynamics of a
957 stream or river are typically defined by high winter/spring flows and flooding patterns, width
958 of the available floodplain, geology/soils in the adjacent floodplain and watershed, and the
959 degree of human impact on water quantity, water quality, and the surrounding landscape.
960 Healthy streams are typically connected with their floodplain, exhibit natural variability in
961 flow amount and timing, and are dynamic and free to evolve based on natural events such

962 as wood falling into the river, channels changing course, high flow events, and landslides.
963 The complexity of the flowing water habitat directly contributes to the health and function
964 of fish-bearing streams.

965 Climate-related changes in precipitation patterns, snowmelt cycles, and fire frequency, as
966 well as increased demand for out-of-stream water use will alter flowing water systems
967 relative to historical conditions. A changing climate has the potential to alter hydrologic
968 regimes and water availability, leaving less water to meet various flowing water and
969 floodplain needs.

970 Protection, maintenance, and restoration of our ecosystems is needed to enhance
971 resiliency by increasing natural storage capacity, improving instream habitat and fish
972 passage, protecting and restoring wetlands and water instream, eradicating invasive
973 species, protecting native plant communities, and protecting groundwater-dependent
974 ecosystems. Land management activities need to protect and improve water quality,
975 including protecting our watersheds and drinking water sources from contamination and
976 pollution.

977 **Riparian Habitats**

978 Riparian habitat zones are adjacent to flowing water in creeks, rivers, and streams as well
979 as springs, seeps, and terraces. They occur at all elevations, from valley bottom
980 floodplains to alpine torrents, and are shaped through seasonal flooding, scour, and soil
981 deposition.

982 Floodplains are diverse habitats adjacent to rivers, streams, lakes, estuaries, or other water
983 bodies that are subject to flooding. In their undisturbed, natural state, these areas act to
984 store excess floodwater, which can protect downstream property from flooding, and
985 release water slowly, extending availability. Floodplains also provide essential habitat for
986 fish and wildlife, including refugia from high stream flows and corridors for wildlife.

987 Riparian zones are the dynamic interface between land and flowing water. The plant
988 assemblages and communities in riparian zones help buffer inputs and the cycling of
989 nutrients, as well as provide habitat for aquatic and terrestrial life. The vegetative
990 composition and structure of riparian zones varies, and is a function of elevation,
991 precipitation pattern, stream gradient, aspect, floodplain width, storage capacity of the
992 soil, groundwater supply, and disturbance (i.e., flooding).

993 In many areas of the state, native riparian vegetation is comprised mostly of deciduous
994 trees and shrubs, such as big-leaf maple, alders, aspen, cottonwood, dogwood, willows,
995 and Oregon ash. Conifers, such as pines, firs, and spruce, dominate some riparian zones
996 at higher elevations and are important in some lower elevation areas as well (e.g., interior
997 Rogue basin). Riparian shrublands may include willows, red osier dogwood, western birch,
998 hawthorn, alder, and chokecherry. Riparian meadows are dominated by grasses, sedges,

999 and rushes. Riparian habitats provide food, cover, and/or breeding sites for many fish and
1000 wildlife species throughout the year.

1001 The Flowing Water and Riparian Key Habitat does not include irrigation structures (e.g.,
1002 ditches) or other man-made waterbodies such as reservoirs. **Natural lakes** are covered
1003 separately, as are **Springs, Seeps, and Headwaters** and **Spring-fed Streams**. The riparian
1004 zones around the edges of natural lakes are included within the **Wetlands** Key Habitat.

1005 CONSERVATION OVERVIEW

1006 Flowing water and the riparian habitat found along its banks are defined together as a **Key**
1007 **Habitat** because their distribution and conservation roles are interconnected. Water is
1008 crucial for all fish and wildlife, and high-quality freshwater aquatic systems provide
1009 essential habitat to many at-risk species, including important spawning and rearing habitat
1010 for salmonids, breeding habitat for amphibians, and habitat for freshwater mussels and
1011 other invertebrates. Flowing water is important to connect ecosystems across elevations
1012 throughout the year. In many locations throughout Oregon, water flow and hydrology have
1013 been impacted by development, including barriers (e.g., roads, dams, and culverts) and
1014 water diversions for out-of-stream uses that can reduce instream flow, increase summer
1015 stream temperatures, and interfere with **fish and wildlife migration**. Channelization and
1016 floodplain development can restrict the natural ability of streams to meander over time,
1017 limiting the quality and availability of these habitats, as well as affecting floodplain
1018 function.

1019 Riparian habitats often have high species diversity and are critical for fish and wildlife,
1020 especially for those species that prefer moist shrubby or forested habitats. Riparian
1021 habitats can maintain favorable water temperature for fish and provide a cooler
1022 temperature refuge for terrestrial species in a warming climate through shading. These
1023 areas also provide essential travel corridors for birds, amphibians, reptiles, mammals, and
1024 other wildlife. In arid areas, such as the **Blue Mountains**, **Northern Basin and Range**, and
1025 **Columbia Plateau** ecoregions, riparian habitats can provide abundant insects, plants, and
1026 moisture throughout the year. Riparian meadows include natural spring-seep habitats that
1027 are extremely important for a wide variety of species, including **Greater Sage-Grouse**.

1028 In addition to providing habitat for birds and other wildlife, riparian habitats have important
1029 ecological functions. Healthy riparian vegetation serves an important role in slowing water
1030 velocities during periods of high runoff and protecting streambeds from scouring and
1031 downcutting. Riparian vegetation also protects stream banks from erosion, influences in-
1032 channel aquatic habitats, filters run off, drives channel complexity in valley bottoms, and
1033 provides nutrients to support terrestrial and aquatic life. Riparian habitats often link upland
1034 and aquatic habitats, which facilitates the role upland habitats play in watershed function.

1035 Riparian habitats have declined from historical levels and are now greatly reduced in area
1036 and connectivity, especially those in low-elevation areas and valley bottoms. Non-native

1037 and often invasive vegetation dominates in many areas. Development, logging, roads,
1038 agricultural practices, beaver removal, and grazing can further degrade riparian habitat.
1039 Removal or reduction of riparian habitat allows runoff containing contaminants such as
1040 fertilizers and pesticides to reach streams and rivers where it can negatively impact
1041 aquatic life.

1042 **[Spotlight] Beaver Habitat and Beaver-Modified Habitat**

1043 Beavers are widely distributed across Key Habitats statewide, including Flowing Water &
1044 Riparian, Wetlands, and Aspen Woodlands. **Beaver habitat**, or habitat for beaver, is the
1045 specific combination of water, food, cover, and space that beaver need to support their
1046 survival on the landscape through time. Beaver are semi-aquatic species that require still
1047 or slow-moving, perennial water at stable depths for cover, protection from predators,
1048 access to food resources, and food storage in the winter. Beavers are slow on land and
1049 prefer to forage within 100 feet of their water source. They need sufficient early seral stage
1050 stream buffers of deciduous and herbaceous riparian vegetation for food and foraging
1051 activities. Beavers are highly territorial and require adequate lateral and longitudinal
1052 habitat quality and stability to support their occupancy on the landscape. In rivers and
1053 stream networks, one beaver family unit (on average two adults, two sub-adults, and two
1054 kits) needs approximately 0.5 to 1.5 linear stream miles for ample space to survive,
1055 reproduce, and thrive. **Beaver habitat**, habitat for beaver, supports the building blocks that
1056 beaver need to create **beaver-modified habitats**, or habitat by beaver.

1057 **Beaver-modified habitat**, or habitat by beaver, are the specific conditions beaver create
1058 when they alter their terrestrial and aquatic habitat to improve their fitness and survival.
1059 Habitat modifications include denning, damming and ponding water, creating canals or
1060 side-channels, importing woody and vegetative materials into flowing water and wetlands,
1061 and changing the structure of riparian vegetative communities. This suite of habitat
1062 modifications and their cumulative effects can provide benefits such as increased
1063 complexity and connectivity of Key Habitats and habitat, structure, and refugia for SGCN.
1064 Nevertheless, beaver activity can also result in flooding, loss of vegetation, economic loss
1065 on working lands, and conflict with private landowners. Actions focused on beaver habitat
1066 and beaver-modified habitats should also include efforts to mitigate negative impacts and
1067 reduce potential conflicts.

1068 Factors that are currently limiting habitat for beaver include declining surface water
1069 availability, altered floodplain disturbance regimes, conversion and loss of wet meadow
1070 and wetland habitats, and altered riparian vegetation communities.

1071 **Oregon Planning and Regulatory Background for Flowing Waters**

1072 Under Oregon law, water is a public resource, meaning that all water belongs to the public.
1073 Cities, irrigators, businesses, and other water users must obtain a permit or license from
1074 the **Oregon Water Resources Department** (OWRD) to use water from any source whether
1075 it is underground, or from lakes or streams, with some exceptions. OWRD is responsible
1076 for allocating new uses of water, whether in cities, farms, factories, or for improvement of
1077 fish habitat, and follows a careful process to preserve the investments already made in the
1078 state.

1079 Oregon's Water Code, established in 1909, created a system of water allocation and
1080 distribution that did not consider water for instream uses, leading to degradation of
1081 Oregon's flowing water and freshwater habitats. Over time, it became clear that a legal
1082 system was needed to protect flows in support of ecological uses. In response, the 1987
1083 Instream Water Rights Act officially recognized instream flows as a beneficial use that
1084 could be protected by a water right, giving them the same legal status as consumptive
1085 water rights. Instream water rights are the state's mechanism to provide water for fish and
1086 wildlife needs and healthy ecosystems that support multiple public uses (e.g., recreation,
1087 fishing, tourism). If there is a conflict between users, however, the date of priority
1088 determines who may use the available water and most instream water rights are quite
1089 junior compared to many out-of-stream water rights.

1090 **Oregon's Planning and Regulatory Background for Riparian Habitats**

1091 Oregon's planning and regulatory framework provides tools to address riparian habitat
1092 conservation issues. Riparian habitat is considered a **Goal 5** resource, where local
1093 governments can adopt protective ordinances through comprehensive plans to establish
1094 riparian buffers. Streamside buffers implemented through the **Northwest Forest Plan**
1095 (NWFP) on public land and the **Oregon Forest Practices Act** on state and private land are
1096 designed to protect riparian health in forested landscapes. On agricultural lands,
1097 Agricultural Water Quality Management Area Rules and Plans have been adopted across
1098 the state to address riparian conditions and other **water quality issues**. While each
1099 riparian rule is slightly different depending on the local area, the riparian rules generally
1100 require agricultural activities to allow establishment, development, and maintenance of
1101 riparian vegetation consistent with site capability to provide moderation of solar heating,
1102 filtration of overland flow, and streambank stability. The Oregon Water Resources
1103 Department also has rules that require the riparian area to be restored or enhanced if it is
1104 disturbed in the process of developing a point of diversion.

1105 **LIMITING FACTORS AND RECOMMENDED APPROACHES**

1106 **Limiting Factor: Water Quantity**

1107 Multiple factors are affecting the amount of water in Flowing Water Habitats. **Water**
1108 **availability** is currently limited in much of the state, especially during the low flow summer
1109 and fall months, and this is expected to increase under a changing climate. Riparian

1110 bottomland habitats compete for water with other uses, particularly in the Blue Mountains,
1111 Columbia Plateau, East Cascades, Klamath Mountain, and Northern Basin and Range
1112 ecoregions. Water diversions for out-of-stream uses occur on all major streams, and valley
1113 bottoms often have multiple canals that divert water away from the natural channel. As a
1114 result, low flows are associated with higher water temperatures and higher nutrient and
1115 contaminant concentrations in creeks, streams, and rivers.

1116 **Recommended Approach**

1117 Conduct instream flow studies to develop ecological flow targets and apply for associated
1118 instream water rights. Engage with regulatory agencies to ensure consideration of fish and
1119 wildlife needs in water right and hydropower processes. Identify priority locations for
1120 voluntary instream transfers and leases and continue to apply for new instream water
1121 rights. Implement water conservation actions to protect or increase instream flows
1122 (quantity, timing, and duration) following the natural hydrological cycle. Increase pace and
1123 scale of voluntary flow restoration through instream leases, transfers, and irrigation
1124 efficiency improvements. Manage beaver populations to contribute to water storage and
1125 availability, when compatible with existing land uses. Pursue collaborative water planning
1126 and implementation processes to secure balanced solutions for water management.
1127 Provide incentives and information about water conservation and sharing at key times of
1128 low flow conditions (e.g., late summer). Assess riparian habitat condition and consider
1129 planting projects to promote shade, which can limit thermal maxima in summer months.

1130 **Limiting Factor: Invasive Aquatic Species**

1131 Alterations in hydrology can make flowing water habitat more susceptible to invasive
1132 plants, invertebrates, and fish. Invasive fish species (e.g. bass, crappie, bluegill, yellow
1133 perch, bullhead, carp, brook trout, fat head minnow, non-native ringed crayfish) can
1134 compete with native fish and amphibians for food resources and habitat, prey on native
1135 species, alter habitat, or hybridize with native fish. For example, non-native carp can
1136 overgraze aquatic vegetation and stir up sediment, depriving native fish and amphibians of
1137 egg-laying sites or preventing eggs from absorbing enough oxygen to develop. Invasive
1138 mollusks (e.g. zebra mussel, quagga mussels) can disrupt food chains by reducing the
1139 availability of food for larval and juvenile fishes. They also attach easily to boats, docks and
1140 buoys and can clog intake pipes as well as drains. Invasive plants (e.g. ludwigia,
1141 watermillfoil, parrot feather, hydrilla) can reduce light penetration, lower species diversity,
1142 alter temperature, reduce dissolved oxygen & pH, and disrupt nutrient cycling, leading to
1143 algae blooms and toxicity.

1144 **Recommended Approach**

1145 Restoration and maintenance of historical hydrological regimes ensures that habitat
1146 conditions best support native fish and wildlife. Work with community partners to restore
1147 flow and water input levels. Continue working with the public to stress the importance of

1148 preventative measures for excluding and detecting quagga and zebra mussels from Oregon
1149 waterways. Where appropriate, work to minimize predation on sensitive native species.
1150 Where non-native aquatic species threaten SGCN, consider site-appropriate tools (e.g.,
1151 mechanical or chemical treatment) in locations and during seasons where they will not
1152 harm native amphibians, fish, or invertebrates. Educate and inform people about the
1153 problems that can be caused by invasive species, including the harm of releasing
1154 aquarium fish or nonnative fish into our rivers or dumping nonnative aquarium plants in
1155 waterbodies and the importance of having boats cleaned before moving to a different
1156 waterbody.

1157 **Limiting Factor: Passage Barriers and Channel Complexity**

1158 Fish and wildlife depend on natural flow regimes and substrates for breeding, foraging,
1159 cover, and migration. Channel complexity is important for fish and wildlife. For example,
1160 woody debris and other natural structures provide nutrient cycling and refugia from
1161 predators and high temperatures. Dams, road culverts, or other human-made barriers can
1162 restrict movement of fish and wildlife, alter or affect instream flow, and restrict bedload
1163 movement and the fluvial processes necessary to create the types of riparian and stream
1164 habitats to which native species are adapted. The large dams disrupt natural hydrologic
1165 regimes, which decreases the amount of bottomland habitat, and impacts anadromous
1166 and other migratory fish passage upstream and downstream. Additionally, altered flow
1167 regimes can contribute to higher temperatures in some streams, making habitat
1168 inhospitable.

1169 Misaligned culverts with the downstream end above the water level disconnect stream
1170 passage corridors, block fish passage, and may force wildlife to cross over roads where
1171 they are vulnerable to vehicles and predators. Undersized or improperly sized culverts can
1172 alter the transport of sediment and wood, creating an uneven distribution of instream
1173 habitat.

1174 **Recommended Approach**

1175 Work with landowners and regulatory agencies to protect and restore natural flow and
1176 channel conditions on streams impacted by barriers. **Eliminate passage barriers** or
1177 improve passage at existing barriers to provide travel corridors for fish and wildlife. Design
1178 future projects with appropriately sized culverts to accommodate adaptation to modeled
1179 hydrologic regimes with climate change. Replace culverts or other passage barriers with
1180 structures that mimic natural conditions as closely as possible (e.g., bridges or open-
1181 bottom arch culverts). Provide additional passage structures for fish and wildlife at
1182 culverts. Provide sufficient channel complexity to maintain ecological benefits for fish and
1183 wildlife.

1184 **Limiting Factor: Pollution**

1185 Point and non-point source pollution are of concern in both rural and urban areas. Non-
1186 point source pollution in flowing waters and adjacent floodplains can contain fertilizers,
1187 pesticides, or oil-based pollutants at levels high enough to cause significant lethal or sub-
1188 lethal effects in native fish and wildlife. Point source pollution from industrial, domestic,
1189 and stormwater treatment may contain high levels of contaminants. High concentrations
1190 of livestock in or near streams can degrade water quality through excessive nutrient input
1191 and may also contaminate water with bacteria. Agricultural runoff from irrigated fields also
1192 increases nutrients in streams and carries pesticides from treated fields into flowing
1193 waterways. High nutrient concentrations in streams can cause anoxic conditions,
1194 excessive aquatic vegetation, and harmful algae blooms. Pesticides in flowing waterways
1195 have the potential to damage all forms of aquatic life and may accumulate in the tissue of
1196 fish and waterfowl consumed by other wildlife and humans.

1197 **Recommended Approach**

1198 Increase awareness of the impacts of urban and rural runoff. Treat stormwater runoff that
1199 flows directly into streams to address tire-wear particles and their associated
1200 contaminants (e.g. 6PPD-q), an emerging concern in the Pacific Northwest. Reduce
1201 stormwater runoff and increase permeability in urban areas with bioswales. Use
1202 stormwater catchment basins designed for larger volume, longer residence, and a high
1203 degree of shading to mimic the delay, treatment, infiltration, and cooling functions of
1204 natural wetlands. Reduce sewage overflows during major rain events where raw sewage is
1205 discharged directly into streams. Increase awareness and manage timing of pesticide
1206 applications that have the potential to harm aquatic communities. Improve compliance
1207 with water quality standards and pesticide use labels administered by the **DEQ** and **U.S.**
1208 **Environmental Protection Agency (EPA)**. Reduce water pollution from agricultural
1209 sources and improve watershed conditions throughout the state through implementation
1210 of ODA rules and **DEQ Total Maximum Daily Load** water quality plans. Establish riparian
1211 buffer zones along streams adjacent to livestock feeding operations and farmland.
1212 Improve efficiency of irrigation systems to reduce agricultural runoff and increase instream
1213 flow. Increase interaction of rivers and floodplains. Encourage opportunities for restoration
1214 of “fringe” wetlands and channel meander to increase water storage. During restoration,
1215 remove pipes and provide stream channels to promote flow, nutrient, and oxygen
1216 exchange. Where possible, provide sufficient room to **restore meanders and other**
1217 **functions**.

1218 **Limiting Factor: Water Temperature**

1219 High water temperatures, particularly summer stream temperatures, are a major threat to
1220 self-sustaining populations of native species and can severely limit population viability for
1221 Oregon’s native anadromous and cold-water species. Aquatic animals have specific
1222 requirements for a tolerable temperature range. Moreover, warmer water holds less
1223 dissolved oxygen, resulting in hypoxic conditions. Hypoxia, which refers to low or depleted
1224 oxygen in a water body, may be lethal to organisms that extract oxygen from water, such as

1225 fish and amphibians. Water temperature may become too warm for native aquatic life
1226 because of alterations in stream flow, thermal pollution, or reduced riparian shading. This
1227 threat to native species is likely to increase with predicted regional climate change effects
1228 that include prolonged droughts, higher air temperatures, lower snowpack, and shifts in
1229 timing of rainfall and snowmelt.

1230 **Recommended Approach**

1231 Maintain or increase native riparian habitat cover to provide shading and other benefits.
1232 Maintain and restore in-stream flow to help preserve favorable water temperatures
1233 (KCI: **Water Quality and Quantity**). Advance real-time water temperature monitoring and
1234 forecasting techniques and conduct monitoring in priority areas. Identify and protect cold-
1235 water resources and refugia. Minimize release of unnaturally warm water from dams and
1236 reservoirs when instream temperatures are high by altering intake/release structures and
1237 management.

1238 **Limiting Factor: Sedimentation**

1239 Sediment flows into streams from natural processes; however, it is exacerbated through
1240 human activities. Deposition of fine sediment in gravel-bottom rivers and streams fills the
1241 interstices of the gravel, reduces the velocity of water flow through the gravel, and
1242 decreases the dissolved oxygen content. An excess of fine sediments can cover eggs of
1243 native fish and amphibians, reduce cover and protection from predators, and create
1244 adverse physical conditions. Salmonids such as salmon and trout rely on clean gravel to
1245 build redds. When fine sediment fills the spaces between gravel it prevents water from
1246 flowing through redds and oxygenating trout and salmon eggs, which reduces egg survival.
1247 In more severe cases, sediment fills the spaces between gravel and can harden the
1248 streambed rendering it unusable to spawning salmonids. Sediment can also bury aquatic
1249 mollusks and freshwater mussels. Aquatic insects rely on interstitial spaces between
1250 boulders, cobble, and gravel and many feed on periphyton that grows on these hard
1251 substrates. When sediment fills the spaces between large substrate or covers it
1252 completely, this habitat is lost, and streams can no longer support the invertebrate
1253 communities that feed fish and other wildlife.

1254 **Recommended Approach**

1255 Reduce run-off of fine sediment from logging, agriculture, grazing, roads, and other
1256 activities that could disturb soil or destabilize streambanks. Implement strategies and best
1257 management practices to reduce sedimentation including filtering run-off before it enters
1258 aquatic systems, decommissioning roads, installing green infrastructure, terracing fields,
1259 installing sediment control basins to reduce erosion, planting cover crops, and practicing
1260 conservation tillage. When constructing new roads, consider sediment removal
1261 capabilities in road design. Maintain and restore native riparian and wetland vegetation to
1262 filter sediments. Utilize large wood instream to improve stream channel complexity by

1263 increasing sediment retention, creating gravel bottom habitat, and promoting the
1264 formation of pool habitat.

1265 **Limiting Factor: Loss of Riparian Habitat, Floodplain Function, and Habitat Complexity**

1266 A large percentage of Oregon’s low-elevation and valley bottom riparian habitats have
1267 been altered or lost. Riparian habitat is often cleared, diked and converted into developed
1268 land, including urban areas, agricultural fields, or grazing pastures. Extensive removal of
1269 riparian habitat can lead to altered hydrological regimes, warmer water temperatures,
1270 erosion promoting downcutting or widening of stream banks, and loss of habitat
1271 complexity as floodplains and side channels become disconnected from streams. This
1272 loss of floodplain connectivity is a key limiting factor for nearly all listed anadromous fish
1273 species and many wildlife species. In addition, the increases in stream temperatures as a
1274 result of depleted riparian habitat often provide ideal habitat for non-native species (e.g.,
1275 game fish such as bass), allowing the non-native species to thrive and out compete native
1276 cold-water salmon and steelhead. Development within historical floodplains can restrict
1277 the natural ability of streams and riparian habitats to meander, limiting the creation and
1278 maintenance of new aquatic and riparian habitats. Lack of channel forming and flushing
1279 flows resulting from flood control efforts have also reduced floodplain processes, habitat
1280 creation, and habitat complexity. Developed floodplains are less effective in storing water
1281 and slowly releasing it back into the streams, filtering sediment and pollutants from
1282 surface water, and providing habitat for fish and wildlife. Losses of riparian habitat
1283 complexity and connectivity limit the value of these important places for wildlife to meet
1284 crucial life history needs.

1285 **Recommended Approach**

1286 Enhance or restore the extent and connectivity of existing riparian habitats. Promote lateral
1287 connectivity of the floodplain to off and side channel habitat. Use voluntary cooperative
1288 efforts and incentive programs (e.g., Conservation Reserve Enhancement Program,
1289 Riparian Lands Tax Incentive Program) to conserve, maintain, and restore riparian habitats
1290 on private lands. Identify and apply lessons learned from successful riparian restoration
1291 efforts on private lands to guide future projects. Develop tools and financial incentives to
1292 assist landowners with erosion prevention, as well as riparian area and streambank
1293 management best management practices. Provide outreach and education on the
1294 functions of riparian habitat and best management practices for landowners. This may
1295 include coordination with local governments on implementing Goal 5 protections and ODA
1296 for Agricultural Water Quality Management Area Plans.

1297 Maintain and restore riparian buffers and minimize impacts from development actions.
1298 Close and revegetate unused roads on public lands. Support and encourage beaver
1299 occupancy and their canal and dam building activities, where possible, to restore
1300 floodplain-riparian processes and function when compatible with existing land uses.
1301 Maintain channel integrity and natural hydrology. Ensure that adequate native riparian

1302 vegetation remains following management activities to prevent erosion, preserve water
1303 quality, and maintain water temperatures favorable for aquatic life. Restore lost vegetation
1304 through planting of native trees, shrubs, and ground cover. Manage for future sources of
1305 large woody debris. Maintain and/or expand existing tracts of large trees, such as
1306 cottonwoods, to benefit riparian habitat function.

1307 **Limiting Factor: Riparian Habitat Degradation**

1308 In the Blue Mountains, Northern Basin and Range, East Cascades, and Columbia Plateau
1309 ecoregions, historical overgrazing has led to soil erosion, poor regeneration of hardwood
1310 trees and shrubs, changes in plant species composition and structure, and degradation by
1311 invasive plants. Although some areas are slowly recovering, many miles of stream are still
1312 lacking adequate riparian vegetation. Ongoing grazing impacts remain in some areas,
1313 especially at low and mid elevations. Western juniper is encroaching in some riparian
1314 areas of eastern Oregon.

1315 **Recommended Approach**

1316 In cooperation with landowners, land managers, and grazing lessees, encourage
1317 approaches such as off-site watering or active management that keep livestock out of
1318 riparian areas. Develop and implement grazing regimes that are compatible with riparian
1319 conservation objectives. Selectively fence restoration sites or other high priority areas to
1320 exclude ungulates. Evaluate impacts by encroaching western juniper and remove juniper
1321 from upper reaches of higher elevation watersheds, if appropriate. Plant riparian
1322 vegetation using native species at priority sites. Work with landowners and grazing
1323 permittees to support riparian conservation and land management objectives.

1324 **Limiting Factor: Invasive Plants in Riparian Habitat**

1325 Invasive plants, such as knapweeds, knotweeds, reed canary grass, Himalayan blackberry,
1326 thistles, poison hemlock, and teasels, degrade riparian habitats by competing with and
1327 replacing native plants. In the Columbia Plateau and Northern Basin and Range
1328 ecoregions, pasture grasses and cheatgrass commonly dominate the understory. Invasive
1329 plants can alter the structure of riparian habitats, creating dense monocultures that hinder
1330 the growth of native vegetation and changing the physical structure of the streambank.
1331 They often provide insufficient food and habitat resources, displacing fish and wildlife and
1332 reducing biodiversity.

1333 **Recommended Approach**

1334 Control key invasive plants using site-appropriate tools, including fire and mechanical,
1335 biological, and chemical treatments. Use chemical treatments carefully and where
1336 compatible with water quality concerns, focusing on spot treatment during the dry season.
1337 Partner with Soil and Water Conservation Districts or other experts to control invasive

1338 weeds and restore riparian habitats. In the Columbia Plateau and Northern Basin and
1339 Range ecoregions, focus control at low-elevation sites. Provide information to local
1340 governments and landowners about potential invasive plants. Where necessary, develop
1341 and implement grazing management regimes that are compatible with riparian
1342 conservation objectives. Replace invasive plants with local native species so there is no
1343 net loss of wildlife habitat in the long term.

1344 RESOURCES FOR MORE INFORMATION

1345 OWEB's Field and Technical Guides Webpage (several guides to inform restoration and
1346 monitoring including Low Tech Process Based Restoration):
1347 <https://www.oregon.gov/oweb/resources/Pages/Field-Tech-Guidance.aspx>

1348 [Oregon Riparian Assessment Framework](#)

1349 DEQ's resources for volunteer water quality monitoring:
1350 <https://www.oregon.gov/deq/wq/Pages/WQ-Monitoring-Volunteer.aspx>

1351 [Oregon Water Resources Department's Water Conservation Tools](#)

1352 [ODFW Water Program Priorities](#)

1353 [ODA Agricultural Water Quality Plans and TMDL Implementation Plans](#)

1354 [ODEQ Total Maximum Daily Loads](#)

1355 [Beaver Created Refugia from Wildfire](#)

1356

1357 GRASSLANDS

1358 Grasslands include a variety of upland grass-dominated habitats, such as upland prairies,
1359 coastal bluffs, and montane grasslands.

1360 ECOREGIONS

1361 Grasslands are a Key Habitat in the **Blue Mountains, Coast Range, Columbia**
1362 **Plateau, Klamath Mountains, West Cascades, and Willamette Valley** ecoregions.
1363 Additional grassland habitats, such as alkali grasslands, perennial bunchgrasses, and
1364 montane grasslands, can also be found in **Specialized and Local Habitats**.

1365 CHARACTERISTICS

1366 Grasslands generally occur on dry slopes or plateaus with well-drained sandy or loamy
1367 soils. Although species vary across Oregon, perennial bunchgrasses and forbs dominate
1368 native grasslands. In some areas, grasslands are similar to wet prairies and wet meadows
1369 in structure and share some of the same prairie-associated plants and animals (wet
1370 prairies and wet meadows are included within the **Wetlands Key Habitat**). In all but the
1371 shallowest rocky soils, grasslands are maintained through disturbances, such as periodic
1372 fire, soil upheaval by rodents, frost heave, wind, or salt spray, and by humans through
1373 prescribed fire, grazing, and mowing.

1374 ECOREGIONAL CHARACTERISTICS

1375 **Blue Mountains**

1376 Bunchgrass grasslands occur primarily in the northeastern portion of the ecoregion,
1377 although other grassy habitats occur throughout the ecoregion. At low elevations, semi-
1378 desert grasslands are dominated by drought-resistant perennial bunchgrasses, such as
1379 needle-and-thread, dropseed, threeawn, and muhly, and may have scattered shrubs. Mid-
1380 elevation plateau grasslands include extensive bunchgrass prairies of Idaho fescue,
1381 junegrass, and bluebunch wheatgrass. At high elevations, ridgetop balds and alpine parks
1382 are dominated by green or mountain fescue, needlegrass, and/or bluegrass species. High-
1383 elevation grasslands often are on south-facing slopes surrounded by subalpine conifer
1384 woodlands. There are several important grassland sites currently being managed for
1385 wildlife and habitat conservation. The Zumwalt Prairie Preserve in northeast Oregon
1386 protects native bunchgrass prairie, with a portion of the reserve designated as a National
1387 Natural Landmark.

1388 **Coast Range**

1389 Coastal bluff and montane grasslands are dominated by low-growing vegetation, such as
1390 perennial bunchgrasses, forbs, mosses, and dwarf shrubs. They occur within a matrix of
1391 conifer forests. In forested ecoregions, such as the Coast Range and West Cascades,
1392 grasslands are particularly important for rare plants and invertebrates. Outer coastal bluffs
1393 and headlands are influenced by wind and salt spray, which limit the growth of woody
1394 vegetation. Montane grasslands include dry meadows and balds and occur on dry, south-
1395 or west-facing slopes with shallow sandy or gravelly soils. They are primarily influenced by
1396 periodic fire, soil upheaval by rodents, and drought conditions.

1397 **Columbia Plateau**

1398 Grasslands include river terrace grasslands, prairies, canyon slopes, and rocky ridges. At
1399 low and mid elevations, semi-desert grasslands are dominated by drought-resistant
1400 perennial bunchgrasses, such as needle-and-thread, dropseed, threeawn, and muhly, and
1401 may have scattered shrubs. Palouse grasslands once dominated most uplands above
1402 1,000 feet in elevation. Palouse grasslands now occur in flat areas with deep soils and are
1403 dominated by bluebunch wheatgrass, Idaho fescue, and other grasses and forbs. Canyon
1404 and foothill grasslands are found on the steeper, rocky slopes surrounding the major rivers
1405 in this region and are dominated by bluebunch wheatgrass, Idaho fescue, Sandberg's
1406 bluegrass, balsamroot, and other forbs.

1407 **Klamath Mountains**

1408 Grasslands in the Klamath Mountains are very diverse. They can be found on valley
1409 bottoms, and include mounded prairie often associated with vernal pools (upper Rogue
1410 Valley and Agate Desert). Dry meadow grasslands and balds occur on south and west
1411 facing mid elevation slopes of the Rogue and Umpqua basins, often in a mosaic with
1412 chaparral and oak savanna. Oak savannas are grasslands with scattered trees that are
1413 usually large with well-developed limbs and canopies. The diversity of grasslands also
1414 includes the open serpentine barrens (such as in the Illinois watershed and eastern
1415 portions of the Kalmiopsis Wilderness), and in high mountain meadows and glades of the
1416 Siskiyou mountains which are a coastal sub-range of the Klamath mountains near the
1417 Oregon/California border. The Cascade Siskiyou national monument in the southern range
1418 of the Klamath Mountains ecoregion has remaining grasslands comprised of
1419 bunchgrasses.

1420 **West Cascades**

1421 Montane grasslands include open dry meadows, grasslands, and balds. Montane
1422 grassland habitats occur in a matrix of mixed conifer forests and woodlands. Mid- and
1423 high-elevation dry meadows tend to have deeper and better-drained soils than the
1424 surrounding forests and are dominated by grasses and wildflowers, such as Roemer's
1425 fescue, alpine or western fescue, California brome, timber oatgrass, broadleaf lupine, and

1426 beargrass. Balds and bluffs generally occur on south- to west-facing slopes on shallow,
1427 well-drained soils and are dominated by bunchgrasses, forbs, and mosses.

1428 **Willamette Valley**

1429 Grasslands, also called upland prairies, are dominated by grasses, forbs, and wildflowers.
1430 Grasslands have well-drained soils and often occur on dry slopes. Willamette Valley
1431 grasslands were historically maintained by cultural burning practices. Some of the primary
1432 species include Roemer's fescue, tufted hairgrass and culturally significant species like
1433 camas, brodiaea, and madia. They are similar to wet prairies in structure and share some
1434 of the same prairie-associated plants and animals (wet prairies are included within
1435 the **Wetlands Key Habitat**). Oak savannas are grasslands with scattered Oregon white oak
1436 trees, generally only one to five trees per acre (denser oak stands are included in the **Oak**
1437 **Habitats**). Oak trees in savannas are usually large with well-developed limbs and
1438 canopies.

1439 **CONSERVATION OVERVIEW**

1440 As a whole, native grasslands are one of the most imperiled habitats in the western United
1441 States and are disappearing rapidly around the globe. In Oregon, the estimated loss of
1442 grasslands ranges from 50 percent to more than 90 percent, depending on the ecoregion.
1443 Compared to historical grassland distributions, grassland loss has been extremely high in
1444 in valley bottoms and foothills in the Coast Range, West Cascades, and Willamette Valley
1445 ecoregions. These historical grasslands have been impacted by conversion to agriculture,
1446 development, succession to forested habitats, and invasive plant species. The deep soils
1447 and moderate climates of many grassland habitats make them especially valuable for
1448 agricultural land uses such as crop, hay, or pasture lands. Areas with deep soil were
1449 disproportionately lost to agricultural cultivation while areas with shallower soils were
1450 more likely to experience intensive grazing. Chronic grazing has impacted grasslands,
1451 affecting plant composition and structure. Also, non-native species were historically
1452 seeded for livestock forage in some grasslands, decreasing the abundance and diversity of
1453 native plants. However, grazing practices have become more sustainable over time, and
1454 carefully managed grazing can help to maintain grassland structure where prescribed fire
1455 is not practical or desired. Disruption of historical fire regimes has allowed for shrubs or
1456 trees to encroach, replacing grasslands with forest. In addition, some foothill grasslands
1457 have been converted to forests through tree planting.

1458 As human population growth increases, urbanization may present a significant challenge
1459 for grassland habitats. While agricultural areas may still mimic some grassland structure
1460 and function and retain some value for wildlife, development and urbanization results in
1461 the direct loss of habitat and habitat fragmentation.

1462 Species of Greatest Conservation Need (SGCN) associated with grasslands vary by
1463 ecoregion but include the: **Burrowing Owl**, **Common Nighthawk**, **Grasshopper**

1464 **Sparrow, Long-billed Curlew, Ferruginous Hawk, Oregon Vesper Sparrow, Streaked**
1465 **Horned Lark, Western Bluebird, Western Meadowlark, Fender's blue butterfly, hoary**
1466 **elfin butterfly, Kincaid's lupine, Oregon silverspot butterfly, Taylor's checkerspot**
1467 **butterfly, Coast Range fawn lily, Cascade Head catchfly, Lawrence's**
1468 **milkvetch, Spalding's campion, and Tygh Valley milkvetch.**

1469 LIMITING FACTORS AND RECOMMENDED APPROACHES

1470 **Limiting Factor: Altered Fire Regimes**

1471 At sites with deep soils, maintenance of grasslands is dependent, in part, on periodic fire.
1472 Fire suppression has led to encroachment by shrubs and conifer trees in some areas and
1473 has aided in an increase in fuel loads, which can lead to high-intensity wildfires. The
1474 introduction and rapid spread of cheatgrass and other non-native grasses throughout
1475 eastern Oregon can increase the frequency, intensity, and spread of fires. In the
1476 Willamette valley in particular, grasslands and the species that inhabit them, are
1477 dependent on managed fire due to coevolution with cultural burning practices. In the
1478 Coast Range, prescribed fire is difficult due to high precipitation and wet conditions. When
1479 conditions are dry enough to use prescribed fire, there may be concerns about risk to
1480 surrounding forests. In the Klamath Mountains and Willamette Valley, prescribed fire
1481 poses challenges, such as conflicts with surrounding land use, smoke management and
1482 air quality, and safety.

1483 **Recommended Approach**

1484 Maintain open grassland structure by using multiple site-appropriate tools, such as
1485 prescribed burns, mowing, controlled grazing, hand-removal of encroaching shrubs and
1486 trees, or thinning. Re-introduce fire at locations and at times where conflicts, such as
1487 smoke and safety concerns, can be minimized. Work with partners to update smoke
1488 management and air quality standards to allow more fall, winter, and spring burn windows.
1489 For all tools, minimize ground disturbance and impacts to native species. Minimize the
1490 spread of cheatgrass. Carefully manage livestock grazing to maintain native plants and soil
1491 crust (cryptogammic crust). Control fires in cheatgrass-dominated areas.
1492 (KCI: **Disruption of Disturbance Regimes**)

1493 **Limiting Factor: Invasive Species**

1494 Invasive plants have degraded grassland habitats, displacing native plants and animals.
1495 Some intentionally planted non-native species, such as crested wheatgrass, are highly
1496 competitive with native bunchgrasses and, once established, limit the growth and
1497 establishment of native plants. Depending on the area, invasive species include
1498 cheatgrass, medusahead, ventenata, rush skeleton weed, spikeweed, Hungarian brome,
1499 yellow star-thistle, knapweeds (diffuse, spotted, and purple), leafy spurge, Canada thistle,
1500 St. John's wort, tansy ragwort, Armenian (Himalayan) blackberry, evergreen blackberry,

1501 Scotch broom, false-brome, Harding grass, and tall oatgrass. Many low-elevation
1502 grasslands are almost entirely dominated by invasive grasses, forbs, and shrubs. In the
1503 Blue Mountains and the Columbia Plateau, juniper encroachment has displaced
1504 grasslands in many areas. Disturbed sites are especially prone to invasive species
1505 establishment.

1506 **Recommended Approach**

1507 Identify remaining native grasslands and work with landowners to maintain quality and
1508 limit the spread of invasive species. Emphasize prevention, risk assessment, early
1509 detection, and quick control to prevent new invasive species from becoming fully
1510 established. To control encroaching junipers, use mastication, cut and pile, lop and
1511 scatter, or cutting for firewood. Develop markets for small juniper trees as a special forest
1512 product to reduce restoration costs. Prioritize control efforts and use site-appropriate
1513 methods to control newly established invasive plant species for which management can
1514 be most effective. Promote the development of additional native seed resources. Re-seed
1515 with site-appropriate native grasses and forbs after control efforts. Conduct research to
1516 determine methods to manage established species, such as cheatgrass, medusahead,
1517 Hungarian brome, and annual ryegrass. Where appropriate, manage livestock grazing and
1518 recreational use, especially motorized use, to minimize new introductions. Support
1519 current prevention programs, such as weed-free hay certification (KCI: **Invasive Species**).
1520 Clean vehicles and other equipment when relocating between sites where invasive species
1521 are present. Establish and implement management plans for all soil-disturbing activities.

1522 **Limiting Factor: Land Use Conversion**

1523 Remnant grasslands are subject to conversion to agricultural, residential, urban, energy,
1524 and infrastructure uses. Grasslands are frequently converted into croplands or
1525 pasturelands because of their deep soils and high productivity. The expansion of cities and
1526 towns often includes converting grasslands into urban and rural residential areas and
1527 associated infrastructure. The conversion of grasslands into other land uses results in
1528 habitat loss and fragmentation, degradation of ecosystem services, such as carbon
1529 sequestration, and biodiversity loss.

1530 **Recommended Approach**

1531 Because many of these areas are privately-owned, **voluntary cooperative**
1532 **approaches** are the key to long-term conservation. Important tools include financial
1533 incentives, technical assistance, regulatory assurance agreements, and conservation
1534 easements. Use and extend existing incentive programs, such as the Conservation
1535 Reserve Program and Grassland Reserve Program, to conserve, manage, and restore
1536 grasslands and to encourage no-till and other compatible farming practices. Support and
1537 implement existing **land use regulations** to preserve natural habitats. Use a landscape
1538 approach in conservation plans and incentive programs to create large, contiguous blocks

1539 of grassland habitat by expanding buffers around key grassland sites. Connect grassland
1540 habitats, such as fallow fields, pastures, and natural meadows, to create contiguous
1541 grassland habitat and improve connectivity between patches.

1542 **Limiting Factor: Land Management Conflicts**

1543 Resource conflicts can arise because high quality grasslands are often high-quality grazing
1544 resources. Although grazing can be compatible with conservation goals, it needs to be
1545 managed carefully because Oregon’s bunchgrass habitats are more sensitive to grazing
1546 than the sod-forming grasses of the mid-western prairies. Overgrazing can lead to soil
1547 erosion, degradation of biological soil crusts, changes in plant species composition and
1548 structure, and establishment of invasive plants. Agricultural management practices, such
1549 as mowing, haying, burning, and herbicide/insecticide application, can be detrimental to
1550 grassland species.

1551 **Recommended Approach**

1552 Use incentive programs and other voluntary approaches to manage and restore grasslands
1553 on private lands. Manage public land grazing to maintain grasslands in good condition.
1554 Conduct research and develop incentives to determine grazing regimes that are
1555 compatible with a variety of conservation goals. Restore native grassland habitat when
1556 possible, removing woody growth and invasive weeds to create a mosaic of clumped
1557 vegetation, bare ground, and a mixture of grasses and forbs with a variety of heights.
1558 Promote use of native plants and seed sources in conservation and restoration programs.
1559 Promote operation of grassland management practices (e.g., mowing, haying, burning, and
1560 herbicide application) to outside of the primary breeding season for grassland-associated
1561 wildlife (roughly April-August).

1562 **Limiting Factor: Loss of Oak Savannas**

1563 In the Klamath Mountains and Willamette Valley ecoregions, large-diameter oak trees with
1564 lateral limb structure and cavities continue to be lost. **Oak Habitats** complement
1565 grassland habitat and should be maintained. Many native wildlife species utilize large-
1566 diameter oaks for nesting, feeding, and shelter. Prior to European settlement, cultural
1567 burning practices helped to maintain the open structure of widely spaced, large-crowned
1568 trees with an understory of perennial native grasses and forbs.

1569 **Recommended Approach**

1570 Maintain large oaks, remove competing conifers or densely stocked small oaks, and create
1571 snags to provide cavity habitat. Management practices like prescribed fires, controlled
1572 grazing, or mowing can maintain oak savanna conditions and help to control invasive
1573 species and encroaching woody vegetation.

1574 HABITAT CHANGE TRENDS ANALYSIS

1575 To investigate juniper encroachment into grassland and sagebrush habitats, the Institute of
1576 Natural Resources (INR) compared the total area and spatial overlap of vegetation classes
1577 in three baseline maps (1851-1937, 1998, 2016). The analysis showed significant increases
1578 in the total area of Juniper Woodlands and corresponding losses of Grassland habitats to
1579 juniper encroachment. By 2016, the total area of Juniper Woodlands had increased by
1580 115% when compared to historical data, largely replacing Sagebrush and Grassland
1581 habitats. An estimated 15% of Grassland habitats were lost to juniper encroachment
1582 between 1851 and 1998. This analysis also showed that significant efforts by land
1583 managers and agencies, such as the BLM and Forest Service, to remove juniper have
1584 slowed encroachment in grassland habitats in recent years.

1585 To address concerns regarding annual grass invasion of both Grassland and Sagebrush
1586 Habitats, INR conducted an analysis using the Rangeland Analysis Platform (RAP) annual
1587 vegetation cover maps to track the total area that is dominated by annual grasses in Oregon
1588 in 1986, 2001, 2016, and 2023. The analysis documented significant increases in annual-
1589 dominated vegetation since 1986. By 2001, the total area of annual-dominated vegetation
1590 increased by 118%. Between 2001 and 2016, the total area of annual-dominated
1591 vegetation increased by an additional 69%. In all years, annual-dominated vegetation was
1592 mostly found in the grassland and sagebrush habitats of southeast Oregon and in the
1593 Columbia Basin ecoregion.

1594 RESOURCES FOR MORE INFORMATION

1595 **[Prairie Vegetation Monitoring Protocol for the North Coast and Cascades Network](#)**

1596 **[The Willamette Valley Landowner's Guide to Creating Habitat for Grassland Birds](#)**

1597 **[Partners in Flight Conservation Strategy for Landbirds in Lowlands and Valleys of](#)**
1598 **[Western Oregon and Washington](#)**

1599 **[Benton County Prairie Species Habitat Conservation Plan](#)**

1600 [Cascadia Prairie-Oak Partnership](#)

1601 [Restoring Oak Habitats in Southern Oregon and North California: A Guide for Private](#)
1602 [Landowners](#)

1603 [Patterns of Vegetation Change in Grasslands, Shrublands, and Woodlands of](#)
1604 [Southwest Oregon](#)

1605 REFERENCES

1606 Allred, B.W., B.T. Bestelmeyer, C.S. Boyd, C. Brown, K.W. Davies, M.C. Duniway, L.M.
1607 Ellsworth, T.A. Erickson, S.D. Fuhlendorf, T.V. Griffiths, V. Jansen, M.O. Jones, J. Karl, A.
1608 Knight, J.D. Maestas, J.J. Maynard, S.E. McCord, D.E. Naugle, H.D. Starns, D. Twidwell,
1609 and D.R. Uden. 2021. Improving Landsat predictions of rangeland fractional cover with
1610 multitask learning and uncertainty. *Methods in Ecology and Evolution*.

1611 Brunner, R. and E. Gaines. 2025. Oregon Vegetation Change 1851-2023. Trends analysis
1612 conducted for Oregon Department of Fish and Wildlife. Institute for Natural Resources,
1613 Portland State University, Portland, OR, USA.

1614

DRAFT

1615 LATE SUCCESSIONAL MIXED CONIFER FORESTS

1616 Late successional mixed conifer forests provide a multi-layered tree canopy, including
1617 large-diameter trees, shade-tolerant tree species in the understory, and a high volume of
1618 dead wood, such as snags and logs.

1619 ECOREGIONS

1620 Late successional mixed conifer forests are a Key Habitat in the **Blue Mountains, Coast**
1621 **Range, East Cascades, Klamath Mountains, and West Cascades** ecoregions.

1622 CHARACTERISTICS

1623 Late successional mixed conifer forests are defined by plant species composition,
1624 overstory tree age and size, and the forest structure. While a range of tree ages are present
1625 in late successional forests, the predominant stand age is over 150 years. They include
1626 characteristics such as a multi-layered tree canopy, shade-tolerant tree species growing in
1627 the understory, large-diameter trees, and a high volume of dead wood, such as snags and
1628 logs. These characteristics can take hundreds of years without stand replacing
1629 disturbance to develop. Historically, fire was the major natural disturbance in all but the
1630 wettest areas. Depending on local conditions, fires in the Coast Range and West Cascades
1631 conifer forests were of moderate- to high-severity, with fire return intervals averaging 100 to
1632 more than 400 years. These stand replacing events were interspersed with periodic low
1633 severity understory burns every 15 to 30 years. This historical fire regime created a complex
1634 mosaic of stand structures across the landscape.

1635 ECOREGIONAL CHARACTERISTICS

1636 **Blue Mountains**

1637 A mixture of conifer species is found throughout many forest sites in the Blue Mountains
1638 ecoregion. Mixed conifer forests can be divided in two subtypes based on temperature and
1639 moisture conditions. Douglas-fir and grand fir are the primary late successional tree
1640 species in the warmer drier climates of this ecoregion. Ponderosa pine and western larch
1641 may also be present. The cool mixed conifer type is indicated by the addition of more
1642 moisture-demanding and cold-tolerant species, such as subalpine fir and Engelmann
1643 spruce, at upper elevations or along streams where cold water-drainage and deep frost
1644 eliminate some species. The understory in this ecoregion generally includes huckleberry,
1645 serviceberry, oceanspray, snowberry, wild ginger, goldthread, starflower, bead lily, and oak
1646 fern.

1647 **Coast Range**

1648 Late successional mixed conifer forests in the Coast Range are generally dominated by
1649 two types of conifer trees: Sitka spruce and Douglas-fir. Sitka spruce forests occur within a
1650 narrow fog- and salt-influenced strip along the coast and extend into some valleys. Soils
1651 tend to be deep, acidic, and well-drained. Sitka spruce dominates the overstory, but
1652 western hemlock, western redcedar, Douglas-fir, big leaf maple, and red alder may be
1653 present. The lush understory has salmonberry, vine maple, salal, evergreen huckleberry,
1654 sword fern, deer fern, and a high diversity of mosses and lichens. Inland, Douglas-fir
1655 forests dominate. The understory of Douglas-fir forests includes shrub and forb species,
1656 such as vine maple, salal, sword fern, Cascade Oregon grape, western rhododendron,
1657 huckleberries, twinflower, vanilla leaf, and oxalis. Due to high precipitation in both Sitka
1658 spruce and Douglas-fir forests, fires are infrequent but do occur during hot, dry, east wind
1659 conditions after prolonged drought. When fires do occur, they are likely to be high severity,
1660 stand-replacing events. Other disturbances include small-scale windthrow events and
1661 floods driven by atmospheric river storms.

1662 **East Cascades**

1663 Late successional mixed conifer forests span the eastern slopes of the Cascade
1664 Mountains. This habitat contains a wide variety of tree species with Douglas-fir, grand fir,
1665 and western hemlock as the most common forest tree species that co-dominate most
1666 overstories. Several other conifers may also be present, including western redcedar,
1667 western white pine, western larch, ponderosa pine, and lodgepole pine. Undergrowth
1668 vegetation in the East Cascades ecoregion includes vine maple, Oregon grape,
1669 huckleberry, oxalis, thimbleberry, manzanita, ceanothus, and twinflower. Many sites once
1670 dominated by Douglas-fir and ponderosa pine (formerly maintained by wildfire) may now
1671 be dominated by grand fir (a fire sensitive, shade-tolerant species).

1672 **Klamath Mountains**

1673 Late successional mixed conifer forests in the Klamath Mountains ecoregion are
1674 characterized by high tree diversity. Douglas-fir is usually dominant. Depending on site
1675 characteristics, other canopy trees may include white fir, sugar pine, ponderosa pine, and
1676 incense cedar. Port-Orford cedar occurs on moist sites, such as riparian areas. Jeffrey pine
1677 and knobcone pine occur on serpentine soils. Broadleaf trees, such as tanoak, canyon live
1678 oak, golden chinquapin, big leaf and vine maple, and Pacific madrone, may occur in the
1679 subcanopy. A range of understory communities may be present, including those mostly
1680 dominated by shrubs, forbs, or grasses, or may be relatively open. However, with an
1681 increase in frequency of droughts, high instances of Douglas-fir mortality in the Klamath
1682 Mountains is quickly shifting the composition of these forests, creating conditions that
1683 result in catastrophic wildfires.

1684 **West Cascades**

1685 Late successional mixed conifer forests are found scattered throughout the **West**
1686 **Cascades Ecoregion**. While Douglas -fir dominates these forests, western hemlock is
1687 almost always co-dominant and usually dominates the understory. In the absence of stand
1688 replacing disturbance, Douglas-fir forests eventually convert to western hemlock. Other
1689 common trees include grand fir and western redcedar in the northern portion of the
1690 ecoregion, or incense cedar, sugar pine, white fir, and western redcedar in the southern
1691 portion of the ecoregion. The understory has shrub and forb species, such as vine maple,
1692 salal, sword fern, Cascade Oregon grape, western rhododendron, huckleberries,
1693 twinflower, vanilla leaf, and oxalis.

1694 CONSERVATION OVERVIEW

1695 Since the 1850s, both timber harvest and large-scale fires have replaced many of the late
1696 successional forests in Oregon with younger forests. Many of the remaining late
1697 successional forests occur in a patchwork with the younger forests that are managed with
1698 shorter rotations to generate timber products. While younger forests still hold value for fish
1699 and wildlife, late successional forests support a wide array of species, many of which
1700 require large patches of older or mature forests to survive and may be sensitive to changes
1701 in the forest seral stage.

1702 The Northwest Forest Plan (**NWFP**) is a comprehensive natural resource planning effort
1703 that includes all or part of the Siuslaw, Rogue River-Siskiyou, Mt. Hood, Willamette,
1704 Deschutes, Umpqua, and Fremont Winema National Forests in Oregon. The
1705 NWFP identifies conservation priorities for species affected by loss and fragmentation of
1706 large patches of late successional forests, assessing over 1,000 species. Late
1707 Successional Reserves established under the NWFP are intended to ensure enough high-
1708 quality habitat to sustain identified species. However, many of the federal lands that are
1709 designated as Late Successional Reserves do not include forests at the late successional
1710 stage, while others are relatively small “checkerboards” of forests embedded in a matrix of
1711 private industrial timber lands, particularly in the Coast Range and Klamath Mountains.
1712 From 1994 until 2020, there was an increase in the acres of late successional stage forest
1713 in the NWFP area. However, in 2020, wildland fires driven by strong east winds in the Mt.
1714 Hood, Willamette, Umpqua, and Rogue Siskiyou National Forests destroyed many acres of
1715 late successional forest. While the 2020 fires were a setback, the gains of 1994 to 2020
1716 demonstrated that the NWFP can successfully increase late successional mixed conifer
1717 forests over time.

1718 The US Forest Service’s Wildfire Crisis Strategy Implementation Plan and Oregon
1719 Department of Forestry’s 20 Year Landscape Resiliency Strategy are examples of federal
1720 and state efforts to address uncharacteristic wildland fire behavior in Oregon’s forests.
1721 These plans address historical fire suppression, exclusion of cultural burning practices,
1722 and the impacts of recent catastrophic and uncharacteristic wildfires by recommending a
1723 variety of active management techniques for forests to increase forest resiliency to
1724 wildland fire.

1725 The Northwest and Southwest State Forest Management Plans provide management
1726 direction for all Board of Forestry Land and Common School Forest Lands. The plans
1727 include management strategies for 16 resources, including fish and wildlife, timber,
1728 recreation, and water resources. The plans describe long-term desired future conditions,
1729 which include older forest structure.

1730 The Private Forest Accord (PFA) is a compromise agreement made between representatives
1731 from Oregon's timber industry, the Oregon Small Woodlands Association, and prominent
1732 conservation and fishing organizations, to modify portions of Oregon's forest practice laws
1733 and regulations in a way that expands protections for fish and amphibians. The changes to
1734 the Oregon Forest Practices Act are aimed to avoid and minimize the effects that timber
1735 harvests and other forest management activities on private forestlands have on these
1736 species and the aquatic habitats they depend on. The PFA included new standards for
1737 stream classification and protection or stream buffers, steep slopes, roads, and culverts,
1738 as well as a grant program to fund riparian and stream habitat restoration projects. These
1739 standards may also help to restore some late successional mixed conifer forests located in
1740 riparian areas adjacent to streams.

1741 In the southwest Cascades, white firs in overstocked stands are often full of disease,
1742 creating a buildup of fuels and putting large ponderosa pines at risk for severe wildfire.
1743 These ladder fuels are a product of fire suppression, exclusion of cultural burning
1744 practices, and past logging practices. Removing these patches will help create more
1745 early seral openings for the benefits of deer and elk and provide a greater mosaic of
1746 habitats across the landscape.

1747 Late successional mixed conifer forests are particularly important for wildlife, mosses, and
1748 lichens. Depending on the ecoregion, **Species of Greatest Conservation**
1749 **Need** associated with late successional conifer forests include **ringtail, fisher, Pacific**
1750 **marten, red tree vole, Marbled Murrelet, Northern Spotted Owl, Oregon slender**
1751 **salamander**, and many others.

1752 **[Spotlight] Pileated Woodpecker**

1753 *The forest's engineer in the Pacific Northwest*

1754 The Pileated Woodpecker plays a vital role in the health and biodiversity of the Pacific
1755 Northwest's mature forests. As the largest woodpecker in the region, it is an ecological
1756 powerhouse that helps shape the forest around it.

1757 Pileated Woodpeckers are expert excavators, carving out large rectangular holes in old
1758 decaying trees to find food and create cavities for nests and roosts. This in turn can create
1759 habitat for secondary cavity nesters/roosters like Pacific marten, Flammulated Owls,
1760 silver-haired bats, of Vaux's Swifts. For many species, availability of cavities for nesting
1761 and roosting is a limiting factor.

1762 Additional impacts of Pileated Woodpecker activity include the creation of foraging
1763 opportunities for other species, acceleration of decomposition and nutrient cycling,
1764 increased heart rot fungi growth and inoculation, and mediation of damaging insect
1765 outbreaks.

1766 The Pileated Woodpecker was included in the first two iterations of Oregon's State Wildlife
1767 Action Plan as a management indicator species of mature and old growth habitats.
1768 Pileated Woodpeckers were once considered an indicator of old growth forests in the
1769 Pacific Northwest, but studies of habitat use and preference suggest that they are more
1770 accurately indicators of the structural elements that are characteristic of mature or old
1771 growth forests, or forestry practices that ensure older trees are retained. While
1772 populations of Pileated Woodpecker in Oregon are currently secure, conservation and
1773 management of this species can provide broad positive impacts to species that rely on
1774 mature forests with these characteristics.

1775 **LIMITING FACTORS AND RECOMMENDED APPROACHES**

1776 **Limiting Factor: Loss of Structural Habitat Elements**

1777 Where historical stands were perpetuated for 200 to more than 1,000 years, commercial
1778 forestlands are now commonly harvested every 60 years or less, which limits the
1779 maintenance and future recruitment of late-successional characteristics. In addition, the
1780 number of large-diameter snags and logs, which contribute to understory structure, has
1781 been reduced over time through wildfire and timber harvest.

1782 **Recommended Approach**

1783 Develop programs, incentives, and market-based approaches to encourage longer
1784 rotations and strategically located large-diameter tree tracts. Where feasible, maintain
1785 structural elements, such as large-diameter tall trees, snags, and logs. Create snags from
1786 green trees or high-cut stumps where maintaining snags is not feasible or where snag
1787 management goals are not being met. Maintain forest stand structures on private industrial
1788 forest lands, and provide technical assistance to landowners to leave large-diameter
1789 downed wood, green trees, or snags in the upland portion of harvested forests, as well as
1790 along riparian areas, to provide benefits for a diversity of wildlife and fish.

1791 **Limiting Factor: Loss of Late Successional Stand Size and Connectivity**

1792 Late successional forest stands have been greatly reduced in size and connectivity,
1793 particularly at lower elevations. This can impact species that are highly adapted to late
1794 successional conditions, require large tracts of intact habitats, and/or species that have
1795 limited ability to move over long distances to find new suitable areas. It also allows edge
1796 species to compete with those adapted to extensive interior forest habitat.

1797 **Recommended Approach**

1798 Maintain existing plans to protect and develop late successional habitat. Use active
1799 management to accelerate development of late successional structural characteristics in
1800 key areas to expand existing patches into larger areas; these will provide greater blocks of
1801 habitat for species with large area requirements or those that require interior forest habitat
1802 and are vulnerable to “edge effects”. Continue to carefully plan forest practices to
1803 maintain connectivity (KCI: **Barriers to Animal Movement**), particularly when species
1804 vulnerable to fragmentation are present. ODFW has mapped **Priority Wildlife**
1805 **Connectivity Areas (PWCAs)** to provide information on places across the landscape with
1806 the highest overall value for facilitating wildlife movement.

1807 Seek opportunities to coordinate management of public and private lands (e.g., All-Lands
1808 Approach) whenever possible to address conservation needs. Use voluntary conservation
1809 tools, such as financial incentives and forest certification to achieve conservation goals on
1810 private lands. Work to maintain a diversity of forest types and ages to support wildlife
1811 habitat connectivity and ecosystem services at a landscape scale.

1812 **HABITAT CHANGE TRENDS ANALYSIS**

1813 Following disturbance, such as timber harvest, regenerating conifer forests often succeed
1814 to deciduous or mixed deciduous-conifer forests without active management (e.g.
1815 replanting with Douglas-fir), especially on the west side of the state. These deciduous
1816 forests are primarily dominated by red alder and bigleaf maple. To investigate the transition
1817 from late successional mixed conifer forests to deciduous forest, the Institute of Natural
1818 Resources (INR) analyzed the change in total area of west-side deciduous and mixed
1819 conifer-deciduous forests between 1851 and 2016. The analysis showed a massive 243%
1820 increase in deciduous forests between 1851 and 1998, likely reflecting a shift towards
1821 earlier successional forests following logging.

1822 **RESOURCES FOR MORE INFORMATION**

1823 [Northwest Forest Plan](#)

1824 [Status and Trends of Late Successional and Old Growth Forests](#)

1825 [Oregon Private Forest Accord](#)

1826 **REFERENCES**

1827 Brunner, R. and E. Gaines. 2025. Oregon Vegetation Change 1851-2023. Trends analysis
1828 conducted for Oregon Department of Fish and Wildlife. Institute for Natural Resources,
1829 Portland State University, Portland, OR, USA.

1830 Hagman, R.K. et. al. 2021. Evidence for widespread changes in the structure, composition,
1831 and fire regimes of western North American forest. Ecological Applications. Volume 31,
1832 Issue 8.

1833

DRAFT

1834 NATURAL LAKES

1835 Natural lakes are relatively large bodies of freshwater surrounded by land that were formed
1836 through geological processes, such as glacial scouring, tectonic movements, volcanic
1837 activity and river meander cutoffs. In Oregon, natural lakes are defined as standing water
1838 bodies larger than 20 acres, including some seasonal lakes. Depth is not a reference for
1839 characterization of a natural lake.

1840 ECOREGIONS

1841 Natural Lakes are identified as a Key Habitat in all inland ecoregions.

1842 CHARACTERISTICS

1843 Natural lakes are distributed throughout Oregon, although the highest concentrations and
1844 largest lakes are found in the **West Cascades**, **East Cascades**, and **Northern Basin and**
1845 **Range** ecoregions. Sources of water for Oregon's natural lakes include rainfall, snowmelt,
1846 seeps and stream flows. The diversity of natural lakes is reflected in the processes that
1847 formed them. These processes include glaciation, volcanism (calderas and lava flows),
1848 coastal dune impoundment, faults and rifts, and riverine erosion (oxbow lakes). Natural
1849 lakes provide important habitat for **Species of Greatest Conservation Need** (SGCN),
1850 contribute to ecosystem services, and attract visitors for tourism and recreation year-round
1851 throughout Oregon's communities.

1852 Crater and Waldo Lakes, Oregon's largest clear water lakes, are both located in the West
1853 Cascades ecoregion, and have been designated by the Department of Environmental
1854 Quality (DEQ) as Outstanding Resource Waters. The designation provides special
1855 protections to maintain the exceptional water quality, ecological, cultural and recreation
1856 values of these lakes. Clear Lake in the McKenzie basin is another naturally clear and cold
1857 lake in the Cascades, with a maximum depth of 175 feet. There are many volcanic lakes in
1858 the Cascade Mountain Range that are also notably clear, supporting diverse aquatic life by
1859 allowing light to penetrate deeper and enhancing their aesthetic value.

1860 The eastern half of the state contains several playa lakes, formed when runoff from
1861 precipitation and mountain snowpack flows into low-lying areas, then evaporates and
1862 leaves mineral deposits. Playas are valuable for their role in water storage, groundwater
1863 recharge, and as critical habitat for migratory birds and other wildlife. In south central
1864 Oregon, Lake Abert has provided essential habitat and food sources for a myriad of
1865 migrating birds in the Pacific Flyway. Lake Abert is also the only hypersaline lake in Oregon
1866 and one of only three hypersaline lakes in the United States.

1867 The Natural Lakes Key Habitat does not include irrigation ditches, reservoirs, or other man-
1868 made water bodies. The wet zone and riparian zones around the edges of natural lakes is
1869 mapped as **Wetlands Key Habitat**.

1870 CONSERVATION OVERVIEW

1871 Many of Oregon's larger natural lakes are important destinations for tourism and
1872 recreation, especially in the summer, and many are desirable locations for year-round
1873 commercial and residential development. These uses can have direct impacts on water
1874 quality and quantity. Pollution and sedimentation are also concerns associated with
1875 development and recreation. Once established, invasive plants and animals can dominate
1876 natural lake environments, reducing biodiversity and impacting recreational activities.
1877 Rising temperatures and altered precipitation regimes associated with climate change will
1878 affect water levels and ecosystem health. In eastern Oregon and at higher elevations, rising
1879 temperatures from climate change are affecting the ice regime (the dates that ice freeze
1880 and thaw) which impacts access to the lake to feed and to drink water. Some of Oregon's
1881 lakes contain unique assemblages of species and habitat features that have high
1882 conservation value. For example, many amphibian and fish SGCN rely upon Oregon's lakes
1883 for breeding each year.

1884 LIMITING FACTORS AND RECOMMENDED APPROACHES

1885 **Limiting Factor: Drought**

1886 Many lakes in south central and southeast Oregon have been drying naturally since the last
1887 glacial period. The effect of this natural drying process is being exacerbated by
1888 anthropogenic climate change. Droughts have resulted in a change in precipitation
1889 patterns, loss of natural runoff, and increase in evapotranspiration from natural lakes.
1890 Increased demand for out-of-stream water use is associated with prolonged drought, and
1891 ground water pumping and management are drying out some natural lakes, especially in
1892 south central and southeastern Oregon. These dry or diminished lake beds impact
1893 waterfowl, recreation, water availability, water quality, aesthetics and human health. Lake
1894 Abert has been affected by many drought-related factors including reduced water flow into
1895 the lake, increased salinity, impacts to food sources for the migrating waterfowl, and
1896 decreased habitat and biodiversity. In Crater Lake, scientists have observed an increase in
1897 the lake's temperature, a decrease in snowfall, and impacts on species such as American
1898 pika and whitebark pine. Drought limits water availability for recreation, fish and wildlife,
1899 and human use and puts stress on the fish and wildlife species that depend on these lake
1900 systems.

1901 **Recommended Approach**

1902 Monitor and measure stream flows, groundwater levels, lake levels, and water use.
1903 Implement modern technology to monitor water use from natural lake systems. Where
1904 applicable, work toward improving irrigation efficiency and delivery systems to use less
1905 water while protecting agricultural interests. Develop and implement groundwater
1906 management plans. Develop and adopt instream water rights for high priority natural lakes
1907 and upstream river reaches contributing flow. Continue to support collaborative water
1908 management solutions to identify, develop and implement voluntary projects that result in
1909 reliable water supplies to natural lakes in late summer, when water quality impacts are the
1910 highest. Manage water in the state with the long-term health and sustainability of natural
1911 lakes as a goal. Continue to educate Oregonians about conscientious water use.

1912 **Limiting Factor: Water Quality**

1913 Nonpoint source pollution may contain fertilizers, pesticides, or oil-based contaminants at
1914 levels high enough to cause significant lethal or sub-lethal effects in native fish and
1915 wildlife. Nonpoint source pollution can enter lakes through runoff from surrounding lands
1916 or streams, and from groundwater. In some lakes, use of motorized recreational watercraft
1917 can degrade **water quality** through pollution.

1918 Cyanobacteria blooms have become more common and prevalent in natural lakes and
1919 waterways. Cyanobacteria can produce cyanotoxins that can cause serious illness or
1920 death in pets, livestock and wildlife. These toxins can also make people sick, and in
1921 sensitive individuals, cause a rash or skin, ear and eye irritation. The frequency, extent, and
1922 magnitude of harmful algal blooms in waterbodies is a response to a variety of individual
1923 and combinations of factors, including changes in water temperature, nutrient loading, and
1924 hydrologic conditions within watersheds.

1925 During wildfires, ash can enter the natural lake and increase heavy metal concentrations in
1926 the water. Ash adds nutrients like nitrogen and phosphorus into the ecosystem. Algae feed
1927 on these nutrients and the nutrients and algae affect the food chain and the clarity of the
1928 lake water.

1929 **Recommended Approach**

1930 The diffuse, intermittent nature of nonpoint source pollutants make traditional
1931 management and control of pollutants very challenging. One of the most effective means
1932 of controlling nonpoint source pollutants is through education and the regulation of land
1933 use and associated land management practices. When recreating, carefully consider
1934 recreational vehicle use and timing of use in sensitive water bodies. Minimize use of
1935 pesticides and herbicides, and follow pesticide use labels for proper application.
1936 Implement Agricultural Water Quality and DEQ Total Maximum Daily Load water quality

1937 plans. Plant vegetation around driveways, shorelines and on slopes, so the vegetation can
1938 absorb nutrients, filter out pollutants, and trap sediment. Maintain septic tanks with regular
1939 pumping and inspection at least every 3-5 years. Pick up pet waste and dispose of it in the
1940 trash. Educate recreational users of projected and actual cyanobacterial blooms in natural
1941 lakes and waterbodies.

1942 **Limiting Factor: Habitat Loss**

1943 Habitat loss has occurred in natural lakes from residential housing, shoreline
1944 development, energy development, agriculture, and infrastructure development. This
1945 results in the loss of natural buffer zones, which are essential for filtering sediments and
1946 nutrients, stabilizing shorelines by slowing water flow, reducing erosion, and providing
1947 critical habitat for various fish and wildlife species by offering food sources and shelter
1948 along lake edges. Water withdrawals (both surface and groundwater), water diversion, and
1949 drought can significantly alter the natural flow and levels of lakes, impacting the availability
1950 of lake habitats.

1951 **Recommended Approach**

1952 Provide outreach and education on avoidance and minimization of impacts from
1953 development actions. This may include limiting development, including residential
1954 housing, road and rail placement along the shoreline of natural lakes to protect fringe
1955 wetlands that buffer the lake, or utilizing a raised trail to maintain hydrology. Support
1956 managed public access to lakes to promote self-education and advocacy of natural
1957 habitats. Limit the footprint of docks and boathouses on natural lakes, follow ODFW
1958 Residential Dock Guidelines, or use established public ramps. Protect native, intact habitat
1959 along the shoreline. Where possible, remove dikes along the shoreline of natural lakes or
1960 modify dike location and structure to restore fringe wetlands along the shoreline of the
1961 lake. Restore damaged habitat by re-grading the shoreline to the natural slope, planting
1962 native vegetation, controlling erosion with better management of stormwater and culvert
1963 replacement. Where natural lakes have been modified with dams, explore options for dam
1964 removal and restoration of more natural hydrology of the lake. Minimize future water
1965 diversion and water use to protect the water flow into and depth of natural lakes.

1966 **Limiting Factor: Invasive species**

1967 Invasive species can compete with native species for food and space, spread diseases,
1968 and produce toxins. Zebra and quagga mussels are highly invasive and can enter natural
1969 lakes from boat hulls, motors, trailers, livewells, and standing water. New Zealand mud
1970 snails can hitchhike on watercraft and fishing gear. Some of the Asian carp species degrade
1971 natural lakes by outcompeting native fish species, increasing water turbidity, and limiting

1972 waterfowl and shorebird production and use. Eurasian watermilfoil spreads through seeds
1973 and vegetative fragments. Invasive turtles and frogs can spread naturally or through
1974 introduction by a previous pet owner.

1975 **Recommended Approach**

1976 Provide outreach and education to the public regarding the impacts of invasive and exotic
1977 species on natural lake ecosystems. This includes outreach about the requirement to stop
1978 at ODFW boat check stations to inspect motorized boats and canoes, kayaks and
1979 paddleboards for invasive species (i.e., aquatic zebra or quagga mussels, snails, and
1980 aquatic plants) to ensure that the invasive species are not being transported between
1981 water bodies. Support programs to prevent carp and other non-native fish (i.e. catfish and
1982 non-native trout), bullfrogs, pet turtles, Eurasian milfoil, purple loosestrife and other
1983 invasive species from being transported and released into natural lakes. Conduct voluntary
1984 monitoring and control efforts.

1985 **Limiting Factor: Wildfires**

1986 Wildfires can have detrimental impacts on natural lakes and can compromise lake water
1987 quality both during active burning and for months and years after the fire is contained.
1988 Accidental human caused fires during the fire season can also create severe impacts to the
1989 lake and surrounding area. Wildfires and accidental human caused fires remove
1990 vegetation that, when intact, helps slow precipitation and hold soil in place, which can lead
1991 to increased stormwater runoff and erosion. Runoff and smoke can carry debris, sediment,
1992 ash, nutrients and other contaminants into the lakes. Wildfires can affect air quality and
1993 the recreation at natural lakes. Forest fires near and at Crater Lake have resulted in such
1994 poor air quality and visibility that the Crater Lake National Park has been closed, through-
1995 hikers on the Pacific Crest Trail have been rerouted, and major events have been cancelled.

1996 **Recommended Approach**

1997 Continue education regarding the impacts of wildfires on natural lakes' air quality, water
1998 quality and recreational opportunities. Continue educating the public to follow recreation
1999 rules for lakes and the surrounding area (i.e., do not build fires outside of an authorized
2000 camping/fire pit or build fires at the lake edge, do not drive vehicle on grass roads or in
2001 natural areas during posted fire seasons). Enforce fire bans and increase awareness when
2002 they go into effect. Where appropriate, thin forests and manage fuels in high priority areas
2003 surrounding lakes.

2004 **Limiting Factor: Water Quantity**

2005 Water is limited in some parts of the state and is projected to become scarcer under a
2006 changing climate and expanded human use. In standing waterbodies, water scarcity can
2007 lead to higher concentrations of contaminants, lowering water quality as less fresh water is
2008 available to dilute nutrients or pollutants. As water quantity diminishes in lakes and
2009 adjacent floodplains, fish are unable to access the shoreline habitat for spawning and
2010 access upstream habitat for life cycle requirements and/or to move to more favorable
2011 conditions. Groundwater pumping and water diversions for out-of-stream uses occur in
2012 every basin, and these uses can impair water quality and quantity, and aquatic species use
2013 and conditions upstream, within and downstream of the lakes. Late summer is a time of
2014 particular concern regarding water quantity.

2015 **Recommended Approach**

2016 Maintain, protect, or restore the natural hydrological cycle for stream flow into lakes.
2017 (KCI: **Water Quality and Quantity**). Develop and adopt instream water rights to increase
2018 quality habitat in and along natural lakes. Develop and implement groundwater
2019 management plans in the lake basins. Manage water in the state with the long-term health
2020 and sustainability of surface and groundwater sources. Continue to educate Oregonians
2021 about conscientious water use.

2022 **RESOURCES FOR MORE INFORMATION**

2023 [Department of Environmental Quality: National Aquatic Resource Surveys: Water](#)
2024 [Quality: State of Oregon](#)

2025 [Oregon Natural Desert Association: Lake Abert](#)

2026 [Oregon Health Authority Website: Cyanobacteria](#)

2027 [Climate Change at Crater Lake](#)

2028 [USGS: Wildfire Impact on Water Quality of California Lakes](#)

2029 [The Center for Lakes and Reservoirs at Portland State University](#)

2030 [Oregon Lake Watch, 2014 Annual Report](#)

2031 [Atlas of Oregon Lakes](#)

2032 [Oregon Lakes Association](#)

2033 For information on boating safety and information about invasive species and other
2034 concerns, see the [Oregon State Marine Board](#)

2035

DRAFT

2036 OAK HABITATS

2037 There are several oak habitat types in Oregon, where oaks comprise most of the canopy.
2038 These can include oak woodlands, oak forest, oak chaparral, and riparian oak. Oak
2039 savannah is covered in the Grassland Key Habitat. Oaks may also co-dominate a canopy in
2040 oak/fir, oak pine, and oak hardwood habitats.

2041 ECOREGIONS

2042 The range of oak habitats are a Key Habitat in the Coast Range, East Cascades, Klamath
2043 Mountains, West Cascades, and Willamette Valley ecoregions.

2044 CHARACTERISTICS

2045 In general, the understory of an Oregon white oak woodland is relatively open with shrubs,
2046 grasses, and wildflowers. The tree canopy of most oak woodland obscures 25-75 percent of
2047 the sky, and an oak forest typically has more than 75 percent cover. Oak chaparral has a
2048 short, shrubby vegetation understory. Riparian oak can tolerate wetter conditions and may
2049 be mixed with other tree species including ash and willow. Oak habitats are ideally
2050 maintained through periodic, low-intensity fire, which removes small conifers and
2051 maintains a moderate cover of low shrubs.

2052 Depending on the ecoregion and site characteristics, oak habitats may also include
2053 ponderosa pine, California black oak, Douglas-fir, madrone, canyon live oak, and tanoak.
2054 Tanoak is closely related to true oaks, sharing a family, but is not a true oak. Tanoak,
2055 however, is an important mast producer often associated with canyon live oak.

2056 ECOREGIONAL CHARACTERISTICS

2057 **Coast Range**

2058 Oak habitats are typically found in drier landscapes, such as south-facing slopes and
2059 foothills bordering the Willamette Valley. The southwestern Oregon coast range is the
2060 northerly extent of the range of canyon live oak and tanoak.

2061 **East Cascades**

2062 In the East Cascades ecoregion, oak woodlands occur primarily on the north end of the
2063 ecoregion and in the south along the Klamath River Canyon. They are located at the
2064 transition between ponderosa pine or mixed conifer forests in the mountains, and the
2065 shrublands or grasslands to the east. Oak habitats in the East Cascades are different in
2066 structure and composition than those in western Oregon but are just as important to a
2067 variety of wildlife and rare plants.

2068 **Klamath Mountains**

2069 Oak habitats are found in lower elevations in the valley floors up to 4000', on dry sites, or in
2070 areas with frequent, low-intensity fires. Oak woodlands may occur in a mosaic with
2071 chaparral and dry conifer woodlands. Nearing the northern extent of its range in this
2072 ecoregion, chaparral is dominated by shrubs species including buckbrush and manzanita
2073 thickets, with deer brush, yerba santa, and silk tassel making up the rest of the shrub
2074 component.

2075 **West Cascades**

2076 Oak woodland habitats are found in drier landscapes, such as south-facing slopes and
2077 foothills bordering the Willamette Valley. Oak habitats extend up to 3500' in southwestern
2078 Oregon in the West Cascades. Portions of the West Cascades may have historically had a
2079 more closed canopy oak habitat as well as very expansive chaparral that filled the
2080 understory.

2081 **Willamette Valley**

2082 In the Willamette Valley, Oregon white oaks were originally found in a mosaic of oak
2083 savanna, forests, and riparian habitats throughout the valley floor and low-elevation
2084 slopes. One variation of oak habitat, that has almost disappeared due to historic harvest, is
2085 white oak and Willamette Valley ponderosa pine. This habitat type is found in valley
2086 bottoms and is tolerant of seasonal flooding. Oaks were most common on flat to
2087 moderately rolling terrain, usually in drier landscapes, and often between prairie remnants
2088 and conifer forests. Today, oak woodlands generally are found in small pockets and some
2089 corridors surrounded by other land uses, such as development or agriculture.

2090 CONSERVATION OVERVIEW

2091 Oak habitats, traditionally managed on a landscape scale by Indigenous people, once
2092 covered almost one million acres in the Coast Range and 400,000 acres in the Willamette
2093 Valley. However, the Coast Range now has very little of its estimated historical oak
2094 woodlands, and the Willamette Valley has even less. Habitat loss has been less severe in
2095 the East Cascades, where fire suppression may have led to expansion of oaks into former
2096 shrub-steppe and grassland habitats. Valuing Traditional Ecological Knowledge and
2097 cultural burning in oak management is critical to protecting and restoring oak habitats.

2098 Oak habitats have been impacted by conversion to other land uses, invasive species, and
2099 vegetation changes due to fire suppression. As a result of conifer plantings and changes in
2100 fire frequency and intensity after European settlement, Douglas-fir now dominates in many
2101 areas of the Coast Range and Willamette Valley foothills.

2102 Oak habitats have been converted to agriculture, residential, and other uses in the
2103 Willamette Valley, the Coast Range foothills, and the coastal hills in southern Oregon.
2104 Development continues to threaten these habitats in all ecoregions where they are found.
2105 The same rolling hills and scenic landscapes that indicate healthy pine-oak habitat also
2106 attract new residents and developers. Because much of the remaining oak habitats are in
2107 private ownership and maintenance of these habitats requires active management,
2108 cooperative incentive-based approaches are crucial to conservation.

2109 Oak habitats provide important food sources, shelter, and resting places for a large range of
2110 birds and other wildlife, including a variety of species that are oak-obligates like Oak
2111 Titmouse and the Acorn Woodpecker. Loss of oaks, particularly large-diameter, open-
2112 structured trees valuable to wildlife, is of particular concern because oak trees have a slow
2113 growth rate, slowing restoration success. In addition, reproduction and recruitment of
2114 younger trees are poor in many areas.

2115 Sudden oak death, a fungal tree pathogen identified in northern California in the 1990s has
2116 been slowly spreading north. In 2001, the pathogen was detected in Curry County, which
2117 continues to be the only area in Oregon where the pathogen is known to occur in a natural
2118 setting. Mediterranean oak borer was found in Oregon in 2018 and is also being tracked.

2119 Depending on the area, Species of Greatest Conservation Need associated with oak
2120 habitats include Columbian white-tailed deer, Chipping Sparrow, Slender-billed White-
2121 breasted Nuthatch, Lewis's Woodpecker, Western Bluebird, Fender's blue butterfly,

2122 Kincaid’s lupine, white rock larkspur, and wayside aster among others. Northern spotted
2123 owl may utilize oak trees in a mixed forest setting.

2124 **[SPOTLIGHT] Acorn Woodpeckers**

2125 Acorn Woodpeckers are striking, social birds found in Oregon’s oak woodlands. Known for
2126 their black-and-white plumage and red cap, they rely on oak trees for food and nesting.
2127 These woodpeckers store acorns in trees, creating “granaries” that serve as both food
2128 storage and breeding sites. They are highly social and often live in groups, working together
2129 to protect their acorn caches and raise their young. The Acorn Woodpecker is a year-round
2130 resident of oak woodland and savanna habitat in western Oregon, primarily found in the
2131 Willamette Valley and Klamath Mountains ecoregions, though occasionally found in the
2132 East Cascades.

2133 Acorn Woodpeckers have been expanding their range in Oregon since the arrival of Euro-
2134 American settlers, expanding north from Roseburg to Eugene between 1920 and 1950, then
2135 further north to Washington County by the early 1990s. Although more than 95% of the oak
2136 woodlands have been lost in the Willamette Valley since European settlement, Acorn
2137 Woodpecker expansion into the Willamette Valley in the last 100 years was likely assisted
2138 by the reduction of fires that maintained grasslands and savanna, transforming some of
2139 those habitats into oak woodlands (mixed with pine or fir) that had larger and older trees
2140 that produce more acorns and provide more dead limbs for granaries and nests.

2141 As one of the most conspicuous residents of Oregon’s oak woodlands, the Acorn
2142 Woodpecker can be an effective catalyst for conservation of this Key Habitat. In Oregon,
2143 much of the remaining oak habitat can be found on private lands. These oak woodlands are
2144 important habitat for many other species.

2145 **LIMITING FACTORS AND RECOMMENDED APPROACHES**

2146 **Limiting Factor: Fire Suppression and Fir Encroachment**

2147 With fire suppression, Douglas-fir encroaches into oak habitats and eventually shades out
2148 oak trees and seedlings, as well as other plants that require open growing conditions. Many
2149 oak woodlands are now dominated by Douglas-fir or have transitioned to fir-oak habitats
2150 due to fire suppression. Without active management, these areas will eventually become
2151 conifer forests. In some areas of the East Cascades, fire suppression combined with
2152 grazing has influenced fine fuel production and led to encroachment by conifers and
2153 establishment of dense patches of small, shrubby oaks.

2154 Large wildfires, like those experienced across the West Cascades and Eastern Oregon have
2155 galvanized public interest in fuels reduction treatments across public and private lands.
2156 When conducted in a manner to retain some understory habitat for wildlife, such as
2157 thinning of small diameter conifers and small diameter oak-on-oak encroachment with
2158 piles and habitat clumps, oak habitats can be restored to fire resiliency and prepped for
2159 low-intensity controlled burns.

2160 **Recommended Approach**

2161 Work with partners to update smoke management and air quality standards to allow more
2162 fall, winter, and spring burn windows for prescribed burning. The Certified Burn Manager
2163 program and cultural waivers have increased equitable access to prescribed fire. Use
2164 multiple tools, including prescribed fire, mowing, grazing, and selective harvest to maintain
2165 open canopy oak-dominated habitats. Ensure that tools are site-appropriate and
2166 implemented to minimize impacts to native species. Re-establish site-appropriate native
2167 grasses, herbaceous plants, and shrubs. (KCI: Disruption of Disturbance Regimes)

2168 **Limiting Factor: Land Use Conversion and Continued Habitat Loss**

2169 Particularly in the Willamette Valley and Klamath Mountains ecoregions, oak habitats
2170 continue to be converted to agricultural (e.g., vineyards), rural residential, urban, and other
2171 land use changes. Remaining oaks can be impacted by soil compaction in agricultural and
2172 residential settings. The conversion of oak habitats into other land uses results in habitat
2173 loss and fragmentation for wildlife, invasive species, and the spread of diseases.

2174 **Recommended Approach**

2175 Much of the remaining oak habitat requires active management and occurs on private land,
2176 where cooperative incentive programs are the best approach. Work with private
2177 landowners to maintain and restore oak habitats and implement outreach and education
2178 efforts. Promote oak conservation on working lands through incentive programs and other
2179 collaborative efforts, such as the Wildlife Conservation and Management Program. Create
2180 new opportunities for acquisition and conservation easements to protect oak habitat, such
2181 as through the Oregon Agricultural Heritage Program. Work with local governments to
2182 protect and conserve oak habitat in local land use planning, through Statewide Planning
2183 Goal 5 as significant wildlife habitat for SGCN.

2184 **Limiting Factor: Loss of Habitat Structure**

2185 Large-diameter oak trees with lateral limb structure and cavities have been lost. In many
2186 areas, there are not sufficient numbers of replacement trees to maintain these habitat
2187 elements over time. In the absence of fire, densely stocked, regenerating oaks often do not
2188 develop open-grown structures due to shading. In the East Cascades, grazing or very hot
2189 fires can lead to development of brushy-structured trees. The shaded or grazed oaks do not
2190 develop the lateral limbs, cavities, and higher acorn crops of open-grown trees, and are
2191 thus less valuable to wildlife. Woodcutting often removes snags, which are necessary for
2192 cavity nesting species.

2193 **Recommended Approach**

2194 Maintain a diversity of tree sizes and ages across the stand, with emphasis on large oak and
2195 other key tree species appropriate to the habitat type. Remove conifers that are competing
2196 with larger oaks. Maintain existing snags and create new snags from competing conifers to
2197 provide cavity habitat. Encourage oak reproduction through plantings or protective
2198 enclosures. It may be appropriate to use nest boxes as temporary cavity habitat in oak
2199 restoration project areas. Improve methods to promote oak reproduction and creation of
2200 open-grown structures.

2201 **Limiting Factor: Invasive Species and Diseases**

2202 In many remaining oak habitats, the overstory is intact but the understory is highly
2203 degraded. Depending on the ecoregion and site, invasive plants, such as Armenian
2204 (Himalayan) blackberry, bird cherry, evergreen blackberry, Scotch broom, English
2205 hawthorn, false brome, yellowstar thistle, diffuse knapweed, and puncturevine, have
2206 established and degraded oak habitats.

2207 Invasive insects, such as the Mediterranean oak borer and carpenter worm moth spread
2208 diseases, cause defoliation, and weaken the structure of the trees. Fungal diseases such
2209 Sudden Oak Death and Armillaria root rot can also significantly impact oak trees.

2210 **Recommended Approach**

2211 Emphasize prevention, risk assessment, early detection, and quick control to prevent new
2212 invasive species from becoming fully established. Prioritize control efforts and use site-
2213 appropriate methods to control newly established invasive plant species for which
2214 management can be most effective. In high-risk areas, use weed-wash stations for

2215 machinery during mechanical restoration or treatment of a site. Re-seed with site-
2216 appropriate native grasses and forbs after control efforts. Prescribed burning may be useful
2217 for management of some invasive species, particularly shrubs. Support efforts toward
2218 expanding native seed resources. (KCI: Invasive Species)

2219 **Limiting Factor: Climate Change**

2220 The mean annual air temperature in the Pacific Northwest is projected to increase under a
2221 changing climate. This warming is projected to be the highest during the summer. Annual
2222 precipitation patterns in the Pacific Northwest are also predicted to change, with
2223 decreases in summer precipitation. While oaks may be tolerant of warmer and drier
2224 summer conditions, the severity of the impact may have detrimental effects.

2225 **Recommended Approach**

2226 Protect and restore a diverse portfolio of oak habitats to preserve genetic diversity.
2227 Continue efforts to restore currently degraded areas and re-establish former oak habitats
2228 to increase resiliency. Engage in strategic, landscape-scale planning efforts to create a
2229 connected network of oak habitats to increase adaptive capacity. Identify where future
2230 climate conditions may support oak habitats, including areas upslope of their current
2231 range where they were not historically found. Identify data gaps and support research
2232 needs, including the protection of oak habitat on natural and working lands to mitigate the
2233 impacts from climate change.

2234 **HABITAT TRENDS ANALYSIS**

2235 Open Oregon white oak and black oak woodlands were common across western Oregon
2236 prior to Euro-American settlement. Subsequent fire suppression and development
2237 pressures of the 20th century led to a loss of oak woodlands. To investigate the magnitude
2238 of loss of oak habitats, the Institute of Natural Resources (INR) compared the total area of
2239 oak habitats in three baseline maps (1851-1937, 1998, 2016). The analysis showed
2240 significant of oak habitats over time. By 2016, the total area of oak habitats had decreased
2241 by 72% when compared to historical data. Oak habitats were largely replaced by mixed
2242 hardwood and conifer forests and agricultural land uses.

2243 **RESOURCES FOR MORE INFORMATION**

2244 [Pacific Northwest Oak Alliance](#)

- 2245 Partners in Flight Conservation of Landbirds and Associated Habitats and Ecosystems in
2246 the East Cascade Mountains of Oregon and Washington
- 2247 Land Manager's Guide to Bird Habitat and Populations in Oak Ecosystems of the Pacific
2248 Northwest
- 2249 Oregon White Oak Restoration Strategy for National Forest System Lands East of the
2250 Cascade Range
- 2251 Cascadia Prairie Oak Partnership
- 2252 Restoring Oak Habitats in Southern Oregon & Northern California v2.0
- 2253 Restoring Oak Habitats in Southern Oregon & Northern California: A Guide for Private
2254 Landowners v3.0
- 2255 Wildlife-friendly Fuels Reduction in Dry Forests of the Pacific Northwest

2256 **REFERENCES**

- 2257 Altman, B. and J. L. Stephens. 2012. Land Managers Guide to Bird Habitat and Populations
2258 in Oak Ecosystems of the Pacific Northwest. American Bird Conservancy and Klamath Bird
2259 Observatory. 82 pp.
- 2260 Brunner, R. and E. Gaines. Institute for Natural Resources. 2025. Oregon Vegetation
2261 Change 1851-2023. Trends analysis conducted for Oregon Department of Fish and Wildlife.
- 2262 Hosten P.E., O.E. Hickman, F. Lang. 2007. Patterns of vegetation change in grasslands,
2263 shrublands, and woodlands of southwest Oregon. USDI, Bureau of Land Management,
2264 Medford District, OR. Website
2265 <https://www.blm.gov/or/districts/medford/files/pattvegchange.pdf>

2266

2267 PONDEROSA PINE WOODLANDS

2268 Ponderosa pine woodlands are common in Oregon's eastside ecoregions. While
2269 dominated by ponderosa pine, these woodlands may also have lodgepole pine, western
2270 juniper, aspen, western larch, grand fir, Douglas-fir, mountain mahogany, incense cedar,
2271 sugar pine, or white fir, depending on ecoregion and site conditions. Known for their open
2272 forest structure, these woodlands generally have fewer than 40 large trees per acre, with
2273 tree canopy cover between 10 and 60%. Understories consist of variable combinations of
2274 fire tolerant shrubs, herbaceous plants, and grasses. Ponderosa pine forests generally
2275 occur in regions with arid conditions with little rainfall during summer months, and can be
2276 found at a range of elevations, from 100 ft to over 6000 ft.

2277 ECOREGIONS

2278 Ponderosa pine woodlands are a Key Habitat in the **Blue Mountains, East Cascades,** and
2279 **Klamath Mountains.**

2280 CHARACTERISTICS

2281 The open structure of ponderosa pine habitats was historically maintained by frequent,
2282 low-intensity surface fires, with some intermittent higher-intensity fires. The thick bark of
2283 mature ponderosa pines provides protection against moderate fires, allowing these trees
2284 to survive and regenerate after fire events. The structure and composition of ponderosa
2285 pine woodlands vary across the state, depending on local climate, soil type and moisture,
2286 elevation, aspect, and fire history. The soils in ponderosa pine woodlands are often well-
2287 drained and sandy or loamy. Ponderosa pine woodlands typically have an open canopy
2288 structure, allowing sunlight to penetrate and support a diverse understory of herbaceous
2289 plants and shrubs.

2290 ECOREGIONAL CHARACTERISTICS

2291 **Blue Mountains**

2292 In the Blue Mountains, ponderosa pine often coexists with other conifers, such as Douglas-
2293 fir, western larch, and grand fir. The understory is diverse, including shrubs like mountain
2294 big sagebrush, bitterbrush, mahogany, snowbrush and various native grasses and forbs
2295 such as Idaho fescue and bluebunch wheatgrass. Ponderosa pine habitats also include
2296 savannas, which have sporadic, widely spaced trees that are generally more than 150
2297 years old. The structure of a savanna is open with an understory dominated by fire-adapted
2298 grasses and forbs as well as shrub fields. Ponderosa pine habitats in the Blue Mountains

2299 generally occur at mid elevation and are replaced by other coniferous forests at higher
2300 elevations.

2301 **East Cascades**

2302 East of the foothills of the Cascades, within the rain shadow cast by the mountains, land
2303 becomes more arid and ponderosa pine woodlands become dominant. In these
2304 woodlands, other conifer species may be present, including Douglas-fir, western larch,
2305 and, in some areas, lodgepole pine. The understory is characterized by a mix of shrubs and
2306 herbaceous plants. Common shrubs include bitterbrush, mountain big sagebrush, and
2307 snowberry. The herbaceous layer often includes native grasses such as Idaho fescue and
2308 bluebunch wheatgrass. Ponderosa pine habitats in the East Cascades generally occur at
2309 mid elevation, where climatic and soil conditions support the growth of these trees, and
2310 are replaced by other coniferous forests at higher elevations.

2311 **Klamath Mountains**

2312 In the Klamath Mountains, pine woodlands are usually dominated by ponderosa pine, but
2313 may be dominated by Jeffery pine, depending on soil mineral content, fertility, and
2314 temperatures. Ponderosa pine and ponderosa pine-oak woodlands occur on dry, warm
2315 sites in the valley margins, foothills, and mountains of southern Oregon. The understory
2316 often has shrubs, including green-leaf manzanita, buckbrush, and snowberry.

2317 **CONSERVATION OVERVIEW**

2318 Ponderosa pine habitats historically covered a large portion of the Blue Mountains
2319 ecoregion, as well as parts of the East Cascades and Klamath Mountains. Ponderosa pine
2320 is still widely distributed in eastern and southern Oregon. However, the structure and
2321 species composition of ponderosa pine woodlands have changed dramatically. In the
2322 past, ponderosa pine habitats had frequent low-intensity fires that maintained an open
2323 understory, as well as some high-intensity fires. Due to certain timber harvest practices,
2324 the exclusion of Indigenous peoples' burning practices, and fire suppression, dense
2325 patches of smaller conifers have overtaken the understory of many ponderosa pine
2326 forests. Depending on the region, these conifers may include shade-tolerant Douglas-fir,
2327 grand fir, white fir, and lodgepole pine as well as young ponderosa pines. These dense
2328 stands are highly vulnerable to drought stress, insect outbreaks, and disease. Many of
2329 these mixed conifer forests are in fire-prone areas where the risk of loss of key ecosystem
2330 components is high. Due to this unnatural density of understory trees in these areas, low
2331 intensity fires can quickly become severe and kill large, mature ponderosa pine trees that
2332 would survive a smaller fire. Fire suppression has also led to the accumulation of a thick
2333 layer of needle duff in the understory. Large trees will send roots into this duff layer, which
2334 are then destroyed when the duff layer burns, often killing mature trees even in lower
2335 intensity fires.

2336 Loss and conversion of ponderosa pine woodlands to shrubland and other habitat types is
2337 occurring largely as a result of the increased scale and frequency of high intensity wildfires.
2338 While ponderosa pines readily re-establish after disturbance, high intensity, landscape
2339 scale wildfires make it difficult for successful regeneration post-fire due to a lack of natural
2340 seed source within seed dispersal ranges. Of particular concern is the loss of large-
2341 diameter ponderosa pine habitats. Most old-growth ponderosa pine stands are greatly
2342 reduced in size and connectivity, occurring in a patchwork with much younger forests.
2343 Younger stands can provide habitat for some wildlife species, but old-growth ponderosa
2344 pine forests provide critical habitat for wildlife that prefer open, dry forests.

2345 Ponderosa pine woodlands support a diversity of wildlife species, including Species of
2346 Greatest Conservation Need (SGCN). One SGCN, the white-headed woodpecker, requires
2347 large-diameter trees and an open understory and is sensitive to changes in the forest seral
2348 stage. As a result, white-headed woodpeckers are entirely dependent on open, late-
2349 successional ponderosa pine woodlands. Other SGCN associated with ponderosa pine
2350 habitats include **Flammulated Owl, Lewis's Woodpecker, long-legged myotis,**
2351 **and pallid bat.**

2352 On federal and private lands, especially in the wildland-urban interface, ponderosa pine
2353 habitats are increasingly being restored or managed in a manner consistent with wildlife
2354 conservation goals through fuel reduction treatments, retention of large-diameter trees,
2355 and maintenance of high densities of snags.

2356 LIMITING FACTORS AND RECOMMENDED APPROACHES

2357 **Limiting Factor: Altered Fire Regimes and Addressing Risk of Uncharacteristically Severe** 2358 **Wildfire**

2359 Certain timber harvest practices, the exclusion of Indigenous peoples' burning practices,
2360 and fire suppression have resulted in dense growth of young pine trees and young mixed
2361 conifer stands, replacing the open understory of ponderosa pine woodlands. These dense
2362 stands are at increased risk of uncharacteristically severe wildfires, drought, disease, and
2363 damage by insects. Over time, some stands will convert to Douglas-fir and grand fir
2364 forests, which do not provide adequate wildlife habitat for species dependent on open
2365 ponderosa pine habitats. While normally drought tolerant, large old-growth ponderosa
2366 pines are dying due to these dense young trees that would historically have been
2367 controlled by frequent, low intensity fires.

2368 These dense understories, along with numerous insect-killed trees, make it difficult to
2369 reintroduce natural fire regimes in some areas, particularly in the Blue Mountains and East
2370 Cascades ecoregions. In parts of the Blue Mountains, East Cascades, and Klamath
2371 Mountains, increasing development of homes and resorts in forested habitats limits the
2372 ability of managers to use prescribed fires due to concerns about smoke and escaped
2373 burns, further increasing the risk of high-intensity wildfires. Some ponderosa pine

2374 woodlands are also being inundated with invasive annual grasses such as cheatgrass and
2375 medusahead, increasing fuel continuity and altering natural fire behavior.

2376 **Recommended Approach**

2377 Use an integrated approach to forest health issues that considers historical conditions,
2378 including roads and human use, wildlife conservation, natural fire intervals, and
2379 silvicultural techniques. Develop implementation plans for thinning overstocked stands
2380 and applying prescribed fire, and ensure plans are acceptable for management of both
2381 game and non-game species. Evaluate individual stands to determine site-appropriate
2382 actions, such as monitoring in healthy stands, or thinning, mowing, and application of
2383 prescribed fire in at-risk stands. Develop markets for small-diameter trees and implement
2384 fuel reduction projects to reduce the risk of forest-destroying wildfires. Manage for a
2385 landscape mosaic that includes structural complexity and species diversity in the
2386 understory and overstory across multiple spatial scales. Fuel reduction strategies need to
2387 consider the habitat structures that are required by wildlife, including snags, downed logs,
2388 and hiding cover. Reintroduce fire where feasible. Engage with Tribal Nations to bring
2389 Traditional Ecological Knowledge of prescribed fire to the overstocked forests. Implement
2390 prescribed fire at a frequency and scale that improves regeneration and establishment of
2391 native shrubs

2392 Support community-based forest collaboratives to increase the pace and scale of forest
2393 restoration. Engage in frequent outreach to educate the public about the ecological
2394 importance of fire to ponderosa pine forests. Monitor forest health initiatives and use
2395 adaptive management techniques to ensure efforts are meeting habitat restoration and
2396 uncharacteristic fire prevention objectives with minimal impacts on wildlife. Work with
2397 landowners and resort operators to reduce vulnerability of properties to wildfires while
2398 maintaining habitat quality. Highlight successful, environmentally sensitive fuel
2399 management programs. Retain features that are important to wildlife, including snags,
2400 downed logs, forage, and hiding cover for wildlife species, and replant with native shrub,
2401 grass, and forb species. Manage reforestation after wildfire to create species and
2402 structural diversity based on local management goals. (KCI: **Disruption of Disturbance**
2403 **Regimes**)

2404 **Limiting Factor: Loss of Size and Connectivity of Large-structure Ponderosa Pine Habitats**

2405 Old-growth ponderosa pine habitats have been greatly reduced in size and connectivity by
2406 timber harvest, the exclusion of Indigenous peoples' burning practices, and fire
2407 suppression, particularly in the Blue Mountains and East Cascades ecoregions. These
2408 changes have led to overstocked stands. Alongside the loss of open understories and
2409 encroachment by dense stands of young trees, many ponderosa pine habitats have been
2410 lost to conversion to rural residential uses and other activities. As a result, few large,
2411 contiguous blocks of ponderosa pine habitat remain.

2412 **Recommended Approach**

2413 Maintain large blocks of large-diameter ponderosa pine habitat. Identify current and
2414 potential movement **corridors** between habitat blocks for protection and restoration. In
2415 areas experiencing rapid development, work with local communities to minimize
2416 development in large blocks of intact habitat.

2417 **Limiting Factor: Invasive Species**

2418 Throughout the state, non-native plants are invading and degrading ponderosa pine
2419 woodlands. In parts of the Blue Mountains and East Cascades, diffuse and spotted
2420 knapweed and Dalmatian and common toadflax are significant invaders. Additionally, in
2421 many areas the spread of cheatgrass and medusahead rye can result in an invasive plant
2422 understory that is highly susceptible to burning, with a high-fuel content vegetation that
2423 carries wildfire more easily than the native vegetation. In the Klamath Mountains,
2424 Armenian (Himalayan) blackberry and Scotch broom are significant invaders, along with
2425 annual invasive grasses.

2426 **Recommended Approach**

2427 Emphasize prevention, risk assessment, early detection, and quick control to prevent
2428 new **invasive species** from becoming fully established. Prioritize efforts and control key
2429 invasive species using site-appropriate methods. Control wildfires in cheatgrass-
2430 dominated areas of the Blue Mountains. Fortunately, many areas within the Blue
2431 Mountains and East Cascades still have few invasive species currently threatening
2432 ponderosa pine habitats. In these areas, invasive plants should be monitored and
2433 controlled as they first arrive when control is more efficient, practical, and cost-effective.
2434 Reintroduce site-appropriate native grasses and forbs after invasive plant control.
2435 Prescribed burning may be useful for management of some invasive species in the
2436 Klamath Mountains.

2437 **RESOURCES FOR MORE INFORMATION**

2438 Oregon Department of Forestry Forest Practices Monitoring Program

2439 Partners in Flight Conservation of Landbirds and Associated Habitats and Ecosystems
2440 in the Northern Rocky Mountains of Oregon and Washington

2441 Partners in Flight Conservation of Landbirds and Associated Habitats and Ecosystems
2442 in the East Cascade Mountains of Oregon and Washington

2443 Managing for Cavity-Nesting Birds in Ponderosa Pine Forests

2444

2445 SAGEBRUSH HABITATS

2446 Sagebrush habitats include all sagebrush steppe- and shrubland-dominated communities
2447 found east of the Cascade Mountains.

2448 ECOREGIONS

2449 Sagebrush habitats are a Key Habitat in the **Blue Mountains, Columbia Plateau, East**
2450 **Cascades**, and **Northern Basin and Range** ecoregions.

2451 CHARACTERISTICS

2452 Sagebrush habitats in eastern Oregon are both extensive and diverse, ranging from low-
2453 elevation valleys to high mountain areas and from grassland-like shrub-steppe to relatively
2454 dense shrublands. Sagebrush-dominated communities differ in structure and species
2455 composition depending on ecoregion, elevation, soils, moisture regimes, and fire history.

2456 Sagebrush habitats are often classified as sagebrush steppe or sagebrush shrublands.
2457 Sagebrush steppe is characterized by grasses and forbs with an open or more dispersed
2458 shrub layer. Sagebrush shrublands are dominated by shrubs, with less understory area
2459 covered by grasses and forbs than in steppe habitats.

2460 In Oregon, sagebrush habitats are dominated by mountain big or Wyoming big sagebrush.
2461 Both mountain big and Wyoming big sagebrush habitats historically experienced natural
2462 fire regimes that maintained a patchy distribution of shrubs and predominance of native
2463 grasses. Big sagebrush, including mountain, Wyoming, and basin, thrives in deep, well-
2464 drained soils. Low sagebrush, however, prefers shallow, wet soils.

2465 ECOREGIONAL CHARACTERISTICS

2466 **Blue Mountains**

2467 The Blue Mountains ecoregion has both mountain big and Wyoming big sagebrush
2468 habitats. These habitats have different plant associations depending on elevation and soils
2469 including several different subspecies of sagebrush such as low sagebrush, silver
2470 sagebrush, rigid sagebrush, basin big sagebrush, and threetip sagebrush. Other common
2471 plant species include **bitterbrush**, mountain mahogany, and rabbitbrush. Soils vary in
2472 depth and texture but are non-saline.

2473 **Columbia Plateau**

2474 Columbia Plateau shrub-steppe habitats are open grass-dominated communities and are
2475 usually found on loamy, wind-deposited (loess) soils. In this ecoregion, shrub-steppe
2476 communities can be broadly divided into two elevational types. Within 10 miles of
2477 the Columbia River, sandy shrub-steppe communities occur on unstable, well-drained
2478 soils with a component of bare ground or open sand present. These communities range
2479 from sagebrush steppe dominated by bitterbrush and needle-and-thread grass or Indian
2480 rice grass, to sand dune communities characterized by sagebrush, bitterbrush, and
2481 western juniper. Further from the Columbia River, both mountain big and Wyoming big
2482 sagebrush communities include basin big sagebrush, needle-and-thread grass, basin
2483 wildrye and bluebunch wheatgrass steppe, and Wyoming big sagebrush and bluebunch
2484 wheatgrass (which formerly occupied the low-elevation, loess uplands in the Columbia
2485 Plateau).

2486 **East Cascades**

2487 Sagebrush habitats occur in the East Cascades transition zone between the Cascade
2488 Mountain forests and the drier sagebrush steppe habitats of the Columbia Plateau and
2489 Northern Basin and Range Ecoregions. The number of species and acreage dominated by
2490 sagebrush is lower in the East Cascades ecoregion than most other east side ecoregions,
2491 especially the Northern Basin and Range. Mountain big sagebrush and Wyoming big
2492 sagebrush habitats are both found in the East Cascades Ecoregion depending on elevation
2493 and soil type with Wyoming big sagebrush habitats found mainly along the eastern edge.

2494 **Northern Basin and Range**

2495 Big sagebrush habitats include mountain, basin, and Wyoming big sagebrush shrublands
2496 and shrub-steppe. Structurally, these habitats are composed of medium-tall to tall (1.5-6
2497 feet) shrubs that are widely spaced with an understory of perennial bunchgrasses. Basin
2498 big sagebrush communities occur on deep silty or sandy soils along stream channels, in
2499 valley bottoms and flats, or on deeper soil inclusions in low sagebrush or Wyoming big
2500 sagebrush stands. Wyoming big sagebrush communities occur on shallower, drier soils.
2501 Mountain big sagebrush communities occur at montane and subalpine elevations on
2502 deep-soiled to stony flats, ridges, nearly flat ridge tops, and mountain slopes. The fire
2503 frequency in big sagebrush habitats ranges from 10-25 years for mountain big sagebrush
2504 and 50-100 years for Wyoming big sagebrush.

2505 Although big sagebrush communities tend to be the dominant habitat type, other
2506 sagebrush types also provide important habitat for wildlife and may need to be considered
2507 at the local and watershed scale, or for the conservation of particular species like

2508 the **Greater Sage-Grouse**. For example, low sagebrush provides critical wildlife habitat for
2509 many sagebrush-obligate species. Low sagebrush habitats cover large areas of the
2510 Northern Basin and Range ecoregion, but low sagebrush communities are slow (150-300
2511 years) to recover from significant soil disturbance or fire. Soil disturbance in these sites
2512 often results in the establishment of invasive annual grasses.

2513 CONSERVATION OVERVIEW

2514 There are many species and subspecies of sagebrush, which are associated with different
2515 grasses and herbaceous plants, depending on site conditions. General ecology and
2516 conservation issues vary by sagebrush community type, so conservation actions must be
2517 tailored to local conditions and conservation goals. Detailed descriptions of the different
2518 sagebrush plant communities are available from sources included in the references.

2519 Although sagebrush habitats are still common and widespread in eastern Oregon, some
2520 sagebrush habitat types have high levels of habitat loss and are of conservation concern.
2521 These types vary by ecoregion. In the **Blue Mountains**, valley-bottom sagebrush types,
2522 including threetip or basin big sagebrush, that occur on deep soils are particularly at risk.
2523 Also important are the valley margin steppe types with Wyoming big sagebrush, squaw
2524 apple, and **bitterbrush**. Overall, the sagebrush habitats in the Blue Mountains ecoregion
2525 have experienced steep declines since colonization.

2526 In the lower elevations of the **Columbia Plateau**, shrub-steppe communities have been
2527 almost entirely replaced by irrigated agriculture. Remnant habitats occur on public lands,
2528 such as the Boardman Bombing Range, and in scattered patches along roadsides and
2529 fields. Loss of sagebrush habitats in the Columbia Plateau is also high compared to
2530 historical acreages.

2531 In the **Northern Basin and Range** ecoregion, several types of big sagebrush are combined
2532 into a single Key Habitat, including mountain, basin, and Wyoming big sagebrush
2533 shrublands and shrub-steppe. This part of Oregon has some of the largest blocks of high-
2534 quality sagebrush habitat left in the United States, but it is estimated that more than half of
2535 this habitat has been lost since the 1800s. Basin big sagebrush communities have had the
2536 greatest loss as compared to historical distribution. These communities historically
2537 occurred on deep soils and have been converted to agriculture, residential housing and
2538 industrial uses in some areas. The deep soils of basin big sagebrush are important for
2539 pygmy rabbits to create burrows.

2540 Although Wyoming big sagebrush habitats are still common and widespread in the
2541 Northern Basin and Range, they have been altered to some degree by unmanaged grazing,
2542 invasive species, and altered fire regimes. With overgrazing and fire suppression, shrub

2543 (mostly sagebrush) density increases, bunchgrass and forb density decreases, and
2544 invasive annual grasses increase. In many areas, these habitats have shifted from mosaics
2545 of native perennial grasses, forbs, and shrubs to landscapes heavily dominated by shrubs
2546 and invasive annual forbs and grasses. Juniper encroachment is an important issue in
2547 mountain big sagebrush communities between 4,500 and 7,000 feet.

2548 Big sagebrush habitats have high structural diversity, thus more places to forage, hide, and
2549 build nests. As a result, the number of bird species generally increases with sagebrush
2550 height. Habitat values are also dependent on a diverse understory of bunchgrasses and
2551 flowering plants.

2552 Throughout eastern Oregon, loss of grassland-shrub mosaics across landscapes and the
2553 degradation of understories have contributed to the decline of species dependent on high-
2554 quality sagebrush habitats. **Species of Greatest Conservation Need (SGCN)** associated
2555 with sagebrush include the **Greater Sage-Grouse, Ferruginous Hawk, Loggerhead**
2556 **Shrike, Sagebrush Sparrow, Brewer's Sparrow, northern sagebrush lizard, Washington**
2557 **ground squirrel, burrowing owls, and pygmy rabbit.**

2558 LIMITING FACTORS AND RECOMMENDED APPROACHES

2559 **Limiting Factor: Altered Fire Regimes**

2560 **Fire suppression** has resulted in undesirable changes in vegetation and contributes to
2561 increases in the intensity of wildfires. In some fire-suppressed areas, western junipers have
2562 encroached into sagebrush habitats. Dense juniper stands are not suitable for species that
2563 require open sagebrush habitats. Replacement of native bunchgrasses by cheatgrass and
2564 other invasive annual grasses has increased fire frequency and intensity in sagebrush
2565 habitats. Prescribed fire, which can be a useful tool when tailored to local conditions, is not
2566 necessarily suitable for all sagebrush habitat types. Some sagebrush habitats, including
2567 low and Wyoming big sagebrush, are extremely slow to recover from disturbance such as
2568 prescribed fire. Fire, both prescribed and natural, can increase dominance by invasive
2569 plants.

2570 **Recommended Approach**

2571 Carefully evaluate sites to determine if prescribed fire is appropriate, taking into
2572 consideration the extent of invasive annual grasses and other fire prone invasive species in
2573 the area and the recovery potential of the sagebrush community. If determined to be
2574 ecologically beneficial, reintroduce natural fire regimes using site-appropriate
2575 prescriptions. Use prescribed fire to create a mosaic of successional stages and avoid
2576 large burn patches. To control encroaching junipers, use treatment methods such as

2577 mastication, cut and pile, lop and scatter, or cutting for firewood. To ensure the long-term
2578 success of juniper removal, it may be necessary to re-treat stands on a regular basis.
2579 Develop markets for small juniper trees as a special forest product to reduce restoration
2580 costs. Maintain juniper trees with old-age characteristics, which are important for nesting
2581 birds, mule deer winter range, and other wildlife.

2582 **Limiting Factor: Invasive Species**

2583 Invasive plants, such as cheatgrass, medusahead, yellow-star thistle, knapweeds (diffuse,
2584 spotted, and purple), rush skeleton weed, spikeweed, leafy spurge, and perennial
2585 pepperweed, invade and degrade sagebrush habitats. The introduction and spread of
2586 annual grasses, such as cheatgrass and medusahead, has increased the frequency,
2587 intensity, and extent of fires in these habitats. Sagebrush and native bunchgrasses are
2588 adapted to infrequent, patchy fires, and are eliminated by hot fires. Invasive grasses also
2589 provide little nutritious value for wildlife and decrease available forage on the landscape.
2590 While not nearly as extensive as invasive plants, non-native animals have also impacted
2591 native fish and wildlife populations. Unregulated horse and burro herds are a concern in
2592 many areas, competing with native wildlife for vegetation and access to limited water
2593 sources, spreading invasive plant seeds via their manure, and trampling sensitive habitats.

2594 **Recommended Approach**

2595 Emphasize prevention, risk assessment, early detection, and quick control to prevent new
2596 invasive species from becoming fully established. Prioritize control efforts and use site-
2597 appropriate methods to control newly established species for which management can be
2598 most effective (e.g., leafy spurge and perennial pepperweed). Cooperate with partners
2599 through habitat programs and County Weed Boards to address invasive species problems.
2600 Oregon's SageCon Invasives Initiative can be used for state-wide planning and coordinating
2601 implementation and funding toward shared priority areas. Reintroduce shrubs, grasses,
2602 and forbs at control sites through seeding and/or planting. In some cases, it may be
2603 desirable to use "assisted succession" strategies, using low seed rates of non-invasive,
2604 non-native plants in conjunction with native plant seeds as an intermediate step in
2605 rehabilitating disturbances to sagebrush habitat. Prevent and control wildfires in areas
2606 where cheatgrass dominates in the understory. Conduct research to determine methods to
2607 manage established species such as cheatgrass and medusahead. Minimize soil
2608 disturbance in high priority areas to prevent the establishment of invasive species. Work
2609 with public land managers to develop effective and enforceable travel management rules
2610 to prevent the spread of noxious weeds. Promote dialogue between wildlife managers,
2611 landowners, and land managers to develop horse management plans based on common
2612 priorities. Provide outreach to explain the issue to the public and the impacts of
2613 unregulated herds on wildlife and habitat.

2614 **Limiting Factor: Damage to Microbiotic Soil Crusts**

2615 The soil surface of many sagebrush habitats is made up of a community of lichens,
2616 bryophytes, algae, bacteria, and fungi that make up the microbiotic soil crust. These soil
2617 crusts contribute to biodiversity and nutrient cycling and improve soil stability and
2618 structure but are sensitive to disturbance. Unmanaged grazing, agricultural practices,
2619 development, and unregulated OHV use can damage soil crusts, which leads to soil
2620 erosion, changes in plant species composition and structure, and degradation by invasive
2621 plants.

2622 **Recommended Approach**

2623 Because most of the Columbia Plateau ecoregion is privately-owned, **voluntary**
2624 **cooperative approaches** are the key to long-term conservation in this ecoregion. Use tools
2625 such as financial incentives, technical assistance, regulatory assurance agreements, and
2626 conservation easements to achieve conservation goals. Work with public land managers to
2627 ensure grazing is carefully managed and that soil crusts are considered in management
2628 plans. Create effective travel management laws for off-highway vehicle use that can be
2629 successful and enforced.

2630 **Limiting Factor: Conversion to Other Land Uses**

2631 Remnant shrub-steppe habitats are subject to **land use conversion**, such as to
2632 agriculture, urban and rural development, and energy projects. For example, in the
2633 Columbia Plateau and Northern Basin and Range ecoregions, thousands of acres are being
2634 converted to largescale solar energy projects. Large solar array installations can impact
2635 wildlife habitats and block migratory corridors with the development footprint or through
2636 exclusion by project fencing. Mining exploration and development also contribute to
2637 sagebrush habitat loss with both a direct loss as a result of the mine development and
2638 extraction processes, and indirect impacts such as runoff as a result of the mining
2639 operations. Recreation can have negative impacts from off-highway vehicles or dispersed
2640 camping in sensitive habitat or during wet seasons. In the Blue Mountains and East
2641 Cascades ecoregions, rapidly growing human populations, especially near Bend,
2642 Redmond, and Madras, are resulting in land use conversion, habitat loss, and habitat
2643 fragmentation.

2644 **Recommended Approach**

2645 Use tools such as financial incentives and conservation easements to conserve priority
2646 sagebrush habitats on private lands. For example, re-establishing the shrub component of
2647 lands enrolled in the Conservation Reserve Program has helped to restore habitat

2648 structure. Work with community leaders and agency partners to ensure that development
2649 is planned and consistent with local conservation priorities. Support and implement
2650 existing **land use regulations** to preserve farm and range land, open spaces, recreation
2651 areas, and natural habitats from incompatible development.

2652 **Limiting Factor: Loss of Habitat Connectivity**

2653 In the Columbia Plateau, remnant shrub-steppe habitats often occur in small patches,
2654 such as roadsides and field edges. These patches are valuable habitat for some species,
2655 especially some SGCN plants. However, small size and poor connectivity of remnant
2656 patches limit dispersal for sagebrush-associated species.

2657 **Recommended Approach**

2658 Maintain high priority patches and improve connectivity. (KCI: **Barriers to Animal**
2659 **Movement**)

2660 **HABITAT CHANGE TRENDS ANALYSIS**

2661 **Transition to Juniper Woodlands**

2662 Western juniper is distributed across most of eastern Oregon and historically formed open
2663 woodlands with sparse understories. After settlement, however, western juniper began
2664 establishing and spreading into new areas, often forming dense stands with substantial
2665 understories, while open, old growth woodlands declined. With this expansion, juniper has
2666 invaded many of the grasslands and shrublands in eastern Oregon, altering the structure
2667 and function of many sagebrush habitats.

2668 To investigate juniper encroachment into grassland and sagebrush habitats, the Institute
2669 of Natural Resources (INR) compared the total area and spatial overlap of vegetation
2670 classes in three baseline maps (1851-1937, 1998, 2016). The analysis showed significant
2671 increases in the total area of Juniper Woodlands and corresponding losses of Big
2672 Sagebrush habitats to juniper encroachment.

2673 By 2016, the total area of Juniper Woodlands had increased by 115% when compared to
2674 historical data, largely replacing Sagebrush and Grassland habitats. An estimated 38% of
2675 Big Sagebrush habitats were lost to juniper encroachment between 1851 and 1998.
2676 Between 1998 and 2016, an estimated 25% of Big Sagebrush habitats were replaced by
2677 Juniper Woodlands. This analysis shows that significant efforts by land managers and
2678 agencies, such as the BLM and Forest Service, to remove juniper have slowed
2679 encroachment in sagebrush habitats in recent years.

2680 RESOURCES FOR MORE INFORMATION

2681 [Partners in Flight Conservation Strategy for Landbirds in the Columbia Plateau of](#)
2682 [Eastern Washington and Oregon](#)

2683 [Characteristics of Western Juniper Encroachment into Sagebrush Communities in](#)
2684 [Central Oregon](#)

2685 For information from the Bureau of Land Management about rangeland issues, fire
2686 management, and fire and invasive species assessment tools,
2687 see: [https://www.blm.gov/learn/blm-library/subject-guides/greater-sage-grouse-](https://www.blm.gov/learn/blm-library/subject-guides/greater-sage-grouse-subject-guide/documents-and-resources)
2688 [subject-guide/documents-and-resources.](https://www.blm.gov/learn/blm-library/subject-guides/greater-sage-grouse-subject-guide/documents-and-resources)

2689 Convened by the Governor’s Office, the [Sage-Grouse Conservation](#)
2690 [Partnership \(SageCon\)](#) is a diverse group of stakeholders working together since 2012 to
2691 develop an “all lands, all threats” plan to address sage-grouse conservation needs and
2692 support community sustainability in Oregon.

2693 [Sagebrush and Sage Grouse | U.S. Geological Survey](#)

2694 [The SageCon Invasives Initiative | Oregon State University](#)

2695 [Threat Based Ecostate Map | Oregon State University](#)

2696 [Cheatgrass Toolkit](#)

2697 [Defend the Core: Maintaining intact rangelands by reducing vulnerability to invasive](#)
2698 [annual grasses | Working Lands For Wildlife](#)

2699 [Partnering to Conserve Sagebrush Rangelands - IWJV](#)

2700 [Sagebrush Conservation Initiative – WAFWA](#)

2701 [Sagebrush Resources | Grassland & Sagebrush Conservation Portal](#)

2702 [Oregon Department of Agriculture Noxious Weeds webpage](#)

2703 <https://sagebrushconservation.org/>

2704

2705 **REFERENCES**

2706 Brunner, R. and E. Gaines. 2025. Oregon Vegetation Change 1851-2023. Trends analysis
2707 conducted for Oregon Department of Fish and Wildlife. Institute for Natural Resources,
2708 Portland State University, Portland, OR, USA.

2709 Gabler, K. I., J. W. Laundré, and L. T. Heady. 2000. Predicting the Suitability of Habitat in
2710 Southeast Idaho for Pygmy Rabbits. *The Journal of Wildlife Management* 64:759–764.

2711 Hagar, J. C., and G. Lienkaemper. 2007. Pygmy Rabbit Surveys on State Lands in Oregon. US
2712 Department of the Interior, US Geological Survey.

2713 Kagan, J., K. Zaret, J. Bernert, E. Henderson, E. Gaines. 2018. 2018 Oregon Habitat Map.
2714 Oregon Biodiversity Information Center, Portland State University, Portland, Oregon, USA.

2715

DRAFT

2716 WETLANDS

2717 Wetlands are habitats that are inundated or saturated by surface water or ground water at a
2718 frequency and duration sufficient to support vegetation typically adapted for life in sodden
2719 soil conditions. While dominated by periods of inundation, the natural ecological cycle
2720 may also include dry intervals. Permanently wet habitats include backwater sloughs,
2721 oxbow lakes, peatlands, and marshes, while periodically wet habitats include seasonal
2722 ponds, vernal pools, and wet prairies.

2723 ECOREGIONS

2724 Wetlands are identified as a Key Habitat in all inland ecoregions.

2725 CHARACTERISTICS

2726 Wetlands are characterized by the presence of water, specific types of vegetation, and soil
2727 conditions. Wetland habitats are highly diverse and include the following general types,
2728 which can be distinguished by differences in their hydrology, vegetation communities and
2729 soils:

2730 **Alkaline wetlands** occur in depressions in more arid areas and are intermittently
2731 saturated. An impermeable soil layer prevents water from percolating through the soil,
2732 concentrating salts in some areas. Soil salinity varies greatly by soil moisture and type and
2733 affects the composition of plant species. Plant species are tolerant of saline conditions
2734 due to the concentration of salts by water evaporation. Alkaline wetland vegetation
2735 includes salt-tolerant grasses, rushes, sedges, and shrubs such as black **greasewood**.
2736 Examples of this habitat type are found in the Klamath Lake and Goose Lake areas of
2737 the **East Cascades** ecoregion, and in the **Northern Basin and Range** , **Blue Mountains**,
2738 and **Columbia Basin** ecoregions.

2739 **Deciduous swamps and shrublands** occur in depressions, around lakes or ponds, or on
2740 river terraces. They generally flood seasonally with nutrient-rich waters and are dominated
2741 by woody vegetation, including willows, hardhack, alder, red osier dogwood, Pacific
2742 crabapple, and ash.

2743 **Marshes (including emergent marshes)** occur in depressions, fringes around lakes, and
2744 along slow-flowing streams, especially in valley bottoms. Marshes are seasonally or
2745 continually flooded and have water-adapted plants, such as sedges, bulrushes,
2746 spikesedges, rushes, cattails, and floating vegetation. Marshes can have mucky soils,
2747 resulting in water with high mineral content and vegetation dominated by herbaceous

2748 species. Saltmarshes and tidal marshes are flooded and drained by tides. In brackish
2749 estuaries, they provide habitats for both freshwater and marine fish and wildlife.

2750 **Off-channel riverine habitats**, such as oxbow lakes, stable backwater sloughs, and
2751 flooded marshes, are created as rivers change course. They have less current than the
2752 main channel, with slower-moving or standing water. These areas provide important rearing
2753 habitats for young fish as well as refuge from high flow events, especially during the
2754 migration of young salmon to the ocean. These habitats may also support an array of
2755 aquatic plants, marsh grasses, and terrestrial vegetation.

2756 **Seasonal ponds and vernal pools** hold water during the winter and spring but typically dry
2757 up during the summer months. Vernal pools occur in complexes of networked depressions
2758 that are seasonally filled with rainwater. They host a variety of plant and animal species
2759 with unique adaptations. These habitats can be very important for native invertebrate
2760 species (e.g., **vernal pool fairy shrimp**), plants (e.g., **big-flowered woolly**
2761 **meadowfoam**, **Cook's desert parsley**), **amphibians**, and **birds**. For example, native
2762 amphibians may be able to reproduce in the short time frames when water is present in
2763 seasonal ponds, while invasive non-native bullfrogs cannot. This reproductive advantage
2764 can help native amphibians that are sensitive to competition and predation from bullfrogs.
2765 Drying vernal pools can provide nesting habitat for **streaked horned larks**.

2766 **Wet meadows (including montane wet meadows)** occur on gentle slopes near stream
2767 headwaters, in mountain valleys, bordering lakes and streams, near seeps, in large river
2768 valley bottoms, and in open wet depressions among montane forests. Montane wet
2769 meadows may have shallow surface water for part of the year, are associated with
2770 snowmelt, and are not typically subjected to disturbance events such as flooding. Wet
2771 meadows are dominated by tufted hairgrass, sedges, certain types of grasses, spikesedge,
2772 rushes, and wildflowers.

2773 **Wet prairies (including wet rock outcrops)** occur in lowlands, especially in floodplains,
2774 whereas wet meadows occur in depressions surrounded by forests and are associated with
2775 snowmelt. Wet prairies are dominated by grasses, sedges, and wildflowers including
2776 camas.

2777 **[Spotlight] Beaver Habitat and Beaver Modified Habitat**

2778 Beavers are widely distributed across Oregon SWAP Key Habitats statewide, including
2779 Flowing Water & Riparian, Wetlands, and Aspen Woodlands.

2780 **Beaver habitat**, or habitat for beaver, is the specific combination of water, food, cover, and
2781 space that beaver need to support their survival on the landscape through time. Beaver are

2782 semi-aquatic species that require still or slow-moving, perennial water at stable depths for
2783 cover, protection from predators, access to food resources, and food storage in the winter.
2784 Beavers are slow on land and prefer to forage within 100 feet of their water source. They
2785 need sufficient early seral stage stream buffers of deciduous and herbaceous riparian
2786 vegetation for food and foraging activities. Beavers are highly territorial and require
2787 adequate habitat quality and stability to support their occupancy on the landscape. In
2788 rivers and stream networks, one beaver family unit needs approximately 0.5 to 1.5 linear
2789 stream miles for ample space to survive, reproduce, and thrive. **Beaver habitat**, habitat for
2790 beaver, supports the building blocks that beaver need to create **beaver-modified habitats**,
2791 or habitat by beaver.

2792 **Beaver-modified habitat**, or habitat by beavers, are the specific conditions beaver create
2793 when they alter their terrestrial and aquatic habitat to improve their fitness and survival.
2794 These habitat modifications include denning, damming and ponding water, creating canals
2795 or side-channels, importing woody and vegetative materials into flowing water and
2796 wetlands, and changing the structure of riparian vegetative communities. This suite of
2797 habitat modifications and their cumulative effects can provide benefits such as increased
2798 complexity and connectivity of Key Habitats and habitat, structure, and refugia for SGCN.
2799 Nevertheless, beaver activity can also result in flooding, loss of vegetation, economic loss
2800 on working lands, and conflict with private landowners. Actions focused on beaver habitat
2801 and beaver-modified habitats should also include efforts to mitigate negative impacts and
2802 reduce potential conflicts.

2803 Factors that are currently limiting habitat for beaver include declining surface water
2804 availability, altered floodplain disturbance regimes, conversion and loss of wet meadow
2805 and wetland habitats, and altered riparian vegetation communities.

2806 **ECOREGIONAL CHARACTERISTICS**

2807 **Blue Mountains**

2808 In the Grand Ronde and Baker Valleys, much of the lower elevation wetlands have been
2809 drained and converted to agriculture. Most remaining wetlands in this ecoregion are found
2810 in high-altitude basins and depressions, although some important valley bottom wetlands
2811 occur along rivers and streams, in floodplains, and in basins and depressions that collect
2812 runoff or groundwater. **Ladd Marsh Wildlife Area** offers an example of the importance of
2813 intact wetland habitat for wetland dependent species.

2814 **Coast Range**

2815 Salt marsh habitats provide vital stopover points for migratory birds and support a diversity
2816 of marine species. Wetlands in this ecoregion are vulnerable to development, especially as
2817 coastal populations grow. The ecological processes that create coastal wetlands, such as
2818 landslides, beaver activity, or logjams blocking streams, often are not compatible with
2819 current land uses, especially in more developed areas. Early planning that allows for
2820 appropriate riparian buffers along coastal rivers and streams can maintain many important
2821 wetland and stream functions, including flood control, water retention and storage,
2822 shading, and decreased contaminant inputs. Many of these functions will help to maintain
2823 higher stream flows and lower water temperatures in months with less precipitation. The
2824 Coast Range ecoregion is also home to **Darlingtonia State Park** where a serpentine
2825 wetland has a population of rare *Darlingtonia* (pitcher plants). This park demonstrates the
2826 variety of wetland habitats and the plants that inhabit them.

2827 **Columbia Plateau**

2828 Historical wetlands along the Columbia River have been inundated by reservoirs, while
2829 floodplain wetlands along the Umatilla and Walla Walla rivers and other tributary streams
2830 have mostly been developed for agriculture. This ecoregion once had extensive springs and
2831 vernal pools, many of which have been lost as water tables lowered. Currently, many
2832 wetlands in this ecoregion are man-made, such as marshes established along the edges of
2833 reservoirs and wetlands created as a result of crop irrigation practices. The **Wanaket**
2834 **Wildlife Area**, managed by the Confederated Tribes of the Umatilla, is a network of
2835 wetlands created through irrigation of pastureland that provides important habitat for many
2836 wetland-dependent species. Similarly, ponds on the Umatilla National Wildlife Refuge use
2837 runoff from the fish hatchery to support seasonally wet shallow pools for migrating
2838 shorebirds and to provide breeding habitat for amphibians. Irrigation wetlands in this
2839 ecoregion can provide important habitat but can also be adversely impacted by runoff
2840 containing fertilizers or other chemicals.

2841 **East Cascades**

2842 The upper Klamath Basin once had an extensive shallow lake and marsh system, but much
2843 of that system has been lost due to drainage and conversion to agriculture and urban uses.
2844 These changes have contributed to the complex issues surrounding water use and species
2845 conservation in the basin. The remaining wetlands in the Klamath Basin support one of the
2846 largest concentrations of waterfowl in North America, with millions of ducks, geese, and
2847 other waterfowl migrating through the basin annually. In the winter, the Klamath Basin
2848 hosts the largest assemblage of wintering Bald Eagles in the continental United States. The
2849 Klamath Basin provides Oregon's only permanent nesting areas for **Red-necked**

2850 **Grebes** and most of the nesting areas for **Yellow Rails**. High quality wetland habitats are
2851 also found in the Upper Deschutes River Basin.

2852 **Klamath Mountains**

2853 Most low-elevation, seasonal wetlands have been lost due to conversion to agriculture,
2854 urban and rural residential uses, energy development, and transportation systems. Altered
2855 hydrology and upland activities like groundwater withdrawal impact many remaining
2856 wetland habitats. High elevation wetlands are severely impacted by livestock grazing.
2857 Scarce vernal pool wetlands in the Agate Desert near Medford support several rare plant
2858 and animal species. These vernal pool wetlands are formed in areas with unusual
2859 topography and soil layering and are very difficult to replace when ground is leveled for
2860 development.

2861 **Northern Basin and Range**

2862 The **Northern Basin and Range** ecoregion contains several large, deep freshwater
2863 marshes. Significant wetlands are associated with the large lake basins, including Lake
2864 Abert, Summer, Malheur, and Harney Lakes, and the Warner Basin. However, many of the
2865 ecoregion's smaller historical wetlands have been lost due to conversion or degradation
2866 from stream channelization, water use, water diversions, and historical overgrazing.
2867 Creation of watering holes for livestock and wildlife has altered the hydrology at many
2868 major alkaline wetlands, making them one of the most altered habitat types in the
2869 ecoregion.

2870 In some areas, flood-irrigation of private pasture and hay meadows provides important
2871 seasonal habitat for migrating and breeding birds. Nevertheless, flood irrigation can
2872 negatively impact water quality, increase sedimentation, and increase water loss due to
2873 evaporation. Cooperative projects, such as settling ponds designed for cleaning flood
2874 irrigation "tail water", or conversion to piped sprinkler systems may offer a way to address
2875 water quality issues.

2876 **West Cascades**

2877 Wetlands in this ecoregion are generally in good condition, although some areas, such as
2878 those located around Mt. Hood and Mt. Jefferson, can be impacted by uncontrolled
2879 livestock grazing, camping, or off-highway vehicle use. **Climate change** projections
2880 estimate that wetland hydrology in the West Cascades could be impacted by shifts in
2881 rainfall and snowmelt and increasing temperatures.

2882 **Willamette Valley**

2883 Almost all remaining wetlands in this ecoregion have been degraded to some degree by
2884 altered water regimes, pollution, and invasive plants and animals. The Willamette Valley
2885 ecoregion also used to have extensive networks of off-channel habitat, deciduous swamps
2886 and shrublands, marshes, seasonal ponds, and vernal pools. However, most of these
2887 habitats have been lost to agriculture and urbanization.

2888 Wetlands in the Willamette Valley serve important ecological functions for communities,
2889 provide habitat for amphibians, reptiles, birds, and fish, and offer key bird and fish
2890 migratory pathways. Once an abundant ecosystem within the Willamette Valley, native
2891 wetland prairies have declined dramatically in extent since the mid-1800s due to a variety
2892 of factors including a growing human population, agricultural conversion, urbanization,
2893 drainage, and colonization by invasive and woody vegetation. In particular, wetland prairie
2894 habitat is regarded as one of the most imperiled in the Willamette Valley ecoregion.

2895 **CONSERVATION OVERVIEW**

2896 Wetlands and wet meadows provide important habitat for migrating and breeding
2897 shorebirds, waterbirds, waterfowl, songbirds, invertebrates, mammals, amphibians, and
2898 reptiles. Floodplain wetlands and backwater sloughs and swamps are important rearing
2899 habitats for juvenile salmon. Wetlands have direct value for people because they improve
2900 water quality by trapping sediments and pollutants, recharge aquifers, store water and
2901 carbon, stabilize erosion, and reduce the severity of floods. Seasonal wetlands that dry up
2902 during the summer provide important ecological functions, such as supporting water
2903 quality and sequestering carbon.

2904 With most wetlands in private ownership, working with landowners to restore and manage
2905 wet meadow systems and other wetlands can increase sustainable production of forage for
2906 livestock and increase late-season stream flows while also providing fish and wildlife
2907 habitats. Restoration and retention of wetlands, especially those with high water storage
2908 potential in arid regions, may help to maintain ground water levels into drought periods.
2909 Even when converted to pasture ponds, highly productive off-channel habitat is present if
2910 hydrologic connections are maintained.

2911 In general, wetlands across the state face a range of pressures from human activity and
2912 environmental changes. Wetland loss is primarily due to land conversion for agricultural,
2913 urban, rural, and energy development, or infrastructure projects. Most wetland habitat loss
2914 has occurred at lower elevations and valley bottoms. Invasive species can quickly take over
2915 degraded wetlands, leading to loss of function. One of the most pervasive invasive plant
2916 species is reed canary grass, which will dominate a wetland, degrading its ecological

2917 function and outcompeting native species. **Climate change** is expected to affect Oregon's
2918 wetlands through shifting precipitation patterns, increased droughts, more high severity
2919 wildfires, and warmer temperatures.

2920 **[SPOTLIGHT]** White-faced Ibis

2921 The White-faced Ibis is a colonial breeding bird that breeds in semi-permanent wetlands
2922 that are regularly impacted by drought and floods. This bird is highly nomadic, allowing it to
2923 compensate for poor conditions at traditional colony sites by moving between years to new
2924 breeding locations, resulting in local population fluctuations and colony abandonment in
2925 response to system dynamics.

2926 Oregon has historically been peripheral to the core of the range of the White-faced Ibis in
2927 the intermountain west, recorded sporadically during the 19th century and with the first
2928 documentation of a breeding colony in the state in 1908. Breeding colonies were
2929 established in Oregon periodically in the following decades, becoming more common
2930 towards the end of the 20th century with an estimate of about 4000 pairs in Oregon in the
2931 early 90s. Prior to 1984, most of Oregon's ibises were located in central Malheur Lake,
2932 though colonies have since been documented in Lake, Harney, and Klamath Counties.

2933 This nomadic nature highlights the importance of considering the regional context in
2934 management decisions and population monitoring, to allow land managers to understand
2935 the bigger picture of how wetland management in their area relates to the whole. With
2936 increasing impacts from climate change and mega-droughts throughout the intermountain
2937 west, wetland conservation in Oregon may become more and more important to the
2938 conservation of White-faced Ibises within the Great Basin.

2939 **LIMITING FACTORS AND RECOMMENDED APPROACHES**

2940 **Limiting Factor: Habitat Loss**

2941 A high percentage of low-elevation and valley bottom wetlands have been lost or degraded
2942 through diking and draining, particularly in the Klamath Mountains and Coast Range
2943 ecoregions. In other areas, overgrazing has led to soil compaction, changes in plant
2944 species composition, and spread of invasive plants. Due to short growing seasons and
2945 other factors, degraded wet meadows can be slow to recover if overgrazed. Saltmarshes
2946 have experienced substantial losses over historical condition from diking, installation of
2947 tide gates, draining, and filling of tidally influenced marshes. Wetlands provide vital habitat
2948 for migrating shorebirds and waterfowl. Loss or degradation of wetland habitat in the
2949 Pacific Flyway could potentially have large impacts on bird populations while early season

2950 haying in wetland habitats can result in poor reproduction of ground-nesting birds due to
2951 destruction of nests and direct mortality of young.

2952 Many wetlands are lost through urbanization, which involves filling or draining the wetland
2953 for development. Unfortunately, this removes wetlands from locations where the functions
2954 they provide might have the most value for SGCN. Maintaining wetland and adjacent
2955 habitats provides social benefits, such as storage of flood water and treatment of
2956 contaminants before reaching streams and ground water.

2957 **Recommended Approach**

2958 Protect and conserve priority wetland habitat that provides vital breeding habitat
2959 for **Species of Greatest Conservation Need** and stopover sites for migrating species
2960 (KCI: **Barriers to Animal Movement**). Identify wetlands that have been altered or lost and
2961 determine their potential for restoration. Build upon current cooperative efforts to maintain
2962 and restore wetlands in partnership with private and public landowners. **Cooperative**
2963 **voluntary approaches** are important for wetland conservation on private lands. Continue
2964 to provide incentives to protect, maintain, or restore wetlands, such as the **Wetland**
2965 **Reserve Enhancement Partnership (WREP)** offered through the Natural Resources
2966 Conservation Service and private mitigation banking. Prioritizing development of **wetland**
2967 **mitigation banks** to support SGCN also provides a strategic landscape approach to
2968 addressing wetland loss.

2969 Develop and implement grazing regimes that are compatible with wet meadow
2970 conservation objectives. Use cooperative efforts and incentive programs to establish semi-
2971 permanent livestock exclusion zones in priority areas. In partnership with landowners,
2972 implement later haying dates in critical bird nesting areas (see **The Willamette Valley**
2973 **Landowner's Guide to Creating Habitat for Grassland Birds**). Manage beaver
2974 populations to contribute to wetland creation and maintenance, when compatible with
2975 existing land uses.

2976 Promote outreach and education programs to educate individuals, communities, city and
2977 county planners, agricultural groups, and forest industries about the function and services
2978 provided by wetlands. Work with the **local planning process** and the **Oregon Department**
2979 **of State Lands** to promote the value of maintaining wetlands and habitat corridors,
2980 especially along floodways, where they can best function to protect structures,
2981 infrastructure, and water quality.

2982 **Limiting Factor: Drought**

2983 Drought affects the quality and extent of wetlands across the state. Drought has resulted in
2984 less precipitation, interception, infiltration, and percolation of water into the soil, falling
2985 water tables, increased evaporation, decreased transpiration, decreased plant and animal
2986 diversity and distribution, and the acidification, cracking, and compaction of wetland soils.
2987 These changes in hydrology and soils affect plant and animal diversity, connectivity with
2988 other aquatic resources and upland habitats, the proportion of invasive plant species in the
2989 wetland, and wetland functions. Climate change patterns are expected to exacerbate and
2990 or extend drought periods resulting in wetland loss of acreage and functional changes.

2991 **Recommended Approach**

2992 Educate the public and water users to conserve water. Look for opportunities to restore
2993 and enhance wetlands (i.e. Barnes and Agency Wetland Restoration- Upper Klamath
2994 National Wildlife Refuge), reuse water for multi-benefit solutions, and improve wetland
2995 habitat through water use efficiencies (i.e. Lower Klamath and Tule Lake National Wildlife
2996 Refuges). Coordinate with ODFW Fish Passage and District Fish Biologist staff on
2997 appropriate approvals for instream, beaver-mimicry (e.g., beaver dam analogs, small
2998 and/or large wood structures) and/or coexistence (e.g., pond levelers, culvert exclusion
2999 devices) structures to raise the water table, restore wetland and waterway connections,
3000 and improve habitat conditions (i.e. Sprague Watershed).

3001 **Limiting Factor: Water Quantity**

3002 Water is extremely limited in much of the Blue Mountains, East Cascades, and Northern
3003 Basin and Range ecoregions. As a result of water availability, there is competition for water
3004 resources, particularly in late summer. Lowered water tables affect wetland habitats.
3005 Competition for water harms both ecological and economic goals. Water diversions for
3006 other uses change the seasonality of flooding, slow habitat recovery, and increase invasion
3007 of non-native grasses. Drought years intensify water shortages.

3008 **Recommended Approach**

3009 Use cooperative efforts and incentive programs, such as financial incentives for wetlands
3010 restoration, water rights acquisition, and wetland mitigation banking, to manage water
3011 allocation and wetland habitats. Implement water conservation actions, where possible, to
3012 increase availability (quantity, timing, and duration).

3013 **Limiting Factor: Degraded Water Quality**

3014 Although wetlands have a role in purifying water, water quality is poor in some wetland
3015 systems. High temperatures affect water quality in some areas. Non-point source runoff
3016 from agricultural and residential areas contains pollutants that can affect water quality and
3017 nutrient levels, and these levels may increase as water evaporates throughout the season.
3018 High nutrient loads can contribute to toxic algal blooms.

3019 **Recommended Approach**

3020 Provide incentives to decrease and manage the release of potential contaminants, such as
3021 fertilizers or pesticides, by controlling the timing of application. Use incentives to promote
3022 substitutes that are less toxic to wildlife and break down quickly in the environment.
3023 Promote the creation of stormwater treatment projects, fencing of aquatic habitats to
3024 exclude livestock, and restoration of riparian buffers and additional wetlands to increase
3025 filtering capacity. Support irrigation systems that conserve, re-collect, and re-use water
3026 more effectively, use gray water, and provide shaded treatment areas that can provide
3027 cooling and habitat. In the Willamette Valley, adopt critical actions recommended by
3028 the **Willamette Restoration Initiative** on Clean Water, such as: reduce the levels of toxins
3029 and other pollutants in the Willamette Basin, provide incentives to decrease water
3030 pollution, and promote education and outreach programs for landowners.

3031 **Limiting Factor: Invasive Species**

3032 Invasive species, such as reed canary grass, purple loosestrife, ludwigia and Japanese
3033 knotweed, invade and degrade wetlands, thereby displacing native plants, reducing plant
3034 community diversity, reducing sources of food for wildlife, and altering water flow and
3035 storage function. Invasions of non-native grasses, such as reed canary grass, can also
3036 create conditions more prone to wildfires.

3037 Invasive, non-native carp can impact wetlands by consuming important plants and by
3038 increasing turbidity, disturbing sediments, and altering biological dynamics for sediment-
3039 associated plants and animals. Turbidity also contributes to higher water temperatures and
3040 lower levels of dissolved oxygen. Non-native bullfrogs have had a devastating impact on
3041 native amphibians and reptiles, leading to the extirpation of Oregon spotted frog and
3042 leopard frogs from much of their historic range. Emerald ash borer is now present in the
3043 Willamette Valley and threatens to cause extensive losses to ash trees, which are a critical
3044 component of off-channel wetland habitats.

3045 Native trees and shrubs can become invasive due to the exclusion of fire from wetlands.
3046 Without fires from natural ignitions and Indigenous peoples cultural burning practices,

3047 encroachment by native vegetation can overwhelm wetlands and out compete wetland
3048 grasses and flowers by reducing water availability, shading, and changing soil chemistry.

3049 **Recommended Approach**

3050 Emphasize prevention, risk assessment, early detection, and quick control to prevent new
3051 invasive species from becoming fully established. Control key invasive plants using site-
3052 appropriate tools, such as flooding (reed canary grass), biological control (purple
3053 loosestrife), and mechanical treatment including mowing. Use chemical treatment
3054 carefully and where compatible with water quality concerns, focusing on spot treatment
3055 during the dry season. Consider screening or adjusting water levels to control carp. Use
3056 revegetation and other means to establish and maintain native plant communities that are
3057 relatively resistant to invasion and that also meet other land use objectives.

3058 **RESOURCES FOR MORE INFORMATION**

3059 **Oregon Department of State Lands, Wetlands**

3060 **Oregon Wetland Program Plan**

3061 **Oregon Statewide Wetlands Inventory**

3062 **Practical Guidelines for Wetland Prairie Restoration in the Willamette Valley,**
3063 **Oregon: Field-tested Methods and Techniques**

3064 **Klamath Wetland Restoration**

3065 **Williamson River Delta Wetland Restoration**

3066 **Harney Basin Wetland Collaborative**

3067 **Partnership for Lake Abert and the Chewaucan**

3068

3069 NEARSHORE HABITATS

3070 Nearshore Habitats encompass the coastal and marine habitats in the area from the 3
3071 nautical mile outer limit of Oregon's territorial sea, where water depths average 66 m and
3072 range from 17 m to 194 m (56 to 308 ft), to the supratidal areas of the shoreline affected by
3073 wave spray and overwash at extreme high tides.

3074 Nearshore Habitats describe the Coastal and Marine Ecological Classification Standard
3075 (CMECS) habitat classification approach (see **Appendix - Marine Habitat Classification**)
3076 for the major habitat types found in Oregon's nearshore, including neritic, soft bottom
3077 subtidal, rocky subtidal, rocky shore, sandy beaches. For information on Estuaries, which
3078 also occur in the nearshore, see the **Estuaries** Key Habitat.

3079 ECOREGIONS

3080 Nearshore Habitats are a Key Habitat in the **Nearshore Ecoregion**.

3081 CHARACTERISTICS

3082 **Neritic Habitat (Open Water)**

3083 The neritic habitat encompasses the waters and biological communities over the
3084 continental shelf, including nearshore and offshore marine subsystems as defined by
3085 CMECS. It spans the surface, upper water column, pycnocline, and lower water column,
3086 extending westward to the continental shelf break at about the 200 m depth contour.
3087 Constantly in motion, this habitat is shaped by the California Current System, seasonal
3088 upwelling and downwelling, El Niño/La Niña events, and the Pacific Decadal Oscillation,
3089 which all drive water movement across various time scales.

3090 The CMECS biotic component identifies planktonic biota as the primary setting, with
3091 species composition varying by water mass. The ecology of the neritic habitat is affected by
3092 processes taking place at scales varying from global to local. The dynamics of the neritic
3093 habitat affect all the other habitats described later in this section.

3094 *Physical Environment*

3095 Many physical and chemical environmental factors affect neritic ecology. These factors
3096 include but are not limited to solar light and radiation influence, salinity, temperature, layer
3097 position, physical mixing, hydrostatic pressure, biogeochemical composition, atmospheric
3098 exposure and influence, surface and underwater currents, swells, waves, and water mass
3099 movements. Many of these factors can change by location and time of year. The neritic

3100 habitat encompasses many water column habitats that shift, expand, and contract over
3101 time and space in both predictable and stochastic patterns.

3102 Coastal upwelling is perhaps the most defining feature of Oregon's neritic habitat with its
3103 alternating upwelling-relaxation events. Upwelling is a water column hydroform, described
3104 by CMECS as an upwardly-directed current caused by divergence of water masses. In
3105 spring and summer months, strong northerly winds push surface and upper water layers
3106 westward towards the deep ocean. This movement causes deep, cold, oxygen-poor but
3107 nutrient-rich waters to rise to the surface near the coast replacing the water that was driven
3108 offshore. These nutrients, brought to the upper layers of the water column, help propagate
3109 and sustain the rich biota of Oregon's coastal waters. The relaxation events, when the
3110 northerly winds briefly cease or reverse, allow the upper water layer to move back towards
3111 shore bringing its rich biotic content with supplies of food, larvae, and juvenile organisms.
3112 In the fall and winter months when winds blow predominantly from the south, the surface
3113 and upper water layers move shoreward and downward in a process called downwelling.
3114 Downwelling is an important part of the annual seasonal cycle that forces oxygen rich
3115 waters from the upper layers downward in the water column. Surface water temperatures
3116 provide a good indication of these seasonal wind forcing differences that bring the cold,
3117 nutrient-rich waters to the surface in the summer and the warmer waters from offshore to
3118 the coast in the winter (see **Nearshore Ecoregion**).

3119 Large-scale changes in water masses, temperatures and currents result in changes in
3120 plankton species composition and abundance, which impact the survival and distribution
3121 of organisms within coastal and oceanic ecosystems. These large-scale oceanic events,
3122 such as El Niño/La Niña and the Pacific Decadal Oscillation, occur at multi-year or decadal
3123 time scales.

3124 Another water column component that affects Oregon's neritic habitats is river plumes.
3125 CMECS does not characterize the marine waters affected by these plumes as estuarine
3126 because they are not meaningfully enclosed by landforms. Riverine waters entering the
3127 ocean often carry high concentrations of nutrients, create gradients in salinity, cause
3128 physical mixing, and create areas of high turbidity. Large river plumes, such as that from
3129 the Columbia River, may serve as a microhabitat within neritic habitats and can potentially
3130 act as biogeographic barriers between marine areas to the north and south. The Columbia
3131 River plume stretches hundreds of miles offshore and shifts predictably over the course of
3132 each year. In the summer the plume spreads south and offshore from the river's mouth,
3133 while during the winter the plume is found to the north of the river mouth and is usually
3134 directly adjacent to the coast. This plume has important ecological effects, not only to
3135 neritic habitats, but to nearshore and offshore habitats as well. The oceanographic fronts
3136 created by the Columbia River plume in the marine systems generate productive
3137 conditions that attract many species of invertebrates, fish, seabirds, and marine
3138 mammals.

3139 *Biological Characteristics*

3140 Neritic habitats support two basic types of marine organisms: plankton and nekton.
3141 Planktonic organisms live in the water column and are incapable of swimming against
3142 currents, instead drifting with them. Plankton are often categorized as either phytoplankton
3143 or zooplankton. Phytoplankton are microscopic photosynthesizing organisms (e.g.,
3144 diatoms), and are the primary producers that form the base of the marine food web. Huge
3145 surges in phytoplankton populations, known as “blooms,” are commonly associated with
3146 upwelling events. Zooplankton are heterotrophic organisms that range in size from
3147 microscopic single-celled organisms to enormous jellyfish a meter or more in diameter.
3148 Some plankton, called holoplankton, like many diatoms, copepods, krill and jellyfish spend
3149 their entire lives as drifters in the water column. Many species like sea urchins, mussels,
3150 crabs, some snails and many fishes have planktonic stages as eggs or larva, called
3151 meroplankton, before either settling to the bottom or growing large enough to be nekton.
3152 The CMECS biotic component uses these planktonic classes and subclasses to describe
3153 the open water neritic zone. They can be further refined by taxonomic groups and
3154 communities that are dominant in any given area of interest. Dramatic changes in plankton
3155 communities occur in Oregon waters with water masses changes. For example, warm
3156 water species are brought into nearshore water with El Niño events.

3157 In contrast, nektonic marine organisms are capable of swimming against currents and
3158 include animals such as adult crustaceans, mollusks, and vertebrates. Highly migratory
3159 and schooling species are typical of nekton in neritic habitats. Many species of
3160 invertebrates, fish, birds, and marine mammals travel and forage within this habitat.

3161 Many nearshore Species of Greatest Conservation Need (SGCN), Watch List and
3162 commonly associated species utilize the open water neritic habitat during their life history
3163 (see **Appendix - Nearshore Species**). Many forage fishes such as northern anchovy,
3164 Pacific herring, topsmelt, Pacific sardine, surf smelt, Pacific sand lance and longfin smelt
3165 feed in this open water neritic habitat. Juvenile rockfish are found in the water column.
3166 Breeding birds such as Tufted Puffin and Common Murre are central place foragers that
3167 feed on the forage fish and other species while nesting. The majority of nearshore SGCN
3168 depend on this habitat for some phase of life. This is also the habitat that supports primary
3169 production by phytoplankton and secondary production by zooplankton, which is at the
3170 base of the food web for the nearshore ecosystem. Ocean currents transport and disperse
3171 larvae and juveniles of many invertebrate and fish species throughout the region.

3172 *Human Use*

3173 Human uses of the neritic habitat include commercial and recreational fishing,
3174 nonconsumptive recreational pursuits such as boating or whale watching, scientific
3175 research, commercial maritime transportation, and military operations. Development of
3176 renewable energy sources from both wind and waves is an emerging use of the neritic
3177 habitat.

3178 **Soft Bottom Subtidal**

3179 Soft bottom subtidal habitat includes all of the unconsolidated substrate areas (e.g., mud,
3180 sand, granule pebbles and various mixes thereof) on the ocean bottom. Soft bottom
3181 subtidal habitats are characterized by CMECS as being within the subtidal zones of the
3182 nearshore and offshore marine subsystems. Subtidal soft bottom habitats are diverse
3183 based on distinct organism assemblages that are influenced by differences in substrate
3184 type (sand vs. mud), organic content and bottom depth. The distribution and relative
3185 abundance and mixes of these substrates are not yet well described for much of Oregon's
3186 nearshore ocean waters.

3187 *Physical Environment*

3188 The primary substrate types in Oregon's soft bottom subtidal areas range from sand to
3189 pebble. CMECS defines unconsolidated mineral substrates based on particle diameter.
3190 Here we consider soft bottom habitats to be composed of the various mixes defined by
3191 CMECS of particles <64 mm in diameter. Because the Oregon coast is primarily an
3192 exposed, high-energy environment, most soft bottom subtidal areas are sandy. However,
3193 mud can be the more prevalent substrate type in areas receiving less energy from water
3194 movement, including isolated and sheltered areas, and deeper areas. The distribution of
3195 these unconsolidated sediment types in Oregon waters is influenced by currents in both
3196 the nearshore and offshore subsystems. Areas close to outfalls and discharge pipes would
3197 be expected to show localized differences based on the displacement of substrate and the
3198 increased availability of organic and small particulate material. The smaller the particle
3199 size, the smaller the pores (or spaces between the particles) are. Pore size dictates the
3200 amount of water and the water chemistry of the substrate, which can define what types of
3201 organisms can live in that sediment.

3202 *Biological Characteristics*

3203 Most soft bottom subtidal communities are dominated by infaunal (burrowing)
3204 invertebrates such as polychaetae worms. However, other organisms such as crustaceans,
3205 echinoderms and mollusks may be locally abundant. Common epifauna (found on the
3206 sediment surface) can include species of shrimp, crabs, snails, bivalves, sea cucumbers,
3207 and sand dollars. Dungeness crab are an important component of soft bottom subtidal
3208 communities and are found both on the surface as well as buried in the substrate. Sea
3209 pens (*Ptilosarcus* sp.), colonial relations to sea anemones, are common on more muddy
3210 bottoms. In some areas of the coast, shallow sandy habitats support extensive beds of
3211 dense sand dollars that may extend miles in length. Common fish in this area include
3212 several species of flatfish (e.g., sanddab, English sole, and sand sole), and important
3213 burrowing forage species such as Pacific sand lance and sandfish.

3214 Species associated with soft bottom subtidal habitats provide a spectrum of ecosystem
3215 services. Most widespread but least apparent of these services are the nutrient cyclers:

3216 deposit feeders and microbes living within the sediments. Emergent species such as sea
3217 pens are only found in this habitat. There are a vast array of worms and other invertebrates
3218 that live in the soft subtidal bottom. Soft bottom habitats are important to many SGCN,
3219 Species of Greatest Information Need (SGIN), Watch List and other commonly associated
3220 species at various life stages (see **Appendix - Nearshore Species**). For example, big skate,
3221 starry flounder, sand sole, Pacific sand lance burrow or cover themselves to hide in these
3222 sediments. Gray whales feed by sifting buried amphipods from the sediments and
3223 scooping clouds of mysid shrimp from above the sediment surface sometimes at the edges
3224 of rocky reefs. Many invertebrates like razor and native littleneck clams live in the subtidal
3225 soft bottom habitat. Both juvenile and adult Dungeness crab forage here and sometimes
3226 hide in these soft sediments. The young of commercially valuable fish species can often be
3227 found here and utilize these areas as nursery habitat. The young of many species use the
3228 nearshore area for foraging and are themselves prey for larger fishes and birds. Sand lance
3229 is a particularly valuable forage species for birds, other fishes, and marine mammals.
3230 Diving birds such as the Common Murre forage for food for their young in soft bottom areas
3231 taking juvenile flat fish back to their chicks while they are nesting.

3232 *Human Use*

3233 Commercial and recreational harvest of Dungeness crab, surf perch, and species of
3234 nearshore flatfish are the principal human uses of the soft bottom subtidal habitat. Sand
3235 and mud from dredging projects are sometimes deposited over soft bottom habitats. Soft
3236 bottom subtidal habitats could also soon be utilized for siting renewable energy projects
3237 and their associated infrastructure. Finally, the soft bottom subtidal offers many
3238 opportunities for scientific research

3239 **Rocky Subtidal**

3240 Rocky subtidal habitat includes all hard substrate areas of the ocean bottom. The geologic
3241 origin substrate components include cobble and boulder in the CMECS unconsolidated
3242 mineral substrate class and bedrock and megaclasts in the rock substrate class.
3243 Anthropogenic origin hard substrates are also here. Anthropogenic reefs include any areas
3244 where hard, persistent material has been placed either purposely or accidentally by
3245 humans. Examples include rock jetties at the entrance to many bays, shipwrecks,
3246 anchoring systems for renewable energy projects, and unburied portions of underwater
3247 cables or pipelines. Rocky subtidal areas are often referred to as reefs, rocky reefs, rocky
3248 banks, pinnacles, or “hard bottom.” Rocky subtidal habitats, including both the natural and
3249 anthropogenic components, are characterized by CMECS as being within the subtidal
3250 zones of the nearshore and offshore marine subsystems. Although most areas are never
3251 exposed to air, the CMECS subtidal definition does include areas that are exposed
3252 intermittently each month when tide levels fall below the Mean Lower Low Water (MLLW)
3253 level. Rocky subtidal habitats are found in both the nearshore subsystem and offshore
3254 subsystem and some of the differences are discussed below.

3255 Some rocky subtidal areas are extensions of shoreline rocky features such as headlands,
3256 cliffs, or rocky intertidal habitat, while others exist as isolated regions of rock surrounded
3257 by habitat with soft bottom substrate. Rocky reefs have varied topography; some may
3258 barely come above the surrounding seafloor, while others may rise from the seafloor many
3259 meters, or extend above the surface to form islands in the Territorial Sea. There are more
3260 than 1,800 islands off the coast of Oregon, the bases of which form rocky subtidal habitat.

3261 *Physical Environment*

3262 The physical characteristics of rocky subtidal habitats reflect proximity to shore, depth of
3263 the water, local seafloor geology, erosional forces, and biological influences. The geology of
3264 many rocky subtidal areas mimics the geology of adjacent landforms, often consisting of
3265 erosion-resistant basalts or metamorphic rock common in Oregon's rocky headlands. Over
3266 geologic time, the underwater rock features have been uplifted, bent, deformed, and
3267 alternately exposed to ocean and terrestrial erosional forces as successive ice ages and
3268 geologic forces caused massive sea level changes. These forces have shaped a variety of
3269 physical habitat features within reefs, including flat rocky benches, stacks, jagged ridges,
3270 broken boulder fields, and a vast number of cracks and crevices that provide shelter and
3271 substrate to abundant life.

3272 Oceanographic processes and features strongly influence the rocky subtidal environment.
3273 Subtidal reefs are exposed to pounding wave action, underwater currents, and the physical
3274 and chemical properties of the water. These factors in turn influence the biological
3275 community on the reefs. Generally, nearshore reefs are more exposed to wave action than
3276 offshore reefs, and the wave action is much stronger in winter than during summer. Wave
3277 action is a key factor in determining the types of organisms that can live on the very shallow
3278 reefs. Ocean currents vary widely by location, time of year, and over tidal cycles. Currents
3279 influence reefs in a variety of ways including direct erosion, sand scour or burial of reef
3280 areas, and movement of organisms to and from reefs, including plankton and larva. Large-
3281 scale or long-term variation in the ocean environment such as upwelling, seasonal current
3282 directional shifts, shifts in ocean circulation, water temperature variation, local and global
3283 weather patterns, ocean acidification, and biological processes combine to determine the
3284 ambient chemical and physical composition of the water in rocky subtidal habitats. The
3285 CMECS water column components can be used to describe important features of the
3286 waters surrounding and overlying rocky reefs that are important in shaping the biological
3287 communities which live there.

3288 The 30 m depth contour is defined by CMECS as the boundary for the nearshore subsystem
3289 and the offshore subsystem. Nearshore rocky reefs differ from offshore reefs in some key
3290 physical characteristics. Light penetration is adequate to support algal life on nearshore
3291 reefs, while offshore reefs support far less algal growth. For example, kelp is only found in
3292 nearshore subsystem rocky areas. Wave action, currents, and storms produce a higher
3293 energy environment on nearshore reefs than their deeper counterparts. Organisms
3294 adapted to higher energy environments are more prevalent in the nearshore area. On some

3295 reefs, strong currents can scour and seasonally bury or expose the rocks with sand,
3296 considerably influencing the types of organisms that can utilize those rocky subtidal
3297 environments.

3298 *Biological Characteristics*

3299 Subtidal rocky reefs are known for their abundant and diverse biological communities. The
3300 variety in topography, substrate characteristics, and depths within and among rocky reefs
3301 produces a plethora of microhabitats, often within relatively small geographic areas. This in
3302 turn provides for a diversity of species adapted to life in these different microhabitats.
3303 Habitat-forming organisms, such as kelp or attached invertebrates, provide additional
3304 microhabitats used by reef species.

3305 Most nearshore rocky reefs have rich algal, invertebrate, fish, bird, and marine mammal
3306 communities. Depending on water depth, light penetration, wave energy, and other
3307 physical and biological processes, algae and macroalgae can provide extensive or sporadic
3308 cover and food for other species in the nearshore subsystem. Algae and macroalgae
3309 include encrusting forms that grow close to the rock surface, turf forms that can create a
3310 dense layer up to a foot thick or more, subcanopy forms that provide added subsurface
3311 habitat structure, and canopy forms that create kelp “forests” which may break the surface
3312 of the water. Offshore rocky reefs in deeper water do not have kelp forests. Free-swimming
3313 (nektonic), drifting (planktonic), and attached invertebrates are common in both the
3314 nearshore and offshore rocky subtidal habitats.

3315 Many Nearshore SGCN, SGIN, Watch List, and other commonly associated species inhabit
3316 rocky subtidal habitats (see **Appendix - Nearshore Species**). These include many fish as
3317 well as a wide variety of filter or suspension feeding invertebrates attach to hard substrates
3318 such as sponges, anemones, barnacles, bryozoans, hydrozoans, tunicates, and cold water
3319 corals. Mobile invertebrates abound here as well. Red and purple urchins, red and flat
3320 abalone eat algae attached to the rocks. Ochre, sunflower and other sea stars forage in
3321 subtidal rocky habitats as do crabs, shrimps, brittle stars, nudibranchs, chitons, and
3322 worms.

3323 The diversity of producers and consumers found in the rocky subtidal creates complex food
3324 webs and interdependence among organisms. Reefs are linked to surrounding
3325 environments by ocean currents and organism movements. Reef topographic structure
3326 often slows currents, enhancing the local community’s ability to capture drifting
3327 organisms, an effect enhanced by the occasional presence of large kelp beds. Many
3328 organisms move on and off reefs, some in large-scale migrations and others in short
3329 feeding forays to other areas. While most nearshore reef fishes occupy both nearshore and
3330 offshore reefs, there are differences in depth preferences of some species and life history
3331 stages.

3332 Several fish species depend on nearshore rocky reefs during early life history stages before
3333 moving off to deeper reefs, the continental shelf, or other areas as they grow. Conversely,
3334 some fish depend on estuaries or rocky intertidal habitat for early life history stages before
3335 moving to rocky subtidal areas as adults. For example, kelp greenling, cabezon, and grass
3336 rockfish tend to be more prevalent on the nearshore reefs. Canary and yelloweye rockfish
3337 move from nearshore to offshore reefs as they grow. Many fish species are entirely
3338 dependent on reefs for parts of their life cycle, while others are visitors. Common visitors
3339 include herring, smelt, sharks, ratfish, and salmon.

3340 Ecological linkages within and between rocky subtidal habitats help to shape their
3341 biological communities and the diversity of species found in this type of habitat. Currents
3342 bring in planktonic organisms and transport drifting larvae to and from disparate rocky
3343 subtidal habitats. The location of reefs with respect to other “upstream” or “downstream”
3344 reefs has a dramatic effect on the types, abundance, and recruitment rates of the reef’s
3345 communities and organisms. This complexity of organism interrelationships makes the
3346 outcome of natural or human disturbance to reefs difficult to measure or predict.

3347 Kelp beds form a small but important subset of Oregon’s rocky subtidal habitat. CMECS
3348 classifies kelp beds as a biotic component of Oregon’s rocky subtidal habitat, and more
3349 specifically as canopy-forming algal beds. Kelp canopies in Oregon consist almost
3350 exclusively of bull kelp (*Nereocystis luetkeana*), a brown macroalgae that grows from the
3351 seafloor to the ocean surface and forms a floating canopy, though a few locations have
3352 historically featured small amounts of giant kelp (*Macrocystis pyrifera*). The strip of coast
3353 from Cape Arago south has historically contained approximately 92 percent of the state’s
3354 kelp beds (Figures 1 and 2). Kelp canopies are relatively scarce habitats in Oregon’s waters,
3355 covering less than one percent of the nearshore area. This distribution is driven historically
3356 by the locations of subtidal rocky seafloor shallow enough for sufficient light penetration to
3357 support kelp growth. More recently, the even more limited distribution reflects reductions
3358 in the total abundance of kelp in response to changes in oceanographic stressors such as
3359 warming ocean temperatures, marine heat waves, changes in ocean chemistry associated
3360 with climate change (kelps need cool, nutrient-rich waters to thrive) and recent increases
3361 in populations of grazing sea urchins (see **Specialized and Local Habitats**).


3362 The presence and attributes of kelp beds depend on a number of physical and biological
3363 variables. The primary variables determining where kelp might exist include water depth
3364 and substrate availability. In Oregon’s waters, kelp beds only form on rocky substrate and
3365 are limited to the nearshore subsystem. Beyond that depth, low light levels on the seafloor
3366 limit the growth of kelp. However, light and substrate are not the only limiting factors; many
3367 rocky reefs in the appropriate depth range rarely or never support kelp beds. Factors that
3368 may limit kelp on these reefs include seasonal sand burial of the reef, sand scour of the
3369 rocks, overexposure to wave and storm energy, locally high turbidity, lack of nutrients,
3370 distance of the reef to “seeding” sources of kelp, abundance of organisms that consume
3371 kelp (e.g., sea urchins), and competition with invertebrates and other algae for rock
3372 substrate available for attachment. Kelp beds in Oregon display pronounced seasonal and

3373 annual variation in extent and density. Bull kelp beds grow rapidly in spring and summer,
3374 followed by a winter period when storms dislodge much of the algae, leaving little or no
3375 surface canopy. The biomass of kelp beds can also vary ten-fold or more from year to year
3376 due to interannual variation in the combinations of physical and biological variables that
3377 affect their growth.

3378 Kelp beds are biologically rich habitats due to both the primary productivity of the kelp and
3379 the effect kelp beds have on the surrounding environment. Bull kelp is one of the fastest
3380 growing organisms in the world, annually providing a large biomass available for
3381 consumption directly or as detritus after the kelp dies. Kelp furnishes a vertical habitat
3382 structure that otherwise would not exist on the reef. Kelp beds also slow water currents
3383 and reduce waves and wind chop, helping to trap drifting larva and nutrients and providing
3384 shelter. Kelp beds and their canopies can also support a rich understory of algal and
3385 attached invertebrate cover. On Oregon reefs, dense understory algae coverage gives way
3386 to dominant invertebrate cover at about 5 to 10 m water depth. Thick kelp cover reduces
3387 light penetration and can limit the density of understory algae. The kelp bed and underlying
3388 reef support a diverse array of fish and invertebrate species and provide cover and foraging
3389 areas for diving seabirds and marine mammals. In Oregon, the mix of fish species on kelp
3390 bed and non-kelp bed reefs is similar, reflecting a lack of kelp-specialist fish species that is
3391 perhaps unsurprising given the relatively low proportion of Oregon's rocky seafloor that is
3392 covered in kelp canopy.

3393 *Human Use*


3394 Human uses of nearshore rocky reefs include fishing, scientific research, sightseeing, and
3395 a number of other recreational and industrial pursuits. Commercial and recreational
3396 fishing for many types of rockfish species, lingcod, cabezon, and kelp greenling are the
3397 primary human uses of this habitat to date. SCUBA diving and underwater photography are
3398 among the other less prevalent uses. Much of the commercial live fish fishery takes place
3399 on shallow nearshore reefs. Recreational anglers also favor shallow nearshore reefs, if they
3400 are available. Commercial fishing effort targeting nearshore species tend to be higher on
3401 the south coast and recreational effort more prevalent on the north coast. Many reefs are
3402 used recreationally by SCUBA divers, sea kayakers, boaters, and surfers. Reefs with
3403 extensive kelp beds and islands provide sightseeing and bird watching opportunities for
3404 coastal residents and visitors. However, many reefs have no features extending to the
3405 ocean surface, and thus many people are unaware of the teeming life existing just below
3406 the water's surface.



3407

3408 **Figure 1.** Maximum historical extent of kelp beds along the north Oregon coast.

3409



3410

3411 **Figure 2.** Maximum historical extent of kelp beds along the south Oregon coast.

3412 **Rocky Shore**

3413 Oregon’s rocky shores, often referred to as rocky intertidal or tidepool areas, form parts of
 3414 the shoreward boundary of the nearshore planning area and can extend from the extreme

3415 low tide to the extreme high tide. They are characterized by CMECS as marine nearshore
3416 areas in the Intertidal and Supratidal zones, which include all hard substrate areas along
3417 the shoreline that are alternately exposed and covered by tides or are affected by wave
3418 splash and overwash, but not areas affected only by wind-driven spray. Everything beyond
3419 the reach of ocean waves is considered terrestrial habitat. The substrates making up
3420 Oregon's rocky shores include both volcanic and sedimentary bedrock as well as
3421 megaclasts, boulder, cobble and human-made (anthropogenic) structures. Some rocky
3422 shore areas are extensions of other shoreline rocky features such as headlands or cliffs,
3423 others exist as isolated regions of rock surrounded by sandy beach habitat, and some are
3424 anthropogenic in origin, having been deposited intentionally or unintentionally by humans.
3425 Oregon's coastline has approximately 152 linear miles of rocky shore habitat, and some 20
3426 miles of jetties.

3427 An example of a naturally occurring geform component found in Oregon's rocky shores
3428 would be a tidepool. Some of the anthropogenic geforms found in Oregon's rocky shores
3429 include breakwaters, jetties, and rip rap deposits. All rocky shore habitats in Oregon are
3430 contained entirely within the SWAP's planning area.

3431 *Physical Environment*

3432 The physical characteristics of rocky shores reflect local shoreline geology, exposure to
3433 ocean waves and currents, and biological influences. The Pacific Ocean exerts tremendous
3434 energy on Oregon's rocky shoreline, eroding coves, widening crevices, and reducing
3435 bedrock to rubble. On the north and central coast volcanic basalt dominates the hard
3436 shoreline, but sedimentary sandstone and mudstone rock can be found at several
3437 locations. Between Coos Bay and the Coquille River the geology is characterized by
3438 sedimentary rock. South of the Coquille River, headlands and rocks are primarily remnants
3439 of ancient metamorphic rocks over 200 million years old. Because of the variety of geologic
3440 origins and processes, Oregon's rocky shores consist of an assortment of cliff faces, wave-
3441 cut platforms, boulder fields, outcrops, and rubble. Each geform presents a unique
3442 mixture of habitats that provide shelter and substrate to support a wide variety of life.

3443 Ocean forces and weather strongly influence rocky intertidal environments. Tides are the
3444 primary influence on organisms and communities. The physical environment of intertidal
3445 areas changes dramatically as the tide rises and falls, alternately covering everything with
3446 salt water or exposing it to air, fresh water from rain and runoff, and the sun. Wave exposure
3447 also has a primary influence on this environment. Intertidal areas protected from waves
3448 due to shoreline orientation or geology provide dramatically different habitat than areas
3449 directly exposed to wave action. Local alongshore currents and ocean circulation
3450 processes introduce additional variables in the habitat, including sand scour of rocks,
3451 seasonal sand burial of rocky areas, and transport of nutrients, larvae, and adult organisms
3452 to and from intertidal sites.

3453 *Biological Characteristics*

3454 Rocky shore habitats are known for and crucial to their abundant and diverse biological
3455 communities. The variety in tidal elevations, wave exposure, and geologic structure within
3456 and among intertidal habitats produces a variety of microhabitats, often within relatively
3457 small geographic areas. This, in turn, provides for a diversity of species adapted to life in
3458 these different microhabitats. Organisms contribute to the variety of habitats as well. For
3459 instance, mussels and algae attach to the rocks, sometimes in huge numbers, providing
3460 additional structure and biogenic habitat used by intertidal species. Anthropogenic
3461 geoforms like jetties often take on similar biological characteristics of natural rocky shore
3462 geoforms, with similar biological communities using them.

3463 Biological communities associated with rocky intertidal habitat include algae, marine
3464 plants, attached and mobile invertebrates, fish, marine mammals and birds (see **Appendix**
3465 **- Nearshore Species**). Algae cover many intertidal areas with dense growth, often layered
3466 with several different species. Surfgrass, a marine vascular plant, often forms thick beds in
3467 lower intertidal areas, providing additional habitat structure for invertebrates and fish. Most
3468 rocky shore areas are extensively covered with attached invertebrates. Common types of
3469 attached organisms include sponges, anemones, barnacles, bryozoans, tunicates, and
3470 mussels. The rocks, algae, and attached invertebrates provide homes for a variety of
3471 mobile invertebrates such as crabs, snails, limpets, sea stars, urchins, brittle stars,
3472 nudibranchs, chitons, and worms. Free-swimming invertebrates, such as shrimps and
3473 drifting (planktonic) invertebrates also occur in tidepools or drift in with the tides. The algal
3474 and invertebrate communities in rocky intertidal areas often form distinct horizontal bands
3475 or zones of life according to the amount of time exposed to the air or covered by the tides.

3476 The upper reaches of the supratidal and intertidal zones experience the greatest variation
3477 in moisture, exposure, and salinity, and are often highly dependent on strong wave action to
3478 bring in nutrients and life. Compared to other shore areas, fewer species are found in
3479 the high intertidal and supratidal. These zones are typically characterized by vegetated
3480 rocks and boulders, along with isolated crevices and tidepools that hold water even during
3481 low tides. Greater abundance and diversity of life is associated with the lower intertidal
3482 areas. The distribution of organisms living in the mid-intertidal is generally limited at upper
3483 elevations by environmental stressors (such as high temperatures and desiccation) and at
3484 lower elevations by biological interactions (such as predation and competition). Organisms
3485 in the lowest parts of the rocky shore area experience almost continual tidal inundation
3486 and must be able to withstand the mechanical and biological stresses associated with this
3487 high-energy environment.

3488 The low intertidal serves as an important connection in the marine food web. Wave activity
3489 helps convert kelp and other organic debris into small fragments that are consumed by
3490 grazers and filter feeders and provide some nutrients to algal communities. Invertebrates
3491 and small fish provide a source of food for numerous bird species that forage along rocky
3492 shores.

3493 Fishes using the rocky shore include species adapted to live in tidepools and subtidal
3494 species that move in and out of the intertidal area with the tides. Tidepool fishes include a
3495 variety of sculpins, gunnels, and pricklebacks, among others. Rockfish species, greenlings,
3496 and surfperch often move into the intertidal area during high tide to feed and take refuge
3497 from subtidal predators. The rocky shore area is especially important to juvenile life stages
3498 of these fishes. The rocks and islands associated with Oregon's rocky shores and the
3499 subtidal rocky reefs provide important seal and sea lion haul out and pupping areas, and
3500 support some of the largest seabird nesting colonies on the contiguous U.S. West Coast.
3501 Islands and rocky intertidal areas are also utilized for nesting by birds that do not nest in
3502 colonies like Black Oystercatchers. Islands are another example of geofoms in the CMECS
3503 framework. Several seabird species that do not nest in colonies in Oregon do feed and take
3504 refuge here, including Black Turnstones and surfbirds.

3505 Rocky shores are linked to surrounding habitats by ocean currents and organism
3506 movements. Currents bring in planktonic organisms that help feed intertidal animals, and
3507 transport drifting larvae to and from intertidal environments. Currents also bring nutrients
3508 that feed the lush algal growth. Many organisms move in and out of intertidal habitats to
3509 feed or take refuge. Fish move in during high tides and terrestrial animals move in during
3510 low tides. Rocky intertidal areas are also linked to each other, primarily through transport of
3511 larvae by ocean currents. The proximity of intertidal habitat to other "upstream" or
3512 "downstream" habitats has dramatic effects on the types, abundance, and recruitment
3513 rates of communities and organisms.

3514 Ecological linkages within and between rocky shore areas help to shape biological
3515 communities and contribute toward the biological abundance of this habitat type. The
3516 diversity of producers and consumers in the intertidal create complex food webs and
3517 interdependencies among organisms. This complexity of organism interrelationships
3518 makes the outcome of natural or human disturbance to rocky shore habitats difficult to
3519 predict or measure. For instance, while human foot traffic can result in inadvertent
3520 trampling of organisms, anthropogenic structures such as jetties provide a unique and
3521 valuable rocky shore habitat at the transition between estuaries and the marine
3522 environment.

3523 *Human Use*

3524 Human uses of rocky intertidal areas include fishing, invertebrate and algae harvest and
3525 collection, education, scientific research, sightseeing, and other recreational, economic,
3526 and social pursuits. Due to their accessibility and the fascinating array of marine life, rocky
3527 intertidal areas receive more public use than many other marine habitats. Visitation by
3528 school groups and others curious about marine life comprises the majority of public use.
3529 For many visitors, their first and sometimes only interaction with the wonders of marine life
3530 comes from tidepool visits. Visitation of rocky shore areas has generally been increasing
3531 over the past five decades.

3532 Rocky shores are used extensively by researchers as a natural laboratory to increase
3533 understanding about general marine ecological principles. In 2025, there are fifteen
3534 intertidal and subtidal sites along the Oregon coast that have special regulations limiting
3535 harvest or collection of organisms in order to enhance scientific research, as well as
3536 education and enjoyment benefits.

3537 Detailed descriptions of types and amount of human use at individual rocky shore sites
3538 along Oregon's coast can be found in the *Oregon Rocky Shores Natural Resources*
3539 *Inventory* (ODFW 1994).

3540 **Sandy Beaches**

3541 Sandy beaches are a widespread feature of the entire Oregon coast and make up
3542 approximately two-thirds of the coastline. Their distribution is interrupted by rocky shores,
3543 rocky headlands, river mouths, estuaries, and human constructions. Oregon's sandy
3544 beaches are characterized by CMECS as marine nearshore areas in the intertidal and
3545 supratidal zones that are composed of very fine to very coarse sand substrate; they extend
3546 in a continuum from the Mean Lower-Low Waterline to the areas above the Mean Higher-
3547 High Waterline that are affected by wave splash and overwash at extreme high tides, but
3548 not areas affected only by wind-driven spray. Sandy beaches stretch inland until they are
3549 stopped by a continuous line of vegetation, debris, rocks, or other barrier. Everything
3550 beyond the reach of the waves and splash zone is considered terrestrial habitat.

3551 *Physical Environment*

3552 Oregon's sandy beaches are high-energy environments that experience significant wave
3553 and wind energy. Several million cubic meters of sand are transported to the nearshore
3554 area annually by river systems. Seasonal variation in wind and wave energy and currents
3555 move substantial amounts of sand onto or off beaches, which results in significant
3556 changes in beach character as underlying rock structures (bedrock and/or cobble) are
3557 exposed. In some areas, patches of ancient forest where the land dropped during past
3558 subduction zone earthquakes may become exposed. Currents and wave energy are other
3559 significant factors in moving sands onto or off of beaches at elevations that are frequently
3560 immersed; the lateral width of the beach will govern the area over which current and wave
3561 energy is dispersed and hence determines the slope of the beach as sands are deposited
3562 or swept away. At higher elevations that are dry and experience infrequent immersion by
3563 tides, wind is the predominant factor in distributing sand and can create windows and
3564 mobile dunes from a few centimeters to several meters tall, while dunes further inland may
3565 be several stories high.

3566 The lateral (north-south) extent of sandy beaches is punctuated by rivers or rocky
3567 headlands where the transition from sand to volcanic rock can be quite abrupt. Rivers can
3568 frequently become "bar-bound" during the summer and early fall months when river flows
3569 diminish due to reduced precipitation, and the energy of flowing water is insufficient to

3570 maintain an open, flowing channel to the sea. In such cases, the river or stream will flow
3571 *through* the sand in its final stages. Bar-bound rivers are generally freed by fall rains on the
3572 Oregon coast that increase river flows and wash sand out of the river mouths to re-
3573 establish a channel of flow. Fall rains and the breaking of blocking bars are important in
3574 restoring access to fresh-water streams for anadromous fishes.

3575 The supratidal zone and upper range of the intertidal zone are subject to the greatest
3576 variation in temperature and moisture and the least physical energy from the ocean. The
3577 intertidal zone, particularly its lower reaches, receives much greater physical energy from
3578 waves and currents, and experiences the least variation in temperature.

3579 *Biological Characteristics*

3580 The movement of sand by water and wind energy makes sandy beaches largely unsuitable
3581 for rooted and attached organisms. However, between the grains of sand in the intertidal
3582 zone is a vast multitude of life too small to see with the naked eye, including diatoms,
3583 harpacticoid copepods, amphipods, and algae, among others. Larger invertebrates can be
3584 found here as well, including crustaceans, mollusks, and diverse worm taxa. Many of the
3585 resident invertebrates burrow in the sand during periods of exposure for protection from
3586 desiccation and/or predation and emerge to forage as tides permit.

3587 Biological communities of the upper intertidal and supratidal zones of sandy beaches are
3588 often based on the resources provided by the incoming tides and deposited at the high tide
3589 line. Once in the intertidal zone, the detritus is broken down by the mechanical force of
3590 waves pounding against the shore and the industry of the many organisms that live and
3591 forage there. Organisms of the mid and lower intertidal, particularly the small invertebrates,
3592 provide food resources for numerous larger invertebrates, fish, and bird species. Some
3593 marine mammals intentionally use this zone to rest, hauling themselves out of the ocean to
3594 lay on the sand.

3595 Several SGCN, Watch List, and commonly associated species are also connected with
3596 general sandy beach habitats, or specific to distinct sandy beach types (see **Appendix -**
3597 **Nearshore Species**). Surf smelt use particular beaches to lay their eggs in the intertidal
3598 zone. Native littleneck and razor clams burrow below the sand and feed on plankton when
3599 the ocean water covers them. Western Snowy Plover nest either in the supratidal zone or
3600 above and feed in the intertidal sandy areas. Sanderlings gather in loose flocks in the winter
3601 months to feed on the rich array of invertebrates under the sand as the waves recede.
3602 Harbor seals rest on sandy beaches and northern elephant seals come ashore to molt,
3603 usually in the supratidal zone.

3604 *Human Use*

3605 Sandy beaches attract substantial human use at all levels of the intertidal and supratidal.
3606 Their easy access and wide variety of organisms and ecological processes attract scientific

3607 interest. Thanks to their uniform, comfortable surface, sandy beaches are valued for a wide
3608 variety of recreational activities including sightseeing, picnicking, walking, running, agate-
3609 hunting, and kite flying. Lower portions of beaches are also launch and recovery areas for
3610 surfers, windsurfers, kite boarders, sea kayakers, and some sailboats, power boats, and
3611 personal watercraft. Wildlife found at sandy beaches is highly valued by humans for
3612 everything from bait or dinner to instructional or aesthetic uses. Driving is permitted on
3613 some Oregon beaches, but not all. All beaches in Oregon are free for the public to access.

3614 CONSERVATION OVERVIEW

3615 There are many Key Conservation Issues (KCI) affecting Nearshore habitats and the
3616 species that live there, including **Climate Change**, **Disruption and Disturbance Regimes**,
3617 **Land Use Changes**, **Invasive Species**, **Pollution**, and **Water Quality and Quantity**.

3618 LIMITING FACTORS AND RECOMMENDED APPROACHES

3619 **Limiting Factor: Public Awareness**

3620 Oregon's nearshore waters are part of the California Current Ecosystem. What occurs in
3621 the ocean waters of the Pacific, such as El Niño and La Niña, and Pacific Decadal
3622 Oscillation, and the timing of spring and fall transition can greatly influence not only the
3623 nearshore habitats and species, but also such things as rainfall, snowpack and drought
3624 conditions throughout the state. A well-informed public helps drive policy and
3625 management decisions as well as individual actions that support a healthy ecosystem and
3626 the many benefits it offers.

3627 **Recommended Approach**

3628 There are a series of recommended approaches in the **Nearshore ecoregion**
3629 Recommendations under the category of education and outreach that include: 1)
3630 developing creative ways to engage with the general public, constituent, and advisory
3631 groups and exploring technologies that support alternative methods of communication and
3632 participation in addition to traditional paths such as issue-specific advisory groups; 2)
3633 broadening outreach materials and information available electronically to deepen public
3634 appreciation of Oregon's nearshore environment; and 3) developing new and expanding
3635 existing partnerships for communication, education, and outreach on nearshore topics
3636 and issues. This approach depends on having the necessary research and monitoring to
3637 provide the public with information about the issues listed above and how those issues
3638 translate into direct threats to fish, wildlife and their habitats in the nearshore.

3639 **Limiting Factor: Climate Change and Disruption and Disturbance Regimes**

3640 Oregon's ocean is already experiencing effects from climate change and increased carbon
3641 dioxide, including ocean acidification, hypoxia, other changes in water chemistry, warming
3642 ocean temperature, and changes in upwelling and other characteristics of the nearshore
3643 ocean and estuaries. These changes will continue to grow and intensify in the future.
3644 Oregon's upwelling ecosystem is experiencing many of these changes sooner and in
3645 greater magnitude than other parts of the nation, increasing the urgency for collecting the
3646 needed information and formulating the necessary management response. This is a global
3647 problem that requires rigorous scientific information to solve, and partnership between
3648 scientists inside and outside of agencies to both understand the phenomena and try to
3649 mitigate its effects. Desired outcomes are to increase ecosystem and community
3650 resilience and sustainability of Oregon's nearshore resources.

3651 **Recommended Approach**

3652 Expanding research and monitoring activities are required to generate the data and
3653 information needed. This is especially true in the areas where human activities are intense
3654 and information on species and their habitats is sparse. Develop and implement research
3655 and monitoring efforts to understand, track, and work toward predicting effects of climate
3656 change and increased carbon dioxide on Oregon's nearshore species and ecosystems.
3657 Focus research on species and ecosystems most at risk, and foster collaboration between
3658 scientists and managers to optimize research outcomes for use in management and
3659 conservation. Continue and expand research and monitoring efforts on nearshore species
3660 and habitats. Gather scientific information on the abundance and distribution of species
3661 and habitats, the interactions among species and between species and their physical
3662 environment, and changes in those resources and interactions over time. Priorities for
3663 research and monitoring needs include oceanographic data, ecosystem data, habitat data,
3664 human dimensions and the impacts of human development (see **Appendix - Nearshore
3665 Research and Monitoring**). Promote use of climate change information in management
3666 decision-making and policy development in statewide, regional and global arenas. Build
3667 climate resilience and climate change adaptation into decision-making to maximize the
3668 long-term benefits of today's public investment in natural resource management.

3669 **Limiting Factor: Land Use Changes (Marine Spatial Planning)**

3670 The **Land Use Changes** KCI provides an overview of the issues associated with land use
3671 throughout the state and information about **Oregon's 19 Statewide Land Use Planning
3672 Goals**. Goals 16-19 are particularly relevant to the Nearshore environment and ecosystem,
3673 especially Goals 16 and 19 on estuarine and ocean resources, respectively. In the wider
3674 marine realm "land use" is often referred to as "marine spatial planning". Oregon utilizes its

3675 **Territorial Sea Plan** to guide state agency actions. Changes to land use in coastal areas
3676 directly and indirectly affect nearshore species and habitats in a variety of ways, that
3677 include such things as shoreline armoring, leasing mariculture plots, siting renewable
3678 energy developments such wave or wind energy facilities, as well as designating marine
3679 reserves, marine gardens, and research areas. In addition to state agencies, federal
3680 agencies also have various roles in marine spatial planning such as USWF, USCG, and
3681 BOEM. There is growing demand for ocean and coastal resources, and competing use of
3682 space has increased the need to move beyond single-sector management and plan for
3683 ocean uses more holistically. Marine planning processes require comprehensive spatial
3684 information on location, abundance and distribution of marine resources and their uses.

3685 **Recommended Approach**

3686 Participate in marine planning processes to ensure Oregon’s interests in marine natural
3687 resource conservation and use are fully represented in marine policy. Develop marine
3688 natural resource spatial information and incorporate it into marine planning processes to
3689 ensure they use the best available science to formulate plans concerning Oregon’s marine
3690 resources and uses. This will require partnerships with State and federal natural resource
3691 agencies, sport and commercial fishing interests, local, state, regional, and federal
3692 governments, community groups, non-governmental organizations, tribes, and the general
3693 public.

3694 **Limiting Factor: Pollution and Water Quality**

3695 Pollution in all its various forms can directly impact nearshore species and their habitats.
3696 **Water quality** is affected not only by pollution of the nearshore environment, but also by
3697 climate change effects that cause ocean warming, ocean acidification and hypoxia, all of
3698 which impact nearshore species and habitats.

3699 **Recommended Approach**

3700 Determining the vulnerability of species and habitats to various types of pollution requires
3701 research and monitoring in the nearshore. Similarly, the goals of monitoring water quality
3702 also depend on research and monitoring efforts in the nearshore. Expanding existing
3703 research and monitoring efforts on these topics will enhance our understanding of their
3704 effects, help inform the public, and drive management and policy choices to help achieve
3705 these goals.

3706 **Limiting Factor: Non-native and Invasive Species**

3707 Many non-native and invasive species have made their way to Oregon’s nearshore waters
3708 or to those of our neighboring states (see **Appendix - Nearshore Species**). These have
3709 been introduced through a variety of mechanisms that include hitch-hiking in ballast water
3710 or in ocean currents. These species can affect food sources, alter habitats, expose native
3711 communities to diseases or toxins, or act as parasites of juvenile and adult members of
3712 coastal species. For many species, the severity of the potential ecological threat is not yet
3713 known. Many of these species could be deemed invasive in the future, but further efforts to
3714 assess impacts are needed.

3715 **Recommended Approach**

3716 Achieving the goals to meet the challenges non-native and invasive species pose take a
3717 collaborative effort. This work need will include education and outreach, research and
3718 monitoring, and policy and management to be successful.

3719 **RESOURCES FOR MORE INFORMATION**

3720 Appendix - Nearshore Climate Change Fact Sheets

3721 Appendix - Nearshore References

3722

3723 SPECIALIZED AND LOCAL HABITATS

3724 Some natural communities and landscape features are not adequately represented
3725 through **Key Habitats**. These communities and features often occur at the local scale and
3726 have a patchy distribution across the landscape. They may be difficult to map from
3727 satellite data and may not be represented well in available datasets. Some of these
3728 habitats provide functions and values that are highly specialized to the local environment,
3729 are limited in quantity, and host a suite of rare or endemic species. To address the
3730 conservation needs of these habitats and their associated species, “specialized and local
3731 habitats” were identified through review of geographic vegetation data, rare plant or animal
3732 occurrences, importance to **Species of Greatest Conservation Need**, and occurrences of
3733 animal concentrations. Many of these habitats are also identified in other state priorities,
3734 such as **Aquatic Resources of Special Concern** through the Oregon Department of State
3735 Lands.

3736 ALPINE HABITATS: MEADOWS, DWARF SHRUBLANDS, ALPINE 3737 TUNDRA, AND WHITEBARK PINE

3738 **Ecoregions**

3739 **BM, EC, KM, NBR, WC**

3740 Alpine habitats provide important foraging and breeding areas for many mammals and
3741 birds as well as critical resources for birds during migration periods. These habitats are at
3742 risk from increased recreational activity and are extremely vulnerable to climate change as
3743 temperatures warm and snowpack decreases. Alpine habitats provide important resources
3744 for many at-risk species, such as the federally listed (threatened) whitebark pine (*Pinus*
3745 *albicaulis*) that is particularly vulnerable to white pine blister rust, outbreaks of mountain
3746 pine beetles, and fire suppression resulting in replacement by more shade-tolerant tree
3747 species.

3748 **Conservation Actions**

3749 Mitigate effects of climate change and provide refugia for fish and wildlife. Manage
3750 recreation, human disturbance, and grazing to minimize impacts to soil and plant
3751 communities. Monitor and control invasive plants. Re-introduce fire into the ecosystem to
3752 prevent fuel build-up and canopy closure as feasible and appropriate to the local area.
3753 Identify blister rust resistant whitebark pine trees and collect the seeds for nursery stock.

3754 AQUATIC VEGETATION BEDS

3755 **Ecoregions**

3756 **All**

3757 Aquatic vegetation beds are a component freshwater ponds, riverine sloughs and alcoves,
3758 and estuaries and nearshore waters. They are at the base of the food chain and provide
3759 habitat for a host of organisms from bacteria, protozoa, and invertebrates to fish,
3760 amphibians, reptiles, birds, and mammals. Vital to maintaining the ecological integrity of
3761 aquatic ecosystems, their preservation and restoration are essential for supporting
3762 biodiversity, improving water quality, and providing numerous ecosystem services that
3763 benefit both wildlife and human communities.

3764 **Conservation Actions**

3765 Retain and restore natural water flow regimes. Maintain consistent water levels. Mitigate
3766 impacts from climate change. Monitor for and control invasive plants such as reed canary
3767 grass and Ludwigia. Mitigate effects of runoff from agricultural fields and roadways. Limit or
3768 prohibit dredging in estuarine algal beds.

3769 **ASH FLOWS, ASH BEDS AND LAVA FIELDS**

3770 **Ecoregions**

3771 **BM, EC, NBR**

3772 Ash flows, ash beds, and lava fields provide habitat for many rare and endemic and other
3773 specialized plants and invertebrates, such as the Oregon lava hole bee (*Atoposmia*
3774 *oregona*). These sites can also be important fossil localities.

3775 **Conservation Actions**

3776 Manage grazing, mining, and off-highway vehicles to minimize erosion and disturbance to
3777 rare plants and invertebrates.

3778 **BALDS AND BLUFFS**

3779 **Ecoregions**

3780 **BM, CR, EC, KM, WC, WV**

3781 Balds and bluffs provide habitat for unique plant communities and invertebrates such as
3782 butterflies. In the Coast Range ecoregion, these habitats include coastal bluffs and
3783 headlands. In the Klamath Mountains ecoregion, these habitats include serpentine barrens

3784 and outcrops. In the Willamette Valley ecoregion, these habitats include wet rock outcrops
3785 dominated by camas (*Camassia spp.*) and other wet prairie species.

3786 **Conservation Actions**

3787 Better mapping and documentation of balds and bluffs are needed. Control encroaching
3788 conifers and shrubs. Monitor for and control invasive plants. Minimize disturbance (e.g.,
3789 trail or road construction, recreation) to help protect rare plant communities. Protect
3790 hydrology to maintain perched wetland and wet rock outcrop function. Consider impacts
3791 from changing fire regimes.

3792 **BAYS**

3793 **Ecoregions**

3794 **CR, NS**

3795 Bays provide winter habitat for waterfowl and other waterbirds, rearing areas for juvenile
3796 anadromous salmonids, and habitat for intertidal and subtidal shellfish beds, including
3797 native oyster beds.

3798 **Conservation Actions**

3799 Provide areas of low disturbance during critical life history needs and time periods.
3800 Minimize impacts from in-water activities such as dredging, as well as impacts from
3801 overwater structures. Coordinate with landowners, communities, local governments,
3802 development interests and other partners to properly plan development to avoid, minimize
3803 and mitigate impacts to bay ecosystems.

3804 **BITTERBRUSH COMMUNITIES**

3805 **Ecoregions**

3806 **BM, EC, KM, NBR, WC**

3807 Antelope bitterbrush (*Purshia tridentata*) is an important habitat component that provides
3808 forage, cover, and nesting habitat for a variety of wildlife. It provides high value winter
3809 forage for deer, elk and pronghorn, supports a variety of insect pollinators, and provides
3810 seeds that support a diversity of small mammals. In some areas, juniper encroachment
3811 threatens bitterbrush communities by outcompeting and shading bitterbrush.

3812 **Conservation Actions**

3813 Improve understanding of bitterbrush regeneration methods. Continue restoration and
3814 monitoring efforts. Manage grazing pressure based on site conditions. Bitterbrush can be
3815 impacted by prescribed fire; caution is needed if considering this tool in proximity.

3816 CANYON SHRUBLANDS

3817 **Ecoregions**

3818 **BM, CP, EC, NBR**

3819 Also known as moist deciduous shrublands, canyon shrublands provide nesting habitat for
3820 songbirds and winter habitat for SGCN such as Columbian Sharp-tailed Grouse
3821 (*Tympanuchus phasianellus columbianus*).

3822 **Conservation Actions**

3823 Maintain healthy shrubs stands and restore degraded stands. Some degraded stands can
3824 benefit from prescribed fire, removal of encroaching invasive junipers, or management of
3825 grazing season timing.

3826 CAVES AND OLD MINES

3827 **Ecoregions**

3828 **BM, CR, EC, KM, NBR, WC**

3829 Caves and old mines provide habitat for rare invertebrates and cave-roosting bats, such as
3830 Townsend's big-eared bat (*Corynorhinus townsendii*) and several *Myotis* species. In the
3831 East Cascades ecoregion, these habitats include lava tubes.

3832 **Conservation Actions**

3833 Use gates or seasonal closures to protect known roost sites from recreational caving and
3834 other disturbance. When mines are closed for human safety, provide openings for bat entry
3835 and exit. Be aware of the potential for white-nosed syndrome; thoroughly sanitize all
3836 clothing, footwear, and equipment between caves to prevent potential contamination.

3837 CHAPARRAL AND CEANOTHUS SHRUBLAND

3838 **Ecoregions**

3839 **BM, CR, EC, KM, WC, WV**

3840 Chaparral and ceanothus shrublands provide cover, nesting, and foraging habitat for
3841 songbirds, kingsnakes, and a variety of invertebrates, including some butterfly species.
3842 These shrublands occur in open areas, so may be found in early successional habitats or
3843 at high elevations, where temperatures and other factors inhibit tree growth. In the
3844 Klamath Mountains ecoregion, chaparral is often removed as a fire hazard, as some
3845 species are highly flammable and dependent on fire for seed germination. Chaparral is
3846 also increasingly removed during development, particularly in lowland valleys. These
3847 habitats are at risk from fire suppression—many ceanothus species become senescent
3848 without the fires needed for regeneration. Chaparral is also unusual habitat in the
3849 Willamette Valley, which makes protecting existing sites important for maintaining local
3850 species diversity.

3851 **Conservation Actions**

3852 Maintain shrub diversity during forest management activities. Delay replanting with
3853 conifers where shrub habitat is limited. Control key invasive plants (e.g., Scotch broom and
3854 Armenian (Himalayan) blackberry) and animals such as feral horses at priority sites.
3855 Implement controlled burns or other fire management techniques where appropriate to
3856 the local area.

3857 **EELGRASS BEDS**

3858 **Ecoregions**

3859 **CR, NS**

3860 Eelgrass beds support the aquatic food chain and provide essential habitat for many
3861 species to fulfill their life history needs. They provide habitat to support intertidal and
3862 subtidal shellfish beds, including native oyster beds. They also provide important rearing
3863 habitat for juvenile fish, including commercially important species, and foraging habitat for
3864 birds, such as Brant (*Branta bernicla*).

3865 **Conservation Actions**

3866 Ensure that development activities that may disturb eelgrass beds avoid, minimize and
3867 mitigate direct and indirect impacts. Discourage dredging or fill of estuaries and eelgrass
3868 beds. Monitor and control invasive species. Restore and monitor eelgrass habitats.
3869 Research the role of eelgrass in mitigating the impacts from climate change. Protect
3870 genetic diversity within eelgrass populations. Also see **Estuaries**.

3871 **FEN PEATLANDS**

3872 **Ecoregions**

3873 **BM, CR, EC, WC**

3874 Fens are peat-accumulating wetlands that form where groundwater discharge is low but
3875 constant, and where appropriate geologic conditions occur, such as glacial deposits with
3876 pumice. Fens provide habitat for sensitive plant species and provide long-term carbon
3877 storage in the form of peat. They are highly sensitive to climate change, which may reverse
3878 the process of peat accretion and lead to carbon loss.

3879 **Conservation Actions**

3880 Maintain groundwater recharge areas, especially at higher elevations. Use conservation
3881 incentives, and where applicable, maintain existing protection standards to provide
3882 buffers around fen areas. Seek opportunities to enhance recharge from local aquifers
3883 supporting the fens.

3884 **FOREST OPENINGS**

3885 **Ecoregions**

3886 **BM, CR, EC, KM, WC, WV**

3887 Forest openings provide essential structural complexity and plant diversity within forests.
3888 Forest openings provide foraging habitat for a variety of species that are adapted to open
3889 meadows, early seral habitat, and forest edges. They support bird species like Olive-sided
3890 Flycatchers, Willow Flycatchers, and Common Nighthawks, as well as species that prefer
3891 open habitat with snags such as Purple Martin and Western Bluebird. Clouded
3892 salamanders live in large logs and stumps in openings, and their populations increase
3893 following wildfires. Disturbances such as wildfire, windthrow, disease, and insect
3894 outbreaks reset succession and often result in large or small openings with high forb and
3895 shrub diversity and woody structure (e.g., large snags and logs). Management of older
3896 successional forest stages on public land typically does not include maintaining forest
3897 openings, and private forestlands are usually intensively managed, often as monocultures,
3898 which leads to a rarity in forest openings with structural complexity and plant diversity.

3899 **Conservation Actions**

3900 During salvage logging or other timber harvest, minimize ground disturbance, and maintain
3901 and create snags and downed logs. Pursue forest management activities that create forest
3902 openings and maintain natural forb, grass, and shrub species. Control key invasive plants
3903 in openings. After burns, reseed with native grasses and forbs, and delay replanting with
3904 conifers. Carefully evaluate salvage logging in burned late successional forests. Provide
3905 education to the public about how not to spread invasive plant species and the importance
3906 of control and management.

3907 UNIQUE GRASSLAND HABITATS

3908 **Ecoregions**

3909 **EC, KM, NBR**

3910 Unique grassland habitats in Oregon include alkali grasslands, perennial bunchgrass, and
3911 montane grasslands. These habitats are important for raptors, grassland birds, and rare
3912 plants.

3913 **Conservation Actions**

3914 Maintain and restore these unique grasslands using site-appropriate tools. Monitor for
3915 invasive species. Manage grazing to minimize impacts to native species.

3916 GREASEWOOD FLATS AND WASHES

3917 **Ecoregions**

3918 **BM, CP, EC, NBR**

3919 Greasewood (*Sarcobatus vermiculatus*) is typically found in flats, washes, and terraces
3920 with saline soils and shallow water tables. Flats, washes, and terraces flood intermittently
3921 but remain dry for most of the growing season, providing habitat for rare plants.

3922 Greasewood is an important browse species for deer and pronghorn, as well as SGCN like
3923 white-tailed jackrabbit and North American porcupine. These habitats are threatened by
3924 changing fire regimes and the spread of invasive annual grasses.

3925 **Conservation Actions**

3926 Maintain and restore greasewood habitats. In the Blue Mountains, include black
3927 greasewood habitats when managing for a mosaic of valley bottom habitats.

3928 INLAND DUNES

3929 **Ecoregions**

3930 **CR, CP, NBR**

3931 Inland dunes include active and partially stabilized dunes in arid inland regions. These
3932 dunes provide habitat for a variety of species including reptiles, small mammals, and rare
3933 plants. In the Columbia Plateau ecoregion, stabilized dunes often support basin big
3934 sagebrush and bitterbrush. In the Northern Basin and Range ecoregion, the Christmas

3935 Valley Sand Dunes are the largest inland shifting sand dune system in the Pacific
3936 Northwest. The alkaline sands of the Northern Basin and Range ecoregion support salt
3937 desert dune shrubs such as greasewood and saltbush. Inland dunes along the Columbia
3938 River have stabilized in recent decades after the damming of the Columbia River.
3939 Historically these dunes were fed by sand transported and deposited annually by the river.
3940 Inland dunes are threatened by the spread of non-native species such as Russian thistle.

3941 **Conservation Actions**

3942 Maintain and enhance existing habitat. Monitor for and control invasive species. Protect
3943 dunes from uncontrolled off-highway vehicle use.

3944 **INTERDUNAL LAKES AND WETLANDS**

3945 **Ecoregions**

3946 **CR**

3947 These habitats are comprised of shallow lakes and wetlands located in areas between
3948 coastal sand dunes. Water levels in interdunal lakes and wetlands are dependent on local
3949 precipitation to recharge sand dune aquifers. These lakes and wetlands provide breeding
3950 habitat for SGCN, including northern red-legged frogs, and support unique wetland plant
3951 communities.

3952 **Conservation Actions**

3953 Maintain groundwater recharge areas at sand dune aquifers. Protect these habitats from
3954 off-road vehicle use and other impacts from human recreation and development.

3955 **INTERTIDAL MUDFLATS**

3956 **Ecoregions**

3957 **CR, NS**

3958 Intertidal mudflats provide foraging habitat for shorebirds, which is critically important
3959 during migration. Mudflats also serve as habitat for a diversity of invertebrate species such
3960 as clams and other shellfish.

3961 **Conservation Actions**

3962 Manage water flows to maintain mudflat habitats. Maintain or restore water quality and
3963 natural sedimentation patterns to preserve habitat quality for invertebrates.
3964 See **Estuaries**.

3965 KELP BEDS

3966 **Ecoregions**

3967 **NS**

3968 Limited to subtidal rocky areas in relatively shallow water, kelp beds are designated as
3969 essential fish habitat for both groundfish and salmon. These areas provide important
3970 habitat for a diversity of other species in the nearshore ecoregion as well.

3971 **Conservation Actions**

3972 Reduce coastal runoff that increases turbidity in nearshore ocean waters. Minimize risk of
3973 oil spills and pollution. Fill data gaps on the gametophyte stage of the kelp life cycle. See
3974 Nearshore Habitats.

3975 MOUNTAIN MAHOGANY WOODLAND AND SHRUBLAND

3976 **Ecoregions**

3977 **BM, EC, NBR**

3978 Mountain mahogany (*Cercocarpus spp.*) communities have expanded in some areas due
3979 to fire suppression but depend on low-intensity fire for long-term maintenance and
3980 regeneration. Many stands are threatened by non-native understory vegetation and juniper
3981 encroachment is a threat in some areas, especially in the Northern Basin and Range. In the
3982 East Cascades ecoregion, mountain mahogany is more diverse than in other ecoregions.
3983 Mountain mahogany in the East Cascades includes birchleaf mountain mahogany, which
3984 is found throughout moist shrublands in the southern portion of the ecoregion. Mountain
3985 mahogany serves as important nesting habitat for birds because it provides tree structure
3986 in otherwise open, shrub-dominated landscapes. Mountain mahogany also provides forage
3987 and cover for a diversity of mammal species.

3988 **Conservation Actions**

3989 Develop methods to manage mahogany stands and encourage regeneration. Restore
3990 native understory vegetation at priority sites. Conduct conifer management within and
3991 adjacent to stands, particularly western juniper management.

3992 OFF-CHANNEL HABITAT

3993 **Ecoregions**

3994 **All inland ecoregions**

3995 Off-channel areas provide critical rearing, security, and foraging habitat for juvenile
3996 salmonids and other native fish, northwestern pond turtles, freshwater mussels, and other
3997 invertebrates.

3998 **Conservation Actions**

3999 Protect and restore off-channel habitat, including restoration of stream hydrology. Avoid,
4000 minimize, and mitigate impacts to off-channel and riparian habitat from development
4001 actions. Manage beaver populations to provide for beaver-modified habitats, while
4002 minimizing conflicts with other land uses. Restoring tidal and riverine inundation to these
4003 areas and restoring/enhancing connectivity are key conservation actions. See **Flowing**
4004 **Water and Riparian Habitat**.

4005 PORT ORFORD CEDAR FORESTS

4006 **Ecoregions**

4007 **KM, CR**

4008 Endemic to southwestern Oregon and northwestern California, Port Orford cedar
4009 (*Chamaecyparis lawsoniana*) forests are associated with serpentine soils and are
4010 characterized by unusual plant and animal associations, co-occurring with SGCN such as
4011 large-flowered rush lily. These habitats have been severely impacted by an introduced,
4012 fungus-like tree disease, the Port Orford cedar root disease, particularly near the coast.

4013 **Conservation Actions**

4014 Maintain and protect existing habitat. Minimize vehicular traffic and/or new road
4015 construction where potential exists to spread the invasive root pathogen.

4016 ROCK HABITATS: CLIFFS, RIMROCK, ROCK OUTCROPS, AND TALUS

4017 **Ecoregions**

4018 **BM, CR, CP, EC, KM, NBR, WC, WV**

4019 Rocky areas provide habitat for peregrine falcons and other cliff-nesting birds, cliff-
4020 roosting bats, rare plants, and wildlife that use rocks for shelter and/or foraging areas.
4021 Talus slopes provide habitat for Larch Mountain salamander, pika, and several
4022 invertebrates. In the Willamette Valley, rock outcrops serve as hibernacula for snakes,
4023 including western rattlesnakes. In dry ecoregions, rock habitats are particularly important
4024 for salamanders as a refuge from hot, dry weather.

4025 **Conservation Actions**

4026 These habitats have few limiting factors in most ecoregions. In the East Cascades,
4027 residential development at the edge of rims alters vegetation and disturbs nesting birds.
4028 Work with local planners to implement existing setback distance standards through the
4029 Statewide Planning Program. Rock mining should be avoided in talus areas where known
4030 populations of Larch Mountain salamander and rare invertebrates occur. For all
4031 ecoregions, if important roosts, hibernacula, or nest sites are known, minimize
4032 disturbance.

4033 **ROCKY SHORES, TIDEPOOLS, AND OFFSHORE ROCKS (E.G., SEA**
4034 **STACKS)**

4035 **Ecoregions**

4036 **CR, NS**

4037 Rocky shores and offshore rocks provide critical nesting, roosting, and foraging habitat for
4038 seabirds and shorebirds, including SGCN like Tufted Puffin and Black Oystercatcher.
4039 These areas also serve as haul-outs for marine mammals, and as roosting areas for
4040 raptors, including peregrine falcons. Rocky shores, tidepools, and offshore rocks also
4041 provide habitat for a variety of marine invertebrates and fish.

4042 **Conservation Actions**

4043 Work with local communities and land management agencies to avoid and minimize
4044 impacts from tidepool viewing, and to minimize disturbance to birds and marine mammals
4045 during sensitive nesting and pupping seasons. Increase research to better understand the
4046 impacts of thermal heatwaves and other climate-related stressors. See Nearshore
4047 Habitats.

4048 **SALT DESERT SCRUB**

4049 **Ecoregions**

4050 **NBR**

4051 This low-to-medium shrub habitat can be found on dry sites with saline soils, such as dry
4052 lake beds, flat desert pavements, low alkaline dunes, around playas, or on gentle slopes
4053 above playas. Salt desert scrub provides habitat for a diversity of reptiles and mammal
4054 species, including species that are primarily or exclusively associated with this habitat,
4055 such as kit fox and long-nosed leopard lizard.

4056 **Conservation Actions**

4057 Salt desert scrub is threatened by invasion of non-native annual grasses, particularly
4058 cheatgrass. Microbiotic soil crusts are particularly critical in these habitats, so it is
4059 important to minimize activities that cause soil disturbance, such as hiking, biking, and off-
4060 highway vehicle use.

4061 **SAND SPITS, SAND BARS, AND SPARSELY VEGETATED ISLANDS**

4062 **Ecoregions**

4063 **CR, EC, NBR, NS**

4064 Sparsely vegetated sandy habitats that are isolated from disturbance due to humans and
4065 mammalian predators are important roosting and nesting sites for colonial waterbirds,
4066 such as American White Pelicans, Brown Pelicans, and Caspian Terns. In eastern Oregon,
4067 this habitat occurs around large lakes and wetlands. Sparsely vegetated island habitat can
4068 be surrounded by either saltwater or freshwater.

4069 **Conservation Actions**

4070 Maintain open habitat characteristics and minimize disturbance at key sites. Manage
4071 water levels to preserve island habitats.

4072 **SPRINGS, SEEPS, AND HEADWATERS**

4073 **Ecoregions**

4074 **All inland ecoregions**

4075 Springs, seeps, and headwaters provide habitat for amphibians, invertebrates, and rare
4076 plants. The isolated nature of springs is one of the factors resulting in high levels of
4077 invertebrate endemism in the East Cascades. Spring systems in the Northern Basin and
4078 Range also contain endemic species, including vertebrates (e.g., Hutton tui chub and
4079 Foscett speckled dace). In dry ecoregions, spring and seep habitats are important as a
4080 source of water for wildlife and as habitat for amphibians and invertebrates. These habitats
4081 have been impacted by livestock watering and agricultural uses. Springs, seeps, and

4082 headwaters are critical to protect for climate resiliency, particularly for water quantity and
4083 quality, and are a refuge for multiple species during and following wildfire.

4084 **Conservation Actions**

4085 Encourage use of incentives, and where applicable, maintain existing protection standards
4086 to provide buffers around springs, seeps, and stream headwaters during development
4087 actions, such as forest management and road building activities. Maintain and protect
4088 groundwater recharge areas and cold water refugia. Use open-bottomed culverts or
4089 bridges when building roads or upgrading culverts to allow fish and wildlife passage. In dry
4090 ecoregions, use cooperative incentive programs to fence spring heads, which provides
4091 benefits to wildlife but allows water to be available for other uses. Minimize impacts from
4092 climate change.

4093 **SPRING-FED STREAMS**

4094 **Ecoregions**

4095 **BM, EC, KM, WC, WV**

4096 Streams dominated by groundwater rather than surface runoff are characterized by more
4097 stable flow and thermal regimes. Spring-fed rivers often display relatively static
4098 morphology compared to runoff systems, and habitat complexity is provided by aquatic
4099 plants and large wood inputs. These factors, along with nutrient rich inputs from underlying
4100 geology, contribute to ecological productivity. These streams support cool-water species
4101 such as bull trout and provide refugia for other temperature-limited species. Spring-fed
4102 streams are also a critical resource for climate resiliency.

4103 **Conservation Actions**

4104 Identify and protect the state's cold-water resources. Conduct real-time flow and
4105 temperature monitoring in priority areas. Maintain and protect groundwater recharge
4106 areas, especially at higher elevations. Maintain, protect, and restore natural water flow
4107 regimes. Maintain and protect supporting aquifers. Minimize impacts from climate change.

4108 **WESTERN JUNIPER SAVANNA WITH MATURE TREES; LATE**
4109 **SUCCESSIONAL WESTERN JUNIPER WOODLANDS**

4110 **Ecoregions**

4111 **BM, CP, EC, NBR**

4112 Western juniper savannas consist of scattered, often large, juniper trees within shrub-
4113 steppe. Late successional juniper woodlands may have a higher density of trees but are
4114 characterized by large-diameter trees. These juniper habitats are important for songbirds
4115 and raptors. In the Columbia Plateau ecoregion, the remaining Ferruginous Hawk nest
4116 sites are primarily juniper trees.

4117 A small percentage of Oregon's juniper woodlands are considered late successional. A
4118 high percentage of old-growth juniper in Central Oregon near Bend, Redmond, and Madras
4119 has been lost. Remaining stands are highly fragmented and are threatened by encroaching
4120 small junipers. In contrast, recruitment of juniper in the sandy shrub-steppe of the
4121 Columbia Plateau is naturally poor, so young juniper trees are not replacing older ones lost
4122 to cutting or natural death.

4123 **Conservation Actions**

4124 Remove small diameter encroaching juniper trees while maintaining larger diameter
4125 junipers and connectivity of juniper patches. Reintroduce fire where practical. Collect
4126 better spatial data on the distribution of mature juniper savanna. In the Columbia Plateau,
4127 maintain existing large juniper trees and examine factors affecting tree recruitment.
4128 Research is underway to determine the age, composition, structure, and wildlife usage of
4129 old growth juniper woodlands (for more information, see the **Eastern Oregon Agricultural**
4130 **Research Center website**).

4131 WESTERN LARCH FOREST AND WOODLAND

4132 **Ecoregions**

4133 **BM, EC**

4134 Western larch (*Larix occidentalis*) forests and woodlands occur on cool, moist sites
4135 interspersed with ponderosa pine habitats. These habitats may have been much more
4136 common historically in the Blue Mountains ecoregion.

4137 **Conservation Actions**

4138 Maintain large-diameter larch trees and patches of larch forest to provide local diversity.
4139 Control key invasive plants.

4140