

HATCHERY AND GENETICS MANAGEMENT PLAN (HGMP)

Hatchery Program:	Clackamas Winter Steelhead
Species or Hatchery Stock:	Winter Steelhead (Stock 122)
Agency/Operator:	Oregon Department of Fish and Wildlife
Watershed and Region:	Clackamas, North Willamette
Date Submitted:	October 25, 2001 February 8, 2017 December 20, 2024
Date Last Updated:	December 20, 2024

Section 1. GENERAL PROGRAM DESCRIPTION

1.1. Name of hatchery or program.

Clackamas winter steelhead program (Stock 122)

1.2. Species and population (or stock) under propagation, and ESA status.

Clackamas winter steelhead (*Oncorhynchus mykiss*)
Lower Columbia River Steelhead ESU – Threatened

1.3. Responsible organization and individuals

Lead Contact:

Name (and title): Scott Patterson, Fish Propagation Program Manager

Agency or Tribe: Oregon Department of Fish and Wildlife

Address: 4034 Fairview Industrial Dr. SE, Salem, OR 97302-1142

Telephone: 503-947-6218

Fax: 503-947-6202

Email: Scott.D.Patterson@odfw.oregon.gov

Operation Staff Leads:

Name (and title): Ben Walczak, District Fish Biologist

Agency or Tribe: Oregon Department of Fish and Wildlife

Address: 17330 SE Evelyn St., Clackamas, OR 7023

Telephone: 971-673-6013

Fax: 971-673-6070

Email: Ben.Walczak@odfw.oregon.gov

Name (and title): Daniel Straw, Clackamas Hatchery Manager

Agency or Tribe: Oregon Department of Fish and Wildlife

Address: 24500 S Entrance Rd., Estacada, OR 97023

Telephone: 503-630-7210

Fax: 503-630-4566

Email: Daniel.E.Straw@odfw.oregon.gov

1.4. Funding source, staffing level, and annual hatchery program operational costs.

The facilities utilized for this program are funded at various levels by the Mitchell Act, State of Oregon, BPA, PGE, City of Portland, Sport Fish Restoration Act, and USACE.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

1.5. Location(s) of hatchery and associated facilities.

Clackamas Hatchery – Clackamas River, RM 22.6 (45.2961, -122.3603)
Willamette Watershed, Oregon

Bonneville Hatchery – Tanner Creek, RM 0.25 (45.6333, -121.9556)
Lower Columbia Watershed, Oregon

Oak Springs Hatchery – Deschutes River, RM 47 (45.2255, 121.0917)
Deschutes Watershed, Oregon

Eagle Creek Hatchery – Eagle Creek, RM 9.94 (45.2762, -122.2017)
Willamette Watershed, Oregon

North Fork Dam – Clackamas River, RM 30 (45.2435, -122.2803)
Willamette Watershed, Oregon

Foster Acclimation Pond – Clackamas River, RM 11 (45.3838, -122.4476)
Willamette Watershed, Oregon

Eagle Fern Acclimation Pond – Eagle Creek, RM, 6.6 (45.2319, -122.2874)
Willamette Watershed, Oregon

1.6. Type of program.

Integrated Harvest

1.7. Purpose (Goal) of program.

Augmentation - The goal of this program is to increase recreational harvest opportunities by releasing artificially propagated smolts.

1.8. Justification for the program.

The program releases fin marked juvenile winter steelhead to produce hatchery adults for harvest.

1.9. List of program “Performance Standards”.

Please refer to sections 1.10.1 and 1.10.2 below.

1.10. List of program “Performance Indicators”, designated by "benefits" and "risks."

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

1.10.1. “Performance Indicators” addressing benefits.

Table 1.10.1. Performance Indicators addressing benefits.

Performance Standard	Performance Indicator	Monitoring and Evaluation
Program provides harvest opportunity	Increased harvest	Estimate annual harvest
Contribution of fish carcasses to the ecosystem	Developed plan for carcass outplants	Record carcass outplants
Satisfy legal harvest while eliminating impacts on wild populations	Developed harvest management plan	Adhere to FMEP
Achieve within-hatchery performance standards	Achievement of IHOT standards	Adhere to IHOT standards

1.10.2. “Performance Indicators” addressing risks.

Table 1.10.2. Performance Indicators addressing risks.

Performance Standard	Performance Indicator	Monitoring and Evaluation
Assess detrimental genetic impacts	Stray rates	Estimate pHOS and record coded-wire tag recoveries
Potentially unpredictable egg supply	Egg take	Record broodstock metrics
Not achieving within-hatchery performance standards	Achievement of IHOT standards	Adhere to IHOT standards
Avoid disease transfer	Application of fish health standards	Adhere to fish health standards and policies

1.11. Expected size of program.

1.11.1. Proposed annual broodstock collection level (maximum number of adult fish).

Up to 150 pairs for production plus additional as needed to support shortfalls in other programs

1.11.2. Proposed annual fish release levels (maximum number) by life stage and location.

Table 1.11.2. Proposed annual fish release levels

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Life Stage	Release Location	Annual Release Level
Yearling	Clackamas Basin	265,000 ¹
¹ The program may exceed the release goal by 5% annually (278,250) and 2% over any 5-year period (270,300) to account for variation during rearing.		

1.12. Current program performance, including estimated smolt-to-adult survival rates, adult production levels, and escapement levels. Indicate the source of these data.

Table 1.12. Smolt to adult survival (SAR) rates, estimated numbers produced, and escapement.

Brood Year	Smolt-to-Adult Survival (%)	Estimated Adult Production	Escapement
2009	2.05	3,634	1,291
2010	2.84	5,093	1,910
2011	3.09	3,648	1,305
2012	2.40	3,298	561
2013	5.54	9,819	3,589
2014	1.06	1,356	389
2015	0.73	1,561	609
2016	0.64	2,085	804
2017	0.71	1,599	562
2018	0.55	1,073	445
2019	1.20	2,656	1,207
2020	0.45	1,394	283

Data Source: HMS and NF Dam Counts

1.13. Date program started (years in operation), or is expected to start.

Started in 1991

1.14. Expected duration of program.

Indefinite

1.15. Watersheds targeted by program.

Willamette

1.16. Indicate alternative actions considered for attaining program goals, and reasons why those actions are not being proposed.

Alternative 1 – Increase program size (Not preferred)

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Increased releases could potentially increase fisheries, financial commitment, and impacts to listed species. This alternative is not preferred due to the increased cost and negative impact to listed species.

Alternative 2 – Reduce program size (Not preferred)

Decreased releases could potentially decrease fisheries, financial commitment, and impacts to listed species. This alternative is not preferred due to the negative impact to fisheries.

Section 2. PROGRAM EFFECTS ON NMFS ESA-LISTED SALMONID POPULATIONS. (USFWS ESA-LISTED SALMONID SPECIES AND NON-SALMONID SPECIES ARE ADDRESSED IN ADDENDUM A)

2.1. List all ESA permits or authorizations in hand for the hatchery program.

This program was evaluated under the Mitchell Act Biological Opinion signed on 1/15/2017. NMFS issued an ESA 4(d) Rule Limit 5 Take Exemption in response to a revised HGMP on 1/17/2018. This 2024 version is intended as re-submission for reinitiation of consultation.

The North Fork Dam ladder and sorting facility has an independent Section 10 authorization for incidental take during operation.

2.2. Provide descriptions, status, and projected take actions and levels for NMFS ESA-listed natural populations in the target area.

2.2.1. Description of NMFS ESA-listed salmonid population(s) affected by the program.

- Identify the NMFS ESA-listed population(s) that will be directly affected by the program.

Table 2.2.1.1 NMFS ESA-listed populations directly affected by the program.

Life Stage	Activity	Affected NMFS ESA-listed Population
Adult	Broodstock	Lower Columbia River Steelhead

- Identify the NMFS ESA-listed population(s) that may be incidentally affected by the program.

Table 2.2.1.2 NMFS ESA-listed populations incidentally affected by the program.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Life Stage	Activity	Affected NMFS ESA-listed Population
Adult	Trap Operations	Upper Willamette Chinook Salmon
		Lower Columbia River Steelhead
		Lower Columbia River Coho Salmon
Juvenile	Ecological Competition	Columbia River Chum Salmon
		Lower Columbia Chinook Salmon
		Lower Columbia River Coho Salmon
		Lower Columbia River Steelhead
		Middle Columbia River Steelhead
		Upper Columbia River Spring-run Chinook
		Upper Columbia River Steelhead
		Upper Willamette Chinook Salmon
		Upper Willamette Steelhead
		Snake River Spring/Summer-run Chinook Salmon
		Snake River Fall-run Chinook Salmon
		Snake River Sockeye Salmon
		Snake River Basin Steelhead

2.2.2. Status of NMFS ESA-listed salmonid population(s) affected by the program.

Due to overlapping broodstock collection seasons at Clackamas Hatchery, the independent authorization for take at North Fork Dam, and the low likelihood that listed populations will be affected through ecological competition, only assessments of listed populations affected annually by the Clackamas Hatchery Trap and angler caught winter steelhead broodstock program are presented here.

- Describe the status of the listed natural population(s) relative to “critical” and “viable” population thresholds

Table 2.2.2.1 Description of listed population thresholds.

Affected NMFS ESA-listed Population	Status
Upper Willamette Chinook Salmon	Moderate to very high extinction risk.
Lower Columbia River Steelhead	Moderate extinction risk.
Lower Columbia River Coho Salmon	Moderate extinction risk.

- Provide the most recent 12 year (e.g. 1988-present) progeny-to-parent ratios, survival data by life-stage, or other measures of productivity for the listed population. Indicate the source of these data.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Productivity assessments are described in the Lower Columbia River Conservation and Recovery Plan for Oregon Populations of salmon and steelhead (ODFW 2010).

- Provide the most recent 12 year (e.g. 1988-1999) annual spawning abundance estimates, or any other abundance information. Indicate the source of these data.

Table 2.2.2.2. Abundance estimates of listed natural origin Clackamas spring Chinook, fall Chinook, winter steelhead, and Coho.

Year	Spring Chinook	Fall Chinook	Winter Steelhead	Coho
2012	2,217	60	2,733	1,663
2013	2,740	360	2,427	4,012
2014	1,231	127	3,404	10,672
2015	3,041	192	3,740	1,784
2016	4,346	711	4,144	1,628
2017	4,534	34	2,531	7,598
2018	2,864	673	3,030	3,159
2019	2,823	880	1,500	4,044
2020	5,068	111	2,005	9,012
2021	3,540	36	990	10,572
2022	4,194	43	1,566	13,796
2023	4,698	37	1,277	19,916

Data Source: ODFW Salmon and Steelhead Recovery Tracker.

- Provide the most recent 12-year (e.g. 1988-1999) estimates of annual proportions of direct hatchery-origin and listed natural-origin fish on natural spawning grounds, if known.

Table 2.2.2.3. Annual estimates of pHOS for Clackamas spring Chinook, fall Chinook, winter steelhead, and Coho.

Year	Spring Chinook	Fall Chinook	Winter Steelhead	Coho
2012	5.5%	81.3%	2.0%	8.3%
2013	4.6%	7.6%	10.1%	2.8%
2014	2.4%	30.6%	0.5%	14.3%
2015	3.8%	37.7%	7.1%	11.4%
2016	2.6%	21.9%	8.1%	8.9%
2017	1.5%	62.2%	1.1%	11.9%
2018	3.4%	5.1%	11.6%	9.5%
2019	3.4%	5.2%	11.9%	4.6%

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

2020	3.4%	0.0%	7.8%	N/A
2021	3.4%	0.1%	8.4%	1.7%
2022	3.4%	0.0%	8.4%	4.3%
2023	2.0%	0.0%	3.1%	2.6%

Data Source: ODFW Salmon and Steelhead Recovery Tracker.

2.2.3. Describe hatchery activities, including associated monitoring and evaluation and research programs, that may lead to the take of NMFS listed fish in the target area, and provide estimated annual levels of take.

- Describe hatchery activities that may lead to the take of listed salmonid populations in the target area, including how, where, and when the takes may occur, the risk potential for their occurrence, and the likely effects of the take.

Listed steelhead will be taken for incorporation into broodstock. Broodstock collection also has potential to take listed steelhead, Chinook, and Coho through migration delay, capture, handling, and release during trap operation.

- Provide information regarding past takes associated with the hatchery program, (if known) including numbers taken, and observed injury or mortality levels for listed fish.

Table 2.2.3.1. Number of listed natural origin coho, steelhead, and spring Chinook salmon captured associated with all programs at Clackamas Hatchery, including angler caught broodstock from Jan. 1 – Dec. 31.

Calendar Year	Coho		Winter Steelhead		Spring Chinook	
	Capture	Mortality	Capture	Mortality	Capture	Mortality
2015	0	0	0	0	4	0
2016	0	0	0	0	1	0
2017	0	0	0	0	0	0
2018	0	0	46	5	0	0
2019	0	0	52	1	13	0
2020	0	0	49	0	10	0
2021	0	0	41	0	5	0
2022	13	0	49	3	15	0
2023	1	0	47	0	0	0
2024	0	0	45	1	0	0

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Data Source: HMS and program records.

- Provide projected annual take levels for listed fish by life stage (juvenile and adult) quantified (to the extent feasible) by the type of take resulting from the hatchery program.

The values below represent all take for Clackamas Hatchery and the angler caught winter steelhead broodstock program from Jan. 1 – Dec. 31 due to overlapping broodstock collection seasons.

Table 2.2.3.2. Estimated annual take levels of listed natural origin Coho.

Listed Species: <u>Coho</u>	ESU: <u>Lower Columbia River</u>		Activity: <u>Trapping</u>	
Location: <u>Clackamas Hatchery</u>	Dates: <u>Jan 1 – Dec 31</u>		Operator: <u>ODFW</u>	
Type of Take	Annual Take of Listed Fish by Life Stage			
	Egg/Fry	Juvenile/Smolt	Adult	Carcass
Observe or harass				
Collect for transport				
Capture, handle, and release				
Capture, handle, tag/mark/tissue sample, and release			100	
Removal (e.g. broodstock)				
Intentional lethal take				
Unintentional lethal take			≤ 3	
Other Take (specify)				

Table 2.2.3.3. Estimated annual take levels of listed natural origin steelhead.

Listed Species: <u>Steelhead</u>	ESU: <u>Lower Columbia</u>		Activity: <u>Trapping/Angling</u>	
Location: <u>Clackamas Hatchery/Willamette Basin</u>	Dates: <u>Jan 1 – Dec 31</u>		Operator: <u>ODFW</u>	
Type of Take	Annual Take of Listed Fish by Life Stage			
	Egg/Fry	Juvenile/Smolt	Adult	Carcass
Observe or harass				
Collect for transport				
Capture, handle, and release				
Capture, handle, tag/mark/tissue sample, and release			200	
Removal (e.g. broodstock)			49	
Intentional lethal take				

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Unintentional lethal take			≤ 6	
Other Take (specify)				

Table 2.2.3.4. Estimated annual take levels of listed natural origin Chinook.

Listed Species: <u>Chinook</u>	ESU: <u>Upper Willamette</u>		Activity: <u>Trapping</u>	
Location: <u>Clackamas Hatchery</u>	Dates: <u>Jan 1 – Dec 31</u>		Operator: <u>ODFW</u>	
Type of Take	Annual Take of Listed Fish by Life Stage			
	Egg/Fry	Juvenile/Smolt	Adult	Carcass
Observe or harass				
Collect for transport				
Capture, handle, and release				
Capture, handle, tag/mark/tissue sample, and release			350	
Removal (e.g. broodstock)			120	
Intentional lethal take				
Unintentional lethal take			≤ 11	
Other Take (specify)				

- Indicate contingency plans for addressing situations where take levels within a given year have exceeded, or are projected to exceed, take levels described in this plan for the program.

Facility operations and fish handling procedures will be modified immediately if listed fish mortality related to operation is identified in or near the trap. This may include, but is not limited to, additional staff training or review of proper procedures, trap modifications, cessation of trapping, modified operation by hatchery personnel, etc.

Section 3. RELATIONSHIP OF PROGRAM TO OTHER MANAGEMENT OBJECTIVES

3.1. Describe alignment of the hatchery program with any ESU-wide hatchery plan (e.g. Hood Canal Summer Chum Conservation Initiative) or other regionally accepted policies. Explain any proposed deviations from the plan or policies.

The hatchery program will be operated consistent with the Lower Columbia River Conservation and Recovery Plan for Oregon Populations of Salmon and Steelhead, ODFW’s Native Fish Conservation Policy, and ODFW’s Hatchery Management Policy.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

3.2. List all existing cooperative agreements, memoranda of understanding, memoranda of agreement, or other management plans or court orders under which program operates.

N/A

3.3. Relationship to harvest objectives.

Juvenile hatchery steelhead releases by this program are mass marked to facilitate selective harvest of hatchery fish.

3.3.1. Describe fisheries benefitting from the program, and indicate harvest levels and rates for program-origin fish for the last twelve years (1988-99), if available. *Also provide estimated future harvest rates on fish propagated by the program, and on listed fish that may be taken while harvesting program fish.*

This program supports fisheries in the Clackamas River, Lower Willamette, and Lower Columbia River with a harvest level described in Table 1.12.

3.4. Relationship to habitat protection and recovery strategies.

Assumption regarding habitat conditions related to this program are that freshwater habitat is at capacity, production is not limited in other habitat for different life stages, and artificially produced populations can coexist without jeopardizing the fitness of natural populations. Considering these assumptions, this program rears juveniles from egg to smolt in a hatchery setting to avoid impacting freshwater habitat availability. Those juveniles are then released at optimized times to emigrate and utilized more available habitat, ultimately minimizing impacts to habitat.

3.5. Ecological interactions. [Please review Addendum A before completing this section. If it is necessary to complete Addendum A, then limit this section to NMFS jurisdictional species. Otherwise complete this section as is.]

Listed fish have potential to both negatively impact and be negatively impacted by the program through competition for resources and attraction of predators. Listed fish in the area may positively impact and be positively impacted by the program through increased nutrient cycling and the subsequent increase in production potential of surrounding habitats.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Section 4. WATER SOURCE

4.1. Provide a quantitative and narrative description of the water source (spring, well, surface), water quality profile, and natural limitations to production attributable to the water source.

Clackamas Hatchery – 50 cfs are authorized from the Clackamas River with an additional 370 gpm of ground water. Operation and discharge are authorized under a NPDES permit.

Eagle Creek Hatchery – 34.02 cfs are authorized from Eagle Creek and spring water. Operation and discharge are authorized under a NPDES permit.

Bonneville Hatchery – 52.22 cfs are authorized from Tanner Creek and ground water. Operation and discharge are authorized under a NPDES permit.

Oak Springs Hatchery – 67.5 cfs are authorized from various springs with an additional 3.23 acre-feet of storage. Operation and discharge are authorized under a NPDES permit.

4.2. Indicate risk aversion measures that will be applied to minimize the likelihood for the take of listed natural fish as a result of hatchery water withdrawal, screening, or effluent discharge.

Hatcheries adhere to water right and NPDES permits.

Section 5. FACILITIES

5.1. Broodstock collection facilities (or methods).

Broodstock are collected from the North Fork Dam and Clackamas Hatchery adult traps. At both locations, adults enter fish ladders leading to temporary holding ponds. The program also incorporates angler caught broodstock.

5.2. Fish transportation equipment (description of pen, tank truck, or container used).

Juvenile and adult transportation is performed by liberation trucks or portable liberation tanks. Liberation trucks are typically 1,000–2,500-gallon capacity units, either mounted on a large flatbed or tanker style truck. The trucks are equipped with oxygen diffusing systems, water re-circulation pumps, and dissolved oxygen meters. Portable liberation tanks have a capacity of 300 gallons and are equipped with oxygen diffusion systems. The transfer of juvenile fish on station is performed using a distribution box, irrigation

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

pipe and a gas-powered water pump. Eggs and milt are transported via passenger vehicles in small, covered containers.

5.3. Broodstock holding and spawning facilities.

Broodstock are held at Clackamas Hatchery in a concrete raceway with multiple dividers. Spawning is conducted under a portable canvas canopy.

5.4. Incubation facilities.

Eggs are incubated at Clackamas, Oak Springs, and Bonneville hatcheries inside incubation rooms via troughs and vertical stack incubators. Each trough has a capacity of 100,000 eggs while each vertical stack incubator has a capacity of 10,000 eggs. Flow through the trough incubators is 3-6 gpm.

5.5. Rearing facilities.

Fry are reared at Oak Springs, Bonneville, and Eagle Creek hatcheries in circular tanks and raceways. Juvenile steelhead reared in concrete raceways.

5.6. Acclimation/release facilities.

Acclimation and release facilities are currently located at Clackamas Hatchery, Foster acclimation facility, Eagle Fern acclimation facility, and Eagle Creek Hatchery. Fish at Clackamas and Eagle Creek hatcheries utilize asphalt and concrete rearing ponds while the Foster and Eagle Fern acclimation facilities utilize above ground rearing tanks.

5.7. Describe operational difficulties or disasters that led to significant fish mortality.

Significant fish mortality could occur due to human error, disease outbreaks, intake failure, high stream flows, drought, high temperatures, low temperatures, wildfire, or various other natural disasters.

5.8. Indicate available back-up systems, and risk aversion measures that will be applied, that minimize the likelihood for the take of listed natural fish that may result from equipment failure, water loss, flooding, disease transmission, or other events that could lead to injury or mortality.

Hatchery-origin winter steelhead propagated under this program are ESA-listed. Back-up generators, automatic alarm and notification systems, UV light treatment, and 24/7 on-call staff are all utilized to avoid and minimize impacts from operational failures at the hatchery. Hatchery and fish health staff continuously work to minimize disease transmission.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Section 6. BROODSTOCK ORIGIN AND IDENTITY

Describe the origin and identity of broodstock used in the program, its ESA-listing status, annual collection goals, and relationship to wild fish of the same species/population.

6.1. Source.

Broodstock for Clackamas hatchery winter steelhead (stock-122) are sourced from listed natural adults returning to North Fork Dam.

6.2. Supporting information.

6.2.1. History.

The Clackamas winter steelhead program first began in 1991 utilizing listed natural adults returning to North Fork Dam.

6.2.2. Annual size.

Up to 150 pairs for production plus additional as needed to support shortfalls in other programs

6.2.3. Past and proposed level of natural fish in broodstock.

Up to 49 natural origin adults will be taken for broodstock. This program utilized 100% natural origin broodstock from 1991 – 1997 and then 70% until 2012. There was a temporary pause in integration of natural origin adults from 2012 – 2014 due to litigation.

6.2.4. Genetic or ecological differences.

Clackamas hatchery winter steelhead are not likely to exhibit differences from the naturally produced Lower Columbia River steelhead due to the incorporation of natural origin broodstock.

6.2.5. Reasons for choosing.

Integration of local, natural origin winter steelhead into the hatchery population will preserve existing hatchery-wild genetic relationships, while boosting diversity within the hatchery population. Integration of wild fish into the hatchery brood will serve to mitigate genetic risk from the program and may improve the performance.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

6.3. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects on listed natural fish that may occur as a result of broodstock selection practices.

The program will be managed as an integrated population, receiving gene flow from the wild population on a regular basis. This management approach will decrease the rate of genetic divergence between the wild and hatchery populations, effectively mitigating for genetic risk.

Section 7. BROODSTOCK COLLECTION

7.1. Life-history stage to be collected (adults, eggs, or juveniles).

Adults

7.2. Collection or sampling design.

The traps described above typically target winter steelhead throughout the run. Broodstock may also be sourced via hook-and-line by anglers.

7.3. Identity.

Adipose fin clips are used to identify returning Clackamas hatchery winter steelhead. Natural origin winter steelhead are identified by the presence of a fully intact and developed adipose fin. Other marks may be considered in the future if necessary.

7.4. Proposed number to be collected:

7.4.1. Program goal (assuming 1:1 sex ratio for adults)

Up to 150 pairs for production plus additional as needed to support shortfalls in other programs. Up to 49 natural origin adults will be taken for broodstock.

7.4.2. Broodstock collection levels for the last twelve years (e.g. 1988-99), or for most recent years available:

Table 7.4.2. Broodstock collection levels.

Brood Year	Adults		Pond Loss
	Females	Males	
2012	77	87	5
2013	72	72	5
2014	72	72	0
2015	124	124	21
2016	120	120	0

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Brood Year	Adults		Pond Loss
	Females	Males	
2017	95	95	0
2018	115	115	5
2019	132	132	1
2020	110	110	3
2021	106	106	0
2022	96	96	3
2023	115	115	0
2024	100	100	17

Data source: HMS

7.5. Disposition of hatchery-origin fish collected in surplus of broodstock needs.

Surplus hatchery fish are distributed to the tribes, donated to foodbanks, recycled to the lower river for additional fishing opportunities, provided for educational purposes, distributed for stream enrichment, or disposed of.

7.6. Fish transportation and holding methods.

Adult winter steelhead collected for broodstock are transported to Clackamas hatchery where they are held until spawning.

7.7. Describe fish health maintenance and sanitation procedures applied.

The fish health monitoring plan is based on the Integrated Hatchery Operations Team for the Columbia Basin Anadromous Salmonid Hatcheries (see Policies and Procedures for the Columbia Basin Anadromous Salmonid Hatcheries, Annual Report 1994. Bonneville Power Administration) and the ODFW Fish Health Management Policy (OAR 635-007-0960 to 635-007-0995). Other resources are used to guide the program and management of diseases such as American Fisheries Society Fish Health Section Blue Book and the World Organisation for Animal Health Manual of Diagnostic tests for Aquatic Animals.

7.8. Disposition of carcasses.

Carcasses are to be used for stream enrichment or disposed of in accordance with ODFW policies and procedures, which include freezing, rendering, or burying.

7.9. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects on listed natural fish resulting from the broodstock collection program.

The risks of disease in broodstock will be minimized by protocols described above in section 7.7. Broodstock will be collected from the entire run period to maintain genetic

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

diversity within the hatchery-produced population. Natural origin males used for broodstock will be released after spawning.

Section 8. MATING

Describe fish mating procedures that will be used, including those applied to meet performance indicators identified previously.

8.1. Selection method.

Adults are collected randomly from throughout the temporal distribution of the run to avoid any timing or size bias.

8.2. Males.

Males are generally only used once during spawning unless there is a shortage of males. No residual males are utilized for spawning.

8.3. Fertilization.

This program intends to utilize 1:1 male-to-female ratios in 3x3 matrices, although alternative matrices will be utilized to maximize genetic diversity and the incorporation of natural genetics.

8.4. Cryopreserved gametes.

Cryopreservation is not utilized for this program.

8.5. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects on listed natural fish resulting from the mating scheme.

Factorial mating schemes and random broodstock selection will be used to reduce the risk of loss of within-population genetic diversity.

Section 9. INCUBATION AND REARING

Specify any management *goals* (e.g. “egg to smolt survival”) that the hatchery is currently operating under for the hatchery stock in the appropriate sections below. Provide data on the success of meeting the desired hatchery goals.

9.1. Incubation:

9.1.1. Number of eggs taken and survival rates to eye-up and/or ponding.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Table 9.1.1. Eggs Taken and Survival Rates

Brood Year	Egg Take	Percent Survival to Eye-up
2011	1,585,552	95.0
2012	1,511,372	97.6
2013	1,259,708	93.7
2014 ¹	1,193,611	92.4
2015	1,450,703	90.0
2016	1,661,557	89.0
2017	614,922	92.2
2018	221,965	94.5
2019	261,694	90.5
2020 ¹	871,830	92.9
2021	611,388	84.1
2022	1,339,058	90.1
2023	581,986	88.6

¹ Inventory error. Egg take was underestimated.

Data Source: HMS

9.1.2. Cause for, and disposition of surplus egg takes.

Surplus eggs are typically collected to compensate for egg to smolt mortality and IHNV culling. If necessary, eggs are culled according to ODFW and IHOT guidelines.

9.1.3. Loading densities applied during incubation.

The average size of steelhead eggs is about 152 - 204 eggs per ounce. Water flows at 3-6 gpm through incubators. Each vertical incubating tray is loaded with up to 10,000 eggs.

9.1.4. Incubation conditions.

The water temperature and flow are monitored daily with a range of 34° – 65°F at 3-6 gpm. Dissolved oxygen (DO) ranges between 9 and 10 parts per million (ppm) at Clackamas Hatchery.

9.1.5. Ponding.

Fry are manually relocated once 100% buttoned up, at approximately 920 – 940 temperature units and approximately 2,700 fish per pound.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

9.1.6. Fish health maintenance and monitoring.

The fish health monitoring plan is based on the Integrated Hatchery Operations Team for the Columbia Basin Anadromous Salmonid Hatcheries (see Policies and Procedures for the Columbia Basin Anadromous Salmonid Hatcheries, Annual Report 1994. Bonneville Power Administration) and the ODFW Fish Health Management Policy (OAR 635-007-0960 to 635-007-0995). Other resources are used to guide the program and management of diseases such as American Fisheries Society Fish Health Section Blue Book and the World Organisation for Animal Health Manual of Diagnostic tests for Aquatic Animals.

9.1.7. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects on listed fish during incubation.

Eggs are incubated with water that meets IHOT water quality standards. Water supplies and the power supply are alarmed to notify hatchery personnel if a failure occurs and hooked up to a back-up generator, in case of a power failure. Hatchery staff are available 24 hr/day for immediate response to any emergency.

9.2. Rearing:

9.2.1. Provide survival rate data (*average program performance*) by hatchery life stage (fry to fingerling; fingerling to smolt) for the most recent twelve years, or for years dependable data are available.

Table 9.2.1 Rearing survival rates.

Brood Year	Number of Fry Poned	Number of Juveniles at Marking	Percent Survival to Marking
2011	299,157	184,032	61.5
2012	188,689	127,626	67.6
2013	189,166	88,617	46.8
2014	299,381	174,688	58.3
2015	371,416	222,296	59.9
2016	439,154	253,777	57.8
2017	419,971	209,920	50.0
2018	350,268	172,068	49.1
2019	359,786	167,499	46.6
2020	417,920	226,619	54.2

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

2021	455,159	290,624	63.9
2022	336,364	238,564	70.9
2023	412,171	308,961	75.0

Data source: HMS

9.2.2. Density and loading criteria (goals and actual levels).

The goal is to maintain less than 1.0 lb of fish per cubic foot of water during rearing.

9.2.3. Fish rearing conditions

Water temperatures range from 40° – 58°F. Dissolved oxygen is only monitored when rearing densities are elevated.

9.2.4. Indicate biweekly or monthly fish growth information (*average program performance*), including length, weight, and condition factor data collected during rearing, if available.

Table 9.2.4. Average end-of-month size

Month	Size (fpp)
1	2193.1
2	1665.7
3	793.6
4	254.4
5	115
6	66.2
7	42.7
8	31.1
9	21.5
10	15.7
11	11.6
12	8.5
13	7.7

Data source: HMS

9.2.5. Indicate monthly fish growth rate and energy reserve data (*average program performance*), if available.

See Table 9.2.4 above for fish growth. No energy reserve data is available.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

9.2.6. Indicate food type used, daily application schedule, feeding rate range (e.g. % B.W./day and lbs/gpm inflow), and estimates of total food conversion efficiency during rearing (*average program performance*).

Fish are fed according to the feed manufacturer's guidance and ODFW's growth program tool.

9.2.7. Fish health monitoring, disease treatment, and sanitation procedures.

The fish health monitoring plan is based on the Integrated Hatchery Operations Team for the Columbia Basin Anadromous Salmonid Hatcheries (see Policies and Procedures for the Columbia Basin Anadromous Salmonid Hatcheries, Annual Report 1994. Bonneville Power Administration) and the ODFW Fish Health Management Policy (OAR 635-007-0960 to 635-007-0995). Other resources are used to guide the program and management of diseases such as American Fisheries Society Fish Health Section Blue Book and the World Organisation for Animal Health Manual of Diagnostic tests for Aquatic Animals.

9.2.8. Smolt development indices (e.g. gill ATPase activity), if applicable.

Weight samples of the fish are taken monthly to ensure proper growth rate.

9.2.9. Indicate the use of "natural" rearing methods as applied in the program.

"Natural" rearing methods are not utilized by this program.

9.2.10. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects on listed fish under propagation.

Rearing ponds are cleaned on weekly basis or as necessary through visual observation of solid wastes on pond bottom.

Section 10. RELEASE

Describe fish release levels, and release practices applied through the hatchery program.

10.1. Proposed fish release levels.

Table 10.1. Proposed fish release levels.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Age Class	Maximum Number	Size (fpp)	Release Date	Location
Yearling	265,000 ¹	6	March - April	Clackamas Hatchery, Foster Acclimation Pond, Eagle Fern Acclimation
¹ The program may exceed the release goal by 5% annually (278,250) and 2% over any 5-year period (270,300) to account for variation during rearing.				

10.2. Specific location(s) of proposed release(s).

Stream, river, or watercourse: Clackamas River
Release point: Clackamas Hatchery
Major watershed: Clackamas
Basin or Region: Willamette

Stream, river, or watercourse: Eagle Creek
Release point: Eagle Fern Acclimation
Major watershed: Clackamas
Basin or Region: Willamette

Stream, river, or watercourse: Clackamas River
Release point: Foster Acclimation
Major watershed: Clackamas
Basin or Region: Willamette

10.3. Actual numbers and sizes of fish released by age class through the program.

Table 10.3. Program releases.

Brood Year	Number of Yearling Released	Average Size (fpp)
2012	172,583	6.0
2013	157,276	5.7
2014	130,674	6.0
2015	222,545	7.5
2016	279,475	6.9
2017	176,350	6.4
2018	233,035	8.6
2019	207,608	7.9
2020	286,372	9.1
2021	280,888	12.4

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Brood Year	Number of Yearling Released	Average Size (fpp)
2022	224,690	11.5
2023	213,692	9.3
Average	215,432	8.1

Data source: HMS

10.4. Actual dates of release and description of release protocols.

Table 10.4. Annual release dates.

Brood Year	Release Date Range
2012	4/23 – 5/01
2013	4/25 – 4/29
2014	4/09 – 4/29
2015	4/13 – 4/20
2016	4/05 – 5/01
2017	4/03 – 4/27
2018	3/11 – 4/23
2019	3/19 – 4/21
2020	3/19 – 4/21
2021	3/18 – 4/06
2022	3/24 – 4/17
2023	4/19 – 4/22

Data Source: HMS

10.5. Fish transportation procedures, if applicable.

All juveniles are transported in liberations trucks described above.

10.6. Acclimation procedures (methods applied and length of time).

Juveniles are acclimated for up to 12 weeks at Clackamas Hatchery, Eagle creek Hatchery, Foster Acclimation, and Eagle Fern Acclimation prior to release.

10.7. Marks applied, and proportions of the total hatchery population marked, to identify hatchery adults.

The program goal is to mass mark 100% winter steelhead smolts with an adipose fin clip and majority of the juveniles retain their marks.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

10.8. Disposition plans for fish identified at the time of release as surplus to programmed or approved levels.

Under current policy, surplus juveniles are destroyed or marked and released into a closed water system such as a lake, reservoir or pond where they contribute to angling opportunities. In general, OAR 635-007-0545 directs disposition of surplus hatchery juveniles.

10.9. Fish health certification procedures applied pre-release.

The fish health monitoring plan is based on the Integrated Hatchery Operations Team for the Columbia Basin Anadromous Salmonid Hatcheries (see Policies and Procedures for the Columbia Basin Anadromous Salmonid Hatcheries, Annual Report 1994. Bonneville Power Administration) and the ODFW Fish Health Management Policy (OAR 635-007-0960 to 635-007-0995). Other resources are used to guide the program and management of diseases such as American Fisheries Society Fish Health Section Blue Book and the World Organisation for Animal Health Manual of Diagnostic tests for Aquatic Animals.

10.10. Emergency release procedures in response to flooding or water system failure.

Emergency releases may occur at any size or point in time during rearing in response water system failure, disease, drought, wildfire, flood, rising air/water temperatures, or any other adverse environmental conditions that may pose a threat to hatchery staff and/or hatchery fish.

Emergency releases will occur after the hatchery crew has exhausted all possibilities for retaining the fish and consulted with the ODFW District Biologist. Emergency releases will be limited to the Willamette basin, or into a closed water body per OAR 635-007-0545.

10.11. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects on listed fish resulting from fish releases.

Juveniles are generally acclimated and released downstream of North Fork Dam at sufficient sizes to minimize potential interactions during emigration and adult homing.

Section 11. MONITORING AND EVALUATION OF PERFORMANCE INDICATORS

11.1. Monitoring and evaluation of “Performance Indicators” presented in Section 1.10.

11.1.1. Describe plans and methods proposed to collect data necessary to respond to each “Performance Indicator” identified for the program.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Table 11.1.1. Methods for evaluating program indicators.

Performance Indicator M&E	Methods
Estimate annual harvest	Query harvest card and ELS data
Record carcass outplants	Keep hatchery records of adult dispositions.
Adhere to FMPEP	N/A
Adhere to IHOT standards	N/A
Estimate pHOS and record coded-wire tag recoveries	Query OASIS spawning ground survey results.
Record broodstock metrics	Keep hatchery records of broodstock numbers and egg take.
Adhere to fish health standards and policies	N/A

11.1.2. Indicate whether funding, staffing, and other support logistics are available or committed to allow implementation of the monitoring and evaluation program.

Current funding is sufficient to implement the activities identified in 11.1.1. Any additional monitoring or evaluations would require additional funding/staffing.

11.2. Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects on listed fish resulting from monitoring and evaluation activities.

Listed fish will not be affected by monitoring or evaluation.

Section 12. RESEARCH

This program does not currently have a research nexus.

Section 13. ATTACHMENTS AND CITATIONS

AFS-FHS (American Fisheries Society-Fish Health Section). 2014. FHS blue book: suggested procedures for the detection and identification of certain finfish and shellfish pathogens, 2020 edition. Accessible at: <https://units.fisheries.org/fhs/fish-health-section-blue-book-2020/>

IHOT (Integrated Hatchery Operations Team). 1996. Operation Plans for Anadromous Fish Production Facilities in the Columbia River Basin. Volume II-Oregon. Annual Report 1995. Portland, OR. Project Number 92-043, Contract Number DE-BJ79-91BP60629.

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

- NMFS. 2017. Endangered Species Act Section 7(a)(2) Biological Opinion and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat (EFH) Consultation. NOAA's National Marine Fisheries Service's implementation of the Mitchell Act Final Environmental Impact Statement preferred alternative and administration of Mitchell Act hatchery funding. January 15, 2017. NMFS Consultation No.: WCR-2014-697. 535p.
- ODFW. 2010a. Final Lower Columbia River Conservation and Recovery Plan for Oregon Populations of Salmon and Steelhead. August 6, 2010. 437p.
- Oregon Administrative Rules (OAR 635-007- -0542 through -0548). 2003. Fish Health Management Policy. Oregon Department of Fish and Wildlife, Salem, OR.
- Oregon Administrative Rules (OAR 635-007-0960 through 1000). 2003. Fish Health Management Policy. Oregon Department of Fish and Wildlife, Salem, OR.
- Oregon Administrative Rules (OAR 635-007-050 2 through -0509). 2002. Native Fish Conservation Policy. Oregon Department of Fish and Wildlife, Salem, OR.
- World Organisation for Animal Health (OIE). (2024). Manual of Diagnostic Tests for Aquatic Animals, eleventh editions 2024. Available at: <https://www.woah.org/en/what-we-do/standards/codes-and-manuals/aquatic-manual-online-access/>

This program shall be implemented according to the Incidental Take Statement, Terms and Conditions, and Hatchery Operation Framework of the NMFS Mitchell Act Biological Opinion. Any deviation from the biological opinion must be approved by NMFS.

Section 14. CERTIFICATION LANGUAGE AND SIGNATURE OF RESPONSIBLE PARTY

“I hereby certify that the information provided is complete, true and correct to the best of my knowledge and belief. I understand that the information provided in this HGMP is submitted for the purpose of receiving limits from take prohibitions specified under the Endangered Species Act of 1973 (16 U.S.C.1531-1543) and regulations promulgated thereafter for the proposed hatchery program, and that any false statement may subject me to the criminal penalties of 18 U.S.C. 1001, or penalties provided under the Endangered Species Act of 1973.”

Name, Title, and Signature of Applicant:

Certified by _____ Date: _____

Attachment 1. Age class designations by fish size and species for salmonids released from hatchery facilities.
(generally from Washington Department of Fish and Wildlife, November 1999)

	Species/Age Class	Size Criteria	
		Number of fish/pound	Grams/fish
X	Chinook Yearling	<=20	>=23
X	Chinook (Zero) Fingerling	>20 to 150	3 to <23
X	Chinook Fry	>150 to 900	0.5 to <3
X	Chinook Unfed Fry	>900	<0.5
X	Coho Yearling ¹	<20	>=23
X	Coho Fingerling	>20 to 200	2.3 to <23
X	Coho Fry	>200 to 900	0.5 to <2.3
X	Coho Unfed Fry	>900	<0.5
X	Chum Fed Fry	<=1000	>=0.45
X	Chum Unfed Fry	>1000	<0.45
X	Sockeye Yearling ²	<=20	>=23
X	Sockeye Fingerling	>20 to 800	0.6 to <23
X	Sockeye Fall Releases	<150	>2.9
X	Sockeye Fry	> 800 to 1500	0.3 to <0.6
X	Sockeye Unfed Fry	>1500	<0.3
X	Pink Fed Fry	<=1000	>=0.45
X	Pink Unfed Fry	>1000	<0.45
X	Steelhead Smolt	<=10	>=45
X	Steelhead Yearling	<=20	>=23
X	Steelhead Fingerling	>20 to 150	3 to <23
X	Steelhead Fry	>150	<3
X	Cutthroat Trout Yearling	<=20	>=23
X	Cutthroat Trout Fingerling	>20 to 150	3 to <23
X	Cutthroat Trout Fry	>150	<3
X	Trout Legals	<=10	>=45
X	Trout Fry	>10	<45

¹ Coho yearlings defined as meeting size criteria and 1 year old at release, and released prior to June 1st.

² Sockeye yearlings defined as meeting size criteria and 1 year old.